

KPV

Alpha-MSH Derived Tripeptide | 5mg & 10mg x 10 Vials

Anti-Inflammatory | GI Protection | Wound Healing | SubQ / Oral

Purity: 99.8% | Endotoxin-Free | Research Use Only

KPV (Lys-Pro-Val) is the C-terminal tripeptide of alpha-MSH, retaining full anti-inflammatory activity of the parent peptide without melanocortin side effects. It acts through melanocortin receptors and direct anti-inflammatory pathways in epithelial and immune cells. Particularly effective for inflammatory bowel disease (IBD), skin inflammation, and wound healing research.

KPV Vial Comparison




5mg
5.0mg/mL



10mg
10.0mg/mL

Parameter	Specification
Class	Alpha-MSH C-terminal tripeptide -- anti-inflammatory MC receptor modulator
Vial Sizes	5mg x 10 10mg x 10 Purity 99.8%
Reconstitution	1 mL BAC Water: 5mg->5mg/mL 10mg->10mg/mL
Standard Dose	500-3,000 mcg per injection SubQ oral also effective for GI
Routes	SubQ (systemic) Oral solution (GI-specific anti-inflammatory)
Frequency	1-2x daily no fasting required
Cycle & Storage	4-12 weeks on, 2-4 off 2-8 deg C use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
MC Receptor Anti-Inflammatory	Binds melanocortin receptors (MC1R, MC3R, MC5R) on immune and epithelial cells; inhibits NF-kB signalling; reduces pro-inflammatory cytokines (IL-6, TNF-alpha, IL-1beta); downregulates inflammatory cascade	IBD; skin inflammation; systemic anti-inflammatory
GI Epithelial Protection	Direct anti-inflammatory action in intestinal epithelial cells; reduces neutrophil infiltration; promotes tight junction integrity; particularly effective in colitis models; oral administration targets GI directly	IBD; leaky gut; colitis research; GI repair
Wound Healing & Skin	Promotes keratinocyte migration; reduces inflammatory damage to healing wounds; anti-inflammatory at wound site without immunosuppression; synergistic with BPC-157 in tissue repair stacks	Wound healing; skin inflammation; psoriasis research

Benefit	Context
IBD Research	KPV is one of the most researched peptides for inflammatory bowel disease -- oral administration provides direct GI anti-inflammatory delivery.
No Melanocortin Side Effects	Unlike full alpha-MSH, KPV retains anti-inflammatory activity without tanning, appetite, or sexual side effects.
Combo Synergy	KPV is included in the BPC+TB+GHKCu+KPV (80mg) combo vial -- adding anti-inflammatory reinforcement to the tissue repair stack.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: Vol(mL)=Dose(mcg)/Conc(mcg/mL) | **Units:** Vol x 100


5mg x 10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)

Dose(mcg)	500mcg	1000mcg	2000mcg	3000mcg
Vol(mL)	0.1000	0.2000	0.4000	0.6000
Units	10.0U	20.0U	40.0U	60.0U
Doses/Vial	10	5	2	1


10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	500mcg	1000mcg	2000mcg	3000mcg
Vol(mL)	0.0500	0.1000	0.2000	0.3000
Units	5.0U	10.0U	20.0U	30.0U
Doses/Vial	20	10	5	3

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Select route	SubQ (systemic) or oral solution (GI-specific). Both effective.
2	Draw/prepare	SubQ: draw volume per dose table. Oral: dilute in water.
3	Administer	SubQ: abdomen 45 deg, rotate sites. Oral: drink solution on empty stomach.
4	Rotate & dispose	8-point rotation for SubQ. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Systemic (SubQ)	500-2,000 mcg	1-2x daily	4-12 wk	No fasting. Any time.
GI-Specific (Oral)	1,000-3,000 mcg	1-2x daily (empty stomach)	4-12 wk	Dissolve in water. Empty stomach for GI effect.

Side Effect	Likelihood	Management
Injection site reaction	Common	Mild. Rotate sites.
Mild GI effects (oral)	Occasional	Rare given anti-inflammatory mechanism.

Contraindications: Cancer history, pregnancy, under 18.

1. 5mg: 1mL BAC->5mg/mL | 10mg: 1mL->10mg/mL.

