

LIPO-C + B12

HIGH-DOSE B12 | 121 mg/mL | 10mL VIAL

MIC + L-Carnitine + Cyanocobalamin + B5 + B6


Pre-Mixed Solution | Ready to Use | Research Use Only

Lipo-C + B12 is a pre-mixed, ready-to-use liquid solution. No reconstitution required — draw directly from the vial. This formulation delivers a significantly higher B12 (Cyanocobalamin) concentration (121 mg/mL total formula) compared to standard Lipo-C, making it ideal for B12-deficiency correction alongside lipotropic support.

Parameter	Specification
Product Name	Lipo-C + B12 (High-Dose B12 Lipotropic Cocktail)
Concentration	121 mg/mL total active formula per mL of solution
Vial Volume	10 mL per vial
Total Active per Vial	1,210 mg total active compounds per vial (121 mg/mL x 10 mL)
Format	Pre-mixed sterile solution — NO reconstitution required
Typical Dose	0.5 mL to 1.0 mL per injection (60.5 mg to 121 mg per dose)
Doses per Vial	10 doses at 1.0 mL 20 doses at 0.5 mL
Route	Intramuscular (IM) preferred Subcutaneous (SubQ) alternative
Frequency	1-3x per week (most common) Daily protocol also researched
Appearance	Clear to light pink/red tint — normal due to B12 (cyanocobalamin); discard if cloudy or contains particles
Storage	2-8 deg C refrigerated Protect from light Do not freeze

ACTIVE INGREDIENTS & MECHANISM OF ACTION

Lipo-C + B12 combines seven synergistic compounds — the standard MIC lipotropic blend plus L-Carnitine and B vitamins — with a significantly elevated B12 (Cyanocobalamin) concentration. This makes it particularly effective for individuals with B12 deficiency, fatigue, or neurological support needs, in addition to lipotropic fat metabolism support.



Ingredient	Role	Key Mechanism	Benefit
B12 (Cyanocobalamin) ★ High-Dose	Energy coenzyme Neurological vitamin	Cofactor in methylmalonyl-CoA mutase; myelin sheath synthesis; red blood cell maturation; homocysteine → methionine conversion; DNA synthesis	Energy production, neurological health, anaemia correction, mood regulation, methylation support
Methionine	Essential sulphur amino acid	Precursor to SAM-e; drives hepatic methylation; promotes liver fat export as VLDL; homocysteine regulation	Liver detox, fat mobilisation, antioxidant support
Inositol	B-vitamin family carbocyclic polyol	Regulates insulin receptor signalling; component of phospholipid membranes; reduces hepatic fat accumulation	Insulin sensitisation, liver fat reduction, mood support
Choline	Essential nutrient	Required for VLDL synthesis and hepatic fat export; precursor to acetylcholine; phosphatidylcholine production	Prevents fatty liver, cognitive function, neurotransmitter synthesis
L-Carnitine	Amino acid derivative	Transports long-chain fatty acids across inner mitochondrial membrane for beta-oxidation	Enhanced fat oxidation, energy production, exercise performance
B5 (Dexpanthenol)	Provitamin B5	Converted to Coenzyme A; essential for fatty acid synthesis and citric acid cycle	Energy metabolism, fat burning, adrenal support
B6 (Pyridoxine)	Water-soluble vitamin	Cofactor in 100+ enzymatic reactions; amino acid metabolism; neurotransmitter synthesis (serotonin, dopamine)	Protein metabolism, mood, immune function, PMS symptom relief

DOSE GUIDE & INJECTION PROCEDURE

No reconstitution needed. Draw directly from vial using a clean syringe. Warm to room temperature 15-20 minutes before injecting. The solution may appear faintly pink or red — this is normal due to B12.



Draw Lipo-C+B12 directly -- no reconstitution required

Protocol	Volume	Active per Dose	Frequency	Route
Standard	1.0 mL	121 mg	3x / week	IM preferred
Half Dose	0.5 mL	60.5 mg	3x / week	IM or SubQ
Maintenance	1.0 mL	121 mg	1-2x / week	IM or SubQ
Daily (Research)	0.5 mL	60.5 mg	5 on / 2 off	SubQ — rotate daily
B12 Repletion (high-deficit)	1.0 mL	121 mg	Daily x 7 days then 1x/week	IM preferred for fastest repletion

Step-by-Step Injection Procedure


Step	Action	Detail
1	Warm the vial	Remove from fridge 15-20 minutes before use. Cold B12 solution causes unnecessary stinging and discomfort at injection site.
2	Wash hands	Wash with soap and warm water for 20 seconds. Dry completely.
3	Select needle & syringe	IM: 23-25G x 1 to 1.5 inch syringe. SubQ: 29-31G x 5/16 inch insulin syringe. 1mL volume syringe for both routes.
4	Clean vial stopper	Wipe rubber stopper with a fresh alcohol swab. Allow 10 seconds to dry before inserting needle.
5	Draw the dose	Invert vial. Insert needle. Draw 0.5 or 1.0 mL as prescribed. Tap syringe and expel all air bubbles.
6	Clean injection site	IM: wipe deltoid or outer thigh with swab, let dry 10-15 sec. SubQ: wipe abdomen 2 inches from navel.
7	Inject	IM: 90-degree dart motion into deltoid; aspirate not required for deltoid/thigh. SubQ: pinch skin, 45-degree angle, release, inject slowly.
8	Withdraw and apply pressure	Remove needle. Press site with dry swab 5-10 seconds. Do NOT rub. Mild warmth, redness, or B12-coloured tint is normal.
9	Dispose safely	Place needle immediately into sharps container. Never recap, reuse, or leave exposed.

INJECTION SITES, ROTATION & SCHEDULE

IM injection into the deltoid (upper arm) is preferred for fastest absorption and best B12 uptake. **SubQ** into the abdomen is a suitable alternative but absorption may be slightly slower. Rotate sites at every session.

Site	Route	Needle Size	Notes
Deltoid (upper outer arm)	IM (Preferred)	23-25G x 1-1.5 in	Fastest B12 absorption. Alternate left/right arm each session.
Vastus Lateralis (outer thigh)	IM (Alternative)	23-25G x 1-1.5 in	Good for self-injection. Use middle outer thigh only.
Abdomen (2 in from navel)	SubQ (Alternative)	29-31G x 5/16 in	Pinch skin. 45-degree angle. Slower absorption than IM.

Injection Sites -- IM Preferred | SubQ Alternative



IM (90 deg) vs SubQ (45 deg) Injection Angles



Recommended Schedule -- 3x Per Week (Mon / Wed / Fri)



Alternative: 1-2x per week. Inject mornings -- B12 boosts energy.

SIDE EFFECTS, STORAGE & GOLDEN RULES

Side Effect	Likelihood	Management
Burning / stinging at injection site	Very Common	Warm vial to room temp before injecting. Inject slowly. Do not rub site.
Pink/red discolouration at site (B12)	Common	Normal B12 pigment. Harmless. Fades within minutes to hours.
Redness or warmth at site	Common	Expected local reaction. Rotate sites. Resolves within 1-2 hours.
Increased energy / alertness	Very Common	Expected B12 + Carnitine effect. Inject morning or midday to avoid sleep disruption.
Nausea (especially higher doses)	Occasional	Inject with a light meal. Reduce to 0.5 mL if persistent.
Urinary odour change	Occasional	Normal sulphur by-product of methionine. Also, B12 may turn urine faintly yellow — harmless.
Acne flare (B12 sensitivity)	Occasional	High-dose B12 can trigger acne in susceptible individuals. Reduce to 0.5 mL or reduce frequency.
Allergic reaction	Rare	Stop immediately. Seek medical advice. Do not rechallenge without assessment.

STOP & SEEK MEDICAL ADVICE: Severe or spreading redness, heat, swelling at injection site (possible infection)

STOP & SEEK MEDICAL ADVICE: Difficulty breathing or swallowing (anaphylaxis)

STOP & SEEK MEDICAL ADVICE: Severe chest pain, palpitations, or irregular heartbeat

STOP & SEEK MEDICAL ADVICE: Numbness or tingling in hands/feet (B6 peripheral neuropathy at high doses)

STOP & SEEK MEDICAL ADVICE: Jaundice or dark urine (hepatic concern)

Contraindications: Known allergy to cobalamin (B12) or any formula component, Leber hereditary optic neuropathy (cyanocobalamin contraindicated), active liver or kidney disease, trimethylaminuria, pregnancy or lactation. Not for use under 18.

Storage Guidelines

State	Temperature	Duration	Notes
Unopened vial	2-8 deg C	Per expiry	Keep upright. Protect from light — B12 is light-sensitive. Do not freeze.
Opened/in-use vial	2-8 deg C	28 days	Cap securely. Store away from light. Inspect before each draw.
Short-term transport	Below 25 deg C	Max 72 hr	Insulated bag with ice packs. Wrap in foil to protect from light.
Do not freeze	No freezing	N/A	Freezing precipitates B12 and disrupts the formula. Discard if frozen.

Golden Rules — Lipo-C + B12

1. Warm the vial to room temperature 15-20 minutes before injecting -- cold B12 stings.
2. Draw directly from the vial -- NO reconstitution is required for this product.

3. A faint pink/red tint to the solution is normal -- this is the B12 (cyanocobalamin).
4. IM injection into the deltoid is preferred for fastest and most complete B12 absorption.
5. Inject slowly over 5-10 seconds to minimise localised burning.
6. Do not rub the site after injection. Press gently with a dry swab for 5-10 seconds.
7. Inject in the morning or midday -- B12 and carnitine can disrupt sleep if injected late.
8. Rotate injection sites every session. Allow at least 1 week rest per site.
9. B12 is light-sensitive -- store vial in original packaging away from direct light.
10. Store at 2-8 deg C. Never freeze. Opened vial must be used within 28 days.

DISCLAIMER: This document is produced by _____ for informational and research reference purposes only. This guide does not constitute medical advice. Lipo-C + B12 must not be used for self-administration or therapeutic purposes without qualified medical supervision. Consult a licensed healthcare professional before beginning any injection protocol. _____ assumes no liability for misuse of this product.

| _____ .com | Research Use Only | For Educational Purposes