


MOTS-C (Human)

Mitochondrial-Derived Peptide | 10mg & 40mg x 10 Vials

AMPK Activation | Metabolic Regulation | Exercise Mimetic


Purity: 99.8% | Endotoxin-Free | Research Use Only

MOTS-C is a mitochondrial-derived peptide (MDP) encoded by the 12S rRNA gene. It is a powerful AMPK activator and metabolic regulator -- called an exercise mimetic for its ability to activate energy-sensing pathways independently of exercise. Doses can range from 1mg to 5mg per injection.



Parameter	Specification
Class	Mitochondrial-derived peptide (MDP) -- AMPK activator
Vial Sizes	10mg x 10 40mg x 10 Purity 99.8%
Reconstitution	10mg: 1mL BAC->10mg/mL 40mg: 2mL BAC->20mg/mL
Standard Dose	1,000-5,000 mcg (1-5mg) per injection no fasting required
Frequency	1x daily or 3-5x per week SubQ morning preferred
Cycle & Storage	8-12 weeks on, 4 weeks off 2-8 deg C 28 days reconstituted

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
AMPK Activation	Activates AMP-activated protein kinase; cellular energy sensor; inhibits anabolic pathways; activates fat-burning; mimics calorie restriction and exercise signalling	Metabolic improvement; fat oxidation; glucose regulation
Folate Cycle	Regulates folate-methionine cycle; affects one-carbon metabolism	Metabolic regulation; longevity research
Anti-Aging & Exercise	Endogenous MOTS-C declines with age; supplementation restores youthful metabolic signalling and physical performance in animal models	Anti-aging research; exercise mimetic; longevity

Benefit	Context
Exercise Mimetic	MOTS-C activates AMPK and metabolic pathways of physical exercise without exercise itself.
Insulin Sensitivity	Improves glucose uptake and insulin sensitivity in multiple research models.
Anti-Aging	Endogenous MOTS-C declines with age; supplementation restores metabolic function in animal models.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$ | **Units:** $\text{Vol} \times 100$


10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	1000mcg	2000mcg	3000mcg	4000mcg	5000mcg
Vol(mL)	0.1000	0.2000	0.3000	0.4000	0.5000
Units	10.0U	20.0U	30.0U	40.0U	50.0U
Doses/Vial	10	5	3	2	2


40mg x 10 -- 2mL BAC -> 20.0mg/mL (20,000mcg/mL)

Dose(mcg)	1000mcg	2000mcg	3000mcg	4000mcg	5000mcg
Vol(mL)	0.0500	0.1000	0.1500	0.2000	0.2500
Units	5.0U	10.0U	15.0U	20.0U	25.0U
Doses/Vial	20	10	6	5	4

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Wash & prepare	Wash hands. 29-31G insulin syringe. Draw exact dose volume.
2	Inject SubQ	Abdomen preferred. Pinch 2-3cm. 45 degrees. Inject slowly.
3	Rotate & dispose	Record site. 8-point rotation. Press 5 sec. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Standard	1,000-2,000 mcg	3-5x/week	8-12wk	Morning injection preferred.
Research	3,000-5,000 mcg	Daily	8-12wk	Higher dose. Monitor response.

Side Effect	Likelihood	Management
Injection site reaction	Common	Rotate sites. Fresh needle each injection.
Fatigue (initial)	Occasional	Metabolic adaptation. Resolves within 1-2 weeks.

Contraindications: Cancer history, pregnancy, under 18.

1. 10mg vial: 1 mL BAC Water yields 10 mg/mL (10,000 mcg/mL).
2. 40mg vial: 2 mL BAC Water yields 20 mg/mL (20,000 mcg/mL).
3. No fasting required -- MOTS-C acts independently of the GH/insulin axis.

4. Inject 1x daily or 3-5x per week SubQ. Morning timing is preferred.
5. Cycle 8-12 weeks on, followed by a mandatory 4-week off-cycle rest period.
6. Store at 2-8 deg C refrigerated. Use reconstituted vial within 28 days.
7. Rotate injection sites through all 8 abdomen points with every injection.
8. Contraindicated in active cancer history -- AMPK and folate cycle modulation.
9. Not for use during pregnancy, lactation, or in individuals under 18 years.

DISCLAIMER: research use only. Not for human therapeutic use. Consult a qualified professional. assumes no liability.
| .com | Research Use Only