

Melanotan II

Cyclic alpha-MSH Analogue | 10mg x 10 Vials

Melanogenesis | Libido Research | Appetite Modulation | SubQ

Purity: 99.8% | Endotoxin-Free | Research Use Only

Melanotan II (MT-II) is a cyclic synthetic analogue of alpha-MSH with broader melanocortin receptor activity (MC1R, MC3R, MC4R). In addition to stimulating melanogenesis (via MC1R), MT-II activates MC4R in the hypothalamus -- producing the well-documented effects on libido/sexual function and appetite suppression. MT-II produces faster and more intense tanning than Melanotan I but with a more complex side-effect profile.


Melanotan II Vial Comparison



10mg
10.0mg/mL

Parameter	Specification
Class	Cyclic alpha-MSH analogue -- MC1R, MC3R, MC4R agonist
Vial Size	10mg x 10 vials Purity 99.8%
Reconstitution	1 mL BAC Water -> 10mg/mL (10,000 mcg/mL)
Standard Dose	250-1,000 mcg per injection SubQ start at 250 mcg
Frequency	Every 1-3 days SubQ no fasting required
vs Melanotan I	Cyclic; broader receptor activity (MC3R/MC4R); stronger tanning; libido/appetite effects
Cycle & Storage	4-12 weeks 2-8 deg C use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
MC1R Melanogenesis	Binds MC1R on melanocytes; activates cAMP; upregulates tyrosinase; stimulates eumelanin synthesis; produces UV-independent skin darkening faster and more intensely than Melanotan I	Rapid melanogenesis; UV-independent tanning research; photoprotection
MC4R Libido/Appetite	MC4R activation in hypothalamus; dopaminergic pathway stimulation; penile/clitoral erection (erectile function research); reduced appetite via MC4R satiety signalling; unique research applications beyond tanning	Sexual function research; erectile dysfunction models; appetite/obesity research
MC3R Metabolic	MC3R activation in arcuate nucleus; involved in energy homeostasis; synergistic with MC4R for appetite and metabolic effects; additional cardiovascular research applications	Energy metabolism; cardiovascular; complementary to MC4R effects

Benefit	Context
Faster Tanning	MT-II produces more rapid and intense melanogenesis than Melanotan I due to its cyclic structure and higher receptor affinity.
Multi-Receptor Research	MC3R/MC4R activity enables research into sexual function, appetite suppression, and energy homeostasis beyond melanogenesis.
Important Distinction	MT-II has a broader side-effect profile than MT-I. Nausea and libido effects are common and must be disclosed in research protocols.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.




1mL Insulin Syringe | 29-31G | SubQ

Formula: $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$ | **Units:** $\text{Vol} \times 100$

10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	250mcg	500mcg	750mcg	1000mcg
Vol(mL)	0.0250	0.0500	0.0750	0.1000
Units	2.5U	5.0U	7.5U	10.0U
Doses/Vial	40	20	13	10

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Wash & prepare	Wash hands. 29-31G syringe. No fasting.
2	Draw dose	Draw exact volume. Start at 250 mcg.
3	Inject SubQ	Abdomen preferred. 45 deg. Rotate sites.
4	Monitor & dispose	Monitor for nausea 20-30 min post-injection. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Loading (Tanning)	250-500 mcg	Daily x 5-7 days	Then reduce	Build melanin. Start 250mcg for first dose.
Maintenance	500-1,000 mcg	Every 2-3 days	4-12 wk	Maintain pigmentation.

Side Effect	Likelihood	Management
Nausea (dose-related)	Very Common	Start at 250 mcg. Take antihistamine 30 min prior.
Facial flushing	Common	Vasodilation. Transient 30-60 min.
Spontaneous erection	Common (males)	MC4R effect. Dose-related. Reduce if unwanted.
Appetite suppression	Common	MC4R satiety signal. Monitor food intake.
Mole changes	Monitor	MANDATORY dermatological monitoring throughout cycle.

Contraindications: Cancer history, pregnancy, under 18.

1. 1 mL BAC Water -> 10mg/mL. ALWAYS start at 250 mcg first injection.
2. No fasting required. Inject any time of day.
3. Nausea is very common. Consider antihistamine 30 min before first injections.
4. MC4R effects (libido, appetite) are expected pharmacological responses.
5. MANDATORY: Regular dermatological mole monitoring throughout the cycle.
6. Contraindicated in personal or family history of melanoma or dysplastic nevi.
7. Rotate SubQ sites through 8-point abdomen map.
8. Store 2-8 deg C. Use within 28 days. Never freeze reconstituted peptide.
9. Do not use with photosensitising medications without physician guidance.
10. Not for use during pregnancy, lactation, or in individuals under 18.

DISCLAIMER:

research use only. Not for human therapeutic use. Consult a qualified professional.

assumes no liability.

|

.com | **Research Use Only**