

Oxytocin

Neuropeptide Hormone | 2mg / 5mg / 10mg x 10 Vials

Social Bonding | Anxiety | Reproductive Research | SubQ / IM / Intranasal


Purity: 99.8% | Endotoxin-Free | Research Use Only

Oxytocin is a hypothalamic nonapeptide released from the posterior pituitary. Central roles: social bonding, trust, anxiety reduction, uterine contraction, lactation. Research applications: autism, PTSD, social anxiety, reproductive medicine. All three vials use 1 mL BAC Water.



Parameter	Specification
Class	Endogenous nonapeptide hormone -- posterior pituitary
Vial Sizes	2mg / 5mg / 10mg x 10 1mL BAC Water each
Concentrations	2mg->2mg/mL 5mg->5mg/mL 10mg->10mg/mL
Standard Dose	500-5,000 mcg SubQ or IM flexible timing
Routes	SubQ IM Intranasal (dilute to 40 IU/mL)
Key Uses	Social bonding, anxiety, PTSD, autism, reproductive medicine research
Storage	2-8 deg C use within 28 days never freeze

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Research Benefit
Social Bonding	Binds OXTR in limbic system; activates reward circuits; increases trust, empathy, social recognition; reduces amygdala fear reactivity	Social bonding; autism; PTSD; social anxiety research
Anxiety & Stress	Inhibits HPA axis; reduces cortisol and amygdala activation; anxiolytic via GABAergic modulation; promotes affiliative behaviour	Anxiety; PTSD; fear extinction models
Reproductive & Uterine	Stimulates uterine smooth muscle contraction; promotes milk ejection reflex; critical for labour induction and lactation research	Labour/lactation research; reproductive medicine

Benefit	Context
Social Neuroscience	Primary neuropeptide for bonding, trust, and empathy -- extensive human clinical trial data.
Anxiety & PTSD	Reduces amygdala reactivity and HPA stress response -- applications in PTSD and social anxiety.
Multi-Route	SubQ, IM, or intranasal -- intranasal provides direct CNS delivery without systemic effects.


RECONSTITUTION & DOSE CALCULATION

All 3 vials use 1 mL BAC Water. Room temp. Swirl gently.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, syringe, swabs, sharps.
2	Clean & draw	Wipe stoppers. Draw 1mL BAC. Aim at glass wall.
3	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Label date. 2-8 deg C, 28 days.



1mL Insulin Syringe | 29-31G | SubQ

2mg x10 -- 1mL BAC -> 2.0mg/mL (2,000mcg/mL)

Dose(mcg)	500mcg	1,000mcg	5,000mcg
Vol(mL)	0.2500	0.5000	2.5000
Units	25.0U	50.0U	250.0U
Doses/Vial	4	2	0


5mg x10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)

Dose(mcg)	500mcg	1,000mcg	5,000mcg
Vol(mL)	0.1000	0.2000	1.0000
Units	10.0U	20.0U	100.0U
Doses/Vial	10	5	1


10mg x10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	500mcg	1,000mcg	5,000mcg
Vol(mL)	0.0500	0.1000	0.5000
Units	5.0U	10.0U	50.0U
Doses/Vial	20	10	2

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Select route	SubQ: 29-31G abdomen. IM: 23-25G deltoid/gluteal. Intranasal: dilute 40IU/mL saline.
2	Draw dose	Exact volume per dose table. No fasting required.
3	Administer	SubQ/IM standard technique. Intranasal: tilt head forward, spray, sniff gently.
4	Rotate & dispose	Rotate SubQ/IM sites. 8-point rotation. Sharps container.

Protocol	Dose	Frequency	Notes
SubQ/IM Research	1,000-5,000 mcg	As needed or 1-2x daily	No fasting. Any time.
Intranasal	40 IU/mL dilution	20-40 IU per admin.	Direct CNS delivery. 10 IU/spray.

Side Effect	Likelihood	Management
PREGNANCY -- UTEROTONIC	CRITICAL	NEVER use unsupervised in pregnancy. Potent labour inducer.
Headache / nausea	Occasional	Dose-related. Reduce dose.
Water retention	High dose	Antidiuretic at very high doses. Monitor.

Contraindications: Unsupervised use in pregnancy (uterotonic). High-dose cardiovascular caution.

1. 2mg: 1mL BAC->2mg/mL | 5mg: 1mL->5mg/mL | 10mg: 1mL->10mg/mL.

2. NEVER use in pregnancy outside supervised medical/research setting -- potent uterotonic.
3. No fasting required. Inject any time.
4. Intranasal: dilute to 40 IU/mL in saline. Use nasal spray device.
5. Rotate SubQ sites through 8-point abdomen map every injection.
6. Store 2-8 deg C. Use within 28 days. Protect from heat.
7. Contraindicated in cardiovascular disease at high doses.
8. Not for unsupervised use in pregnancy, lactation initiation, or under 18.

DISCLAIMER: research use only. Not for human therapeutic use. Consult a qualified professional.
| .com | **Research Use Only**