

# Selank

Anxiolytic Nootropic Peptide | 5mg & 10mg x 10 Vials

Anxiety | Cognition | BDNF Upregulation | SubQ / Intranasal


Purity: 99.8% | Endotoxin-Free | Research Use Only

Selank is a synthetic heptapeptide analogue of tuftsin (Thr-Lys-Pro-Arg-Pro-Gly-Pro) developed by the Institute of Molecular Genetics (Russian Academy of Sciences). It exhibits anxiolytic, nootropic, and immunomodulatory effects with a unique safety profile -- no dependence, sedation, or withdrawal. Also available as an intranasal formulation for direct CNS delivery.



| Parameter       | Specification                                                   |
|-----------------|-----------------------------------------------------------------|
| Class           | Synthetic heptapeptide anxiolytic/nootropic -- tuftsin analogue |
| Vial Sizes      | 5mg x 10   10mg x 10   Purity 99.8%                             |
| Reconstitution  | 1 mL BAC Water: 5mg->5mg/mL   10mg->10mg/mL                     |
| Standard Dose   | 250-1,000 mcg per injection SubQ   intranasal also effective    |
| Routes          | SubQ   Intranasal (dilute, 1-2 drops/nostril)   no fasting      |
| Frequency       | 1-3x daily   acute or ongoing protocols                         |
| Cycle & Storage | Ongoing (low dependence risk)   2-8 deg C   use within 28 days  |

MECHANISM OF ACTION & RESEARCH BENEFITS



| Pathway                        | Mechanism                                                                                                                                                                                                   | Benefit                                                       |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------|
| Anxiolytic (GABAergic)         | Modulates GABA-A receptor sensitivity; anxiolytic without sedation or dependence; reduces anxiety comparable to benzodiazepines in animal models but without tolerance; normalises HPA axis stress response | Anxiety; stress; PTSD; anxiolytic research without dependence |
| Cognitive & BDNF               | Upregulates BDNF (brain-derived neurotrophic factor); enhances memory consolidation; improves learning and attention; nootropic effects independent of anxiolytic mechanism                                 | Cognitive enhancement; memory; BDNF research                  |
| Immunomodulation & Enkephalins | Tufts-in-derived immunostimulant activity; modulates enkephalin metabolism (endogenous opioid regulation); mood-stabilising; antidepressant effects in research                                             | Immune support; mood regulation; endogenous opioid research   |

| Benefit           | Context                                                                                                                  |
|-------------------|--------------------------------------------------------------------------------------------------------------------------|
| No Dependence     | Selank does not cause tolerance, dependence, or withdrawal -- a key advantage over benzodiazepines for anxiety research. |
| BDNF Upregulation | Unique dual mechanism: anxiolytic plus nootropic via BDNF. Most anxiolytics impair cognition -- Selank improves it.      |
| Intranasal Option | Intranasal delivery provides rapid CNS-targeted effect for acute anxiety protocols.                                      |


## RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

| Step | Action           | Detail                                                                                       |
|------|------------------|----------------------------------------------------------------------------------------------|
| 1    | Gather           | Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.                                    |
| 2    | Clean stoppers   | Wipe both stoppers. Air-dry 10-15 sec.                                                       |
| 3    | Draw 1mL BAC     | Exactly 1 mL. Expel air bubbles.                                                             |
| 4    | Inject into vial | Aim at GLASS WALL. Slowly. Never onto powder.                                                |
| 5    | Swirl & label    | Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days. |



1mL Insulin Syringe | 29-31G | SubQ

**Formula:**  $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$  | **Units:**  $\text{Vol} \times 100$


**5mg x 10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)**

| Dose(mcg)  | 250mcg | 500mcg | 750mcg | 1000mcg |
|------------|--------|--------|--------|---------|
| Vol(mL)    | 0.0500 | 0.1000 | 0.1500 | 0.2000  |
| Units      | 5.0U   | 10.0U  | 15.0U  | 20.0U   |
| Doses/Vial | 20     | 10     | 6      | 5       |


**10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)**

| Dose(mcg)  | 250mcg | 500mcg | 750mcg | 1000mcg |
|------------|--------|--------|--------|---------|
| Vol(mL)    | 0.0250 | 0.0500 | 0.0750 | 0.1000  |
| Units      | 2.5U   | 5.0U   | 7.5U   | 10.0U   |
| Doses/Vial | 40     | 20     | 13     | 10      |

## INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



| Step | Action           | Detail                                                           |
|------|------------------|------------------------------------------------------------------|
| 1    | Select route     | SubQ (29-31G, abdomen) or intranasal (dilute, nasal spray).      |
| 2    | Draw/prepare     | SubQ: draw dose. Intranasal: dilute to 0.15% solution in saline. |
| 3    | Administer       | SubQ: 45 deg, pinch skin. Intranasal: tilt head, spray, sniff.   |
| 4    | Rotate & dispose | 8-point rotation for SubQ. Sharps container.                     |

| Protocol              | Dose          | Frequency  | Duration           | Notes                                         |
|-----------------------|---------------|------------|--------------------|-----------------------------------------------|
| Anxiety Protocol      | 250-500 mcg   | 1-3x daily | Ongoing (low risk) | No dependence or tolerance. Safe ongoing use. |
| Cognitive Enhancement | 500-1,000 mcg | 1-2x daily | 8-12 wk            | BDNF upregulation and memory consolidation.   |

| Side Effect                | Likelihood      | Management                              |
|----------------------------|-----------------|-----------------------------------------|
| Mild sedation at high dose | Rare            | GABA modulation. Reduce dose.           |
| Mood improvement           | Expected effect | Anxiolytic and mood-normalising effect. |

**Contraindications: Cancer history, pregnancy, under 18.**

1. 5mg: 1mL BAC->5mg/mL | 10mg: 1mL->10mg/mL.

|                                                                             |
|-----------------------------------------------------------------------------|
| 2. No fasting required. Inject any time -- morning and afternoon preferred. |
| 3. SubQ injection or intranasal delivery -- both effective.                 |
| 4. Intranasal: dilute to 0.15% (1.5mg/mL) in saline. 1-2 drops per nostril. |
| 5. No tolerance, dependence, or withdrawal -- safe for ongoing use.         |
| 6. Upregulates BDNF: cognitive enhancement alongside anxiolytic effects.    |
| 7. Rotate SubQ sites through 8-point abdomen map.                           |
| 8. Store 2-8 deg C. Use within 28 days.                                     |
| 9. Not for use during pregnancy, lactation, or in individuals under 18.     |

---

DISCLAIMER:

research use only. Not for human therapeutic use. Consult a qualified professional.

assumes no liability.

|

**.com | Research Use Only**