

Sermorelin

GHRH Fragment (1-29) | 5mg & 10mg x 10 Vials

Natural GHRH Sequence | Bedtime Injection | Anti-Aging & Body Composition

Purity: 99.8% | Endotoxin-Free | Research Use Only


Sermorelin is the natural N-terminal fragment (amino acids 1-29) of endogenous GHRH. As the biologically active portion of GHRH, it stimulates pituitary GH release with a very short half-life (~11-12 min). Bedtime injection is optimal to amplify the natural deep-sleep GH pulse. Frequently used in anti-aging protocols.

FASTED REQUIRED: Inject at least 1-2 hours after last meal. Insulin blunts GH release. Wait 30 min before eating post-injection.



Parameter	Specification
Half-Life	~11-12 minutes -- very short acting, multiple daily or bedtime only
Vial Sizes	5mg x 10 10mg x 10 Purity 99.8%
Reconstitution	1 mL BAC Water: 5mg->5mg/mL 10mg->10mg/mL
Standard Dose	200-300 mcg per injection (mcg, not mg)
Frequency	1-3x daily -- always fasted bedtime preferred
Best Protocol	Bedtime SubQ -- amplifies natural deep-sleep GH surge
Cycle	12 weeks on, 4 weeks off Long-term safe at standard doses
Storage	2-8 deg C Reconstituted: use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
GHRHR	Natural GHRH(1-29) sequence binds GHRH receptors physiologically; gentle, pulsatile GH release; no desensitisation at standard doses	Safe long-term use; physiological GH rhythm
Sleep Amplification	Bedtime injection amplifies natural slow-wave sleep GH surge; aligns with 10-30 min peak; recovery and cellular repair enhanced	Better sleep quality; overnight recovery; anti-aging
IGF-1 & Body Comp	Cumulative GH elevation raises IGF-1 over weeks; lean mass preservation; fat reduction especially visceral	Progressive body composition improvement

Benefit	Context
Natural Sequence	Sermorelin IS GHRH(1-29) -- the identical biologically active sequence. Most gentle and natural GH stimulation available.
Anti-Aging Profile	Lower receptor desensitisation risk vs synthetic GHRH analogues. Suitable for longer cycles in anti-aging research.
Sleep Enhancement	Bedtime dosing amplifies the natural growth hormone surge during slow-wave sleep, enhancing recovery and cellular repair.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: Vol(mL)=Dose(mcg)/Conc(mcg/mL) | **Units:** Vol x 100


5mg x 10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)

Dose(mcg)	100mcg	150mcg	200mcg	250mcg	300mcg
Vol(mL)	0.0200	0.0300	0.0400	0.0500	0.0600
Units	2.0U	3.0U	4.0U	5.0U	6.0U
Doses/Vial	50	33	25	20	16


10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	100mcg	150mcg	200mcg	250mcg	300mcg
Vol(mL)	0.0100	0.0150	0.0200	0.0250	0.0300
Units	1.0U	1.5U	2.0U	2.5U	3.0U
Doses/Vial	100	66	50	40	33

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Confirm fasted	1-2 hr after last meal. For bedtime: 1hr after dinner.
2	Draw dose	Draw volume per dose table. Expel air bubbles.
3	Inject SubQ	Abdomen preferred. 45 deg. Rotate sites.
4	Wait & sleep	Bedtime: go to sleep within 30 min. Morning: wait 30 min before eating.

Protocol	Dose	Frequency	Duration	Notes
Bedtime (Preferred)	200-300 mcg	1x daily pre-bed	12wk on 4wk off	Best protocol. Amplifies deep-sleep GH pulse.
AM + Bed	200 mcg each	2x daily fasted	12wk on 4wk off	Standard research protocol.

Side Effect	Likelihood	Management
Injection site reaction	Common	Rotate sites. Fresh needle.
Drowsiness (bedtime)	Common	Expected -- GH promotes deep sleep.
Water retention	Mild	Less than synthetic GHRH. Resolves.

Contraindications: Cancer history, pregnancy, under 18.

1. Always inject fasted (1-2hr after last meal). Insulin blocks GH release.
2. Bedtime injection is the preferred protocol -- amplifies natural deep-sleep GH surge.
3. 1 mL BAC Water per vial: 5mg->5mg/mL 10mg->10mg/mL.
4. Stack with Ipamorelin at same injection for amplified GH pulses.
5. Natural GHRH sequence -- gentle, suitable for longer anti-aging cycles.
6. Rotate injection sites. Never shake. Store 2-8 deg C, 28 days max.
7. Cycle 12 weeks on, 4 weeks off. Contraindicated in cancer history.

DISCLAIMER:

research use only. Not for human therapeutic use. Consult a qualified professional.

assumes no liability.

|

.com | **Research Use Only**