

Snap-8


SNAP-8 Octapeptide | 10mg x 10 Vials

Neuromuscular Junction Modulator | Anti-Wrinkle | SubQ / Intradermal

Purity: 99.8% | Endotoxin-Free | Research Use Only


Snap-8 (Acetyl Glutamyl Heptapeptide-3) is a synthetic octapeptide analogue of the N-terminal domain of synaptosomal-associated protein (SNAP-25). It competes with SNAP-25 for positions in the SNARE complex, moderating the release of acetylcholine at neuromuscular junctions. This relaxes facial muscle contractions, reducing the appearance of expression lines -- a peptide alternative in anti-aging research.

Snap-8 Vial Comparison



Parameter	Specification
Class	Synthetic octapeptide -- SNARE complex modulator, SNAP-25 analogue
Vial Size	10mg x 10 vials Purity 99.8%
Reconstitution	1 mL BAC Water -> 10mg/mL (10,000 mcg/mL)
Standard Dose	100-500 mcg per injection intradermal or SubQ
Route	Intradermal (preferred for facial lines) or SubQ (systemic)
Frequency	1-2x daily no fasting required
Cycle & Storage	4-8 weeks on, 2 weeks off 2-8 deg C use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
SNARE Modulation	Snap-8 competes with SNAP-25 for incorporation into the SNARE complex at neuromuscular junctions; partially inhibits formation of the SNARE fusion complex; reduces vesicle docking efficiency; moderates acetylcholine (ACh) exocytosis	Reduced neuromuscular transmission intensity; muscle relaxation at injection site
ACh Release Reduction	Lower ACh release -> reduced motor end-plate stimulation; decreased contraction amplitude in facial expression muscles; milder than botulinum toxin -- reversible and dose-titratable	Softened expression lines; anti-wrinkle; facial rejuvenation research
Collagen Stimulation	Secondary fibroblast activation via reduced mechanical stress from repeated muscle contractions; may upregulate collagen I/III synthesis in overlying dermis	Complementary skin rejuvenation alongside wrinkle-reduction mechanism

Benefit	Context
Anti-Wrinkle Mechanism	Snap-8 reduces the intensity of facial muscle contractions by competing with SNAP-25 in the SNARE complex -- a peptide alternative to botulinum toxin research.
Reversible & Titratable	Unlike botulinum toxin, Snap-8 effects are fully reversible and dose-dependent, making it safer for dose-finding research.
Intradermal Targeting	Intradermal injection near target muscles delivers highest local concentration, maximising neuromuscular junction modulation at the expression line site.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.




1mL Insulin Syringe | 29-31G | SubQ

Formula: $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$ | **Units:** $\text{Vol} \times 100$


10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	100mcg	200mcg	300mcg	400mcg	500mcg
Vol(mL)	0.0100	0.0200	0.0300	0.0400	0.0500
Units	1.0U	2.0U	3.0U	4.0U	5.0U
Doses/Vial	100	50	33	25	20

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Wash & prepare	Wash hands. Use 29-31G needle. No fasting needed.
2	Select route	Intradermal (facial lines): 30-32G needle, bevel up, shallow 15 deg angle. SubQ (systemic): 29-31G, 45 deg, pinch skin.
3	Draw & inject	Draw exact dose. Inject slowly. For intradermal: expect small bleb/wheal.
4	Rotate & dispose	Rotate sites. Do not massage intradermal site. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Facial Lines	100-300 mcg per site	1-2x daily	4-8 wk 2 off	Intradermal at target line.
Systemic	300-500 mcg	1-2x daily	4-8 wk 2 off	SubQ abdomen. General anti-aging research.

Side Effect	Likelihood	Management
Injection site redness	Common	Normal for intradermal. Resolves within hours.
Temporary muscle softening	Intended effect	Gradual onset over 2-4 weeks. Fully reversible.
Bruising (intradermal)	Occasional	Use fine needle (30-32G). Gentle technique.

Contraindications: Cancer history, pregnancy, under 18.

1. 1 mL BAC Water -> 10mg/mL (10,000 mcg/mL).

2. No fasting required. Inject any time of day.
3. Intradermal route preferred for facial expression line research.
4. Use 30-32G needle for intradermal delivery near target muscles.
5. Do not massage intradermal injection site after delivery.
6. Start at 100 mcg per site and titrate upward based on response.
7. Cycle 4-8 weeks on, 2 weeks off. Store 2-8 deg C, use within 28 days.
8. Effects are gradual (2-4 weeks) and fully reversible -- not permanent.
9. Not for use during pregnancy, lactation, or in individuals under 18.

DISCLAIMER:

research use only. Not for human therapeutic use. Consult a qualified professional.

assumes no liability.

|

.com | Research Use Only