


BPC+TB Combo

BPC-157 + TB-500 Pre-Mixed | 4 Formulations x 10 Vials

Gold Standard Tissue Repair Stack | Synergistic Healing | SubQ / IM


Purity: 99.8% | Endotoxin-Free | Research Use Only

Pre-mixed combination vials containing BPC-157 + TB-500 (and optionally GHK-Cu and KPV). BPC-157 and TB-500 work through complementary non-overlapping pathways for synergistic tissue repair. Available in 4 formulations: BPC5+TB5 (10mg), BPC10+TB10 (20mg), BPC10+TB10+GHKC50 (70mg), and BPC10+TB10+GHKC50+KPV10 (80mg).



Parameter	Specification
BPC-157 Component	GI cytoprotection, angiogenesis, tendon repair, NO pathway -- 5mg or 10mg
TB-500 Component	Actin regulation, cell migration, angiogenesis, anti-inflammatory -- 5mg or 10mg
GHK-Cu (70/80mg vials)	Collagen synthesis, wound healing, gene modulation -- 50mg per vial
KPV (80mg vial)	Anti-inflammatory MC-receptor tripeptide -- 10mg per vial
Reconstitution	10mg/1mL BAC 20mg/2mL BAC 70mg/1mL BAC 80mg/1mL BAC
Standard Dose	0.5-1.0mL per injection -- delivers full dose of all components
Frequency & Storage	2-3x/week no fasting 2-8 deg C use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
BPC-157 Pathway	GI mucosal protection; tendon fibroblast activation; eNOS/NO upregulation; VEGF expression; vascular and neural protection	GI repair; tendon healing; vascular protection
TB-500 Pathway	Actin sequestration enabling cell migration to injury; NF-kB anti-inflammatory; cardiomyocyte and oligodendrocyte progenitor activation; VEGF angiogenesis	Cell migration; angiogenesis; anti-inflammatory; cardiac/neural
GHK-Cu + KPV (70/80mg)	GHK-Cu: collagen I/III synthesis via TGF-beta; antioxidant; modulates 4,000+ genes. KPV: direct MC-receptor anti-inflammatory; reduces cytokine cascade	Collagen remodelling; potent anti-inflammatory reinforcement

Benefit	Context
Synergistic Repair	BPC-157 and TB-500 act through completely different, complementary pathways -- combined efficacy significantly exceeds either peptide alone.
Pre-Mixed Convenience	Single vial replaces separate reconstitution and dosing of 2-4 peptides. One injection delivers all components.
Scalable Formulations	4 formulations allow protocol customisation from basic BPC+TB to the comprehensive 4-peptide 80mg vial.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$ | **Units:** $\text{Vol} \times 100$


BPC5+TB5 10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	500mcg	1000mcg	2000mcg
Vol(mL)	0.0500	0.1000	0.2000
Units	5.0U	10.0U	20.0U
Doses/Vial	20	10	5


BPC10+TB10 20mg x 10 -- 2mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	500mcg	1000mcg	2000mcg
Vol(mL)	0.0500	0.1000	0.2000
Units	5.0U	10.0U	20.0U
Doses/Vial	20	10	5

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Reconstitute	Add correct BAC Water volume per spec. Swirl gently. Never shake.
2	Draw dose	0.5-1.0mL per injection for full component dose.
3	Inject SubQ/IM	Near injury (localised) or abdomen (systemic). 45 deg SubQ or 90 deg IM.
4	Rotate & dispose	8-point rotation. No fasting. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Acute Injury	1.0mL (full dose)	2-3x/week	4-8 wk	Near injury site. All components at once.
Maintenance	0.5mL (half dose)	2x/week	8-12 wk	Lower ongoing dose after acute phase.

Side Effect	Likelihood	Management
Injection site reaction	Common	BPC-157 and TB-500 are well tolerated. Rotate sites.
Mild fatigue / healing response	Occasional	Normal systemic repair response. Resolves.

Contraindications: Cancer history, pregnancy, under 18.

1. 10mg/1mL BAC | 20mg/2mL BAC | 70mg/1mL BAC | 80mg/1mL BAC.

2. No fasting required. Inject any time.

3. Standard dose: 0.5-1.0mL per injection delivers full dose of all included components.
4. Inject near target injury for localised effect, or abdomen for systemic benefit.
5. BPC-157 + TB-500 use complementary pathways -- synergy exceeds either alone.
6. GHK-Cu (70/80mg vials) adds collagen synthesis and gene-level tissue repair.
7. KPV (80mg vial) adds potent anti-inflammatory support to the stack.
8. Rotate through 8-point abdomen map. Store 2-8 deg C, 28 days max.
9. Contraindicated in cancer history. Not for pregnancy or under 18.

DISCLAIMER: SaiyanMed research use only. Not for human therapeutic use. Consult a qualified professional. SaiyanMed assumes no liability.

SaiyanMed | saiyanmed.com | Research Use Only