

CJC-1295 + IPA


GHRH + GHS Synergy Stack | 5mg+5mg & 10mg x 10 Vials

Pre-Mixed Stack | 3-10x GH Pulse Amplification

Purity: 99.8% | Endotoxin-Free | Research Use Only


Pre-mixed CJC-1295 (no DAC) + Ipamorelin combo vial. GHRH primes pituitary + GHS triggers release = 3-10x amplified GH pulse. Must be injected fasted. One injection delivers both peptides simultaneously.

FASTED REQUIRED: Inject at least 1-2 hours after last meal. Insulin blunts GH release. Wait 30 min before eating post-injection.



Parameter	Specification
Contents	CJC-1295 (no DAC) + Ipamorelin in equal parts per vial
Reconstitution	1 mL BAC Water -> 10 mg/mL total (5mg/mL each)
Dose	100-300 mcg per injection (equal of each component)
Frequency	2-3x daily -- ALWAYS FASTED (1-2hr after last meal)
Timing	Morning fasted + pre-bed optimal
Cycle	12 weeks on, 4 weeks off Stack/Cycle same as individual peptides
Storage	2-8 deg C Reconstituted: use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
GHRH (CJC)	Binds GHRH receptor; primes somatotrophs; enlarges GH pool via cAMP/PKA	Elevated GH release capacity
GHS (Ipa)	Binds GHSR-1a; triggers acute GH exocytosis; no cortisol/prolactin	Clean GH pulse; no side-pathway activation
Synergy	Two independent pathways fired simultaneously: GHRH primes + GHS releases. Combined GH 3-10x either alone	Maximum GH output per injection

Benefit	Context
Pre-Mixed Convenience	One vial eliminates separate reconstitution and dosing of two peptides.
Proven Synergy	GHRH + GHS is the gold standard combination for GH secretagogue research.
No Cortisol/Prolactin	Ipamorelin component ensures clean GH release -- no hormonal side effects.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: Vol(mL)=Dose(mcg)/Conc(mcg/mL) | **Units:** Vol x 100


5mg+5mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	100mcg	150mcg	200mcg	250mcg	300mcg
Vol(mL)	0.0100	0.0150	0.0200	0.0250	0.0300
Units	1.0U	1.5U	2.0U	2.5U	3.0U
Doses/Vial	100	66	50	40	33


10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	100mcg	150mcg	200mcg	250mcg	300mcg
Vol(mL)	0.0100	0.0150	0.0200	0.0250	0.0300
Units	1.0U	1.5U	2.0U	2.5U	3.0U
Doses/Vial	100	66	50	40	33

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Confirm fasted	1-2 hr after last meal. Mandatory.
2	Draw dose	Draw volume per table. One syringe = both peptides.
3	Inject SubQ	Pinch 2-3cm. 45 deg. Slowly. Rotate sites.
4	Wait 30 min	Before eating. Needle into sharps container.

Protocol	Dose	Frequency	Duration	Notes
Beginner	100 mcg	2x daily AM+bed	12wk on 4wk off	Start here. Both fasted.
Standard	200 mcg	2-3x daily fasted	12wk on 4wk off	200mcg CJC + 200mcg Ipa per injection.

Side Effect	Likelihood	Management
Injection site reaction	Common	Rotate sites. Fresh needle.
Water retention	Occasional	GH-mediated. Reduce dose.
Tingling / headache	Occasional	Transient. Resolves in 1-2 weeks.

Contraindications: Cancer history, pregnancy, under 18.

1. Always fasted (1-2hr after last meal). Insulin blocks GH release.
2. 1 mL BAC Water per vial -> 10mg/mL total (5mg/mL each component).
3. Dose = both peptides combined: 200mcg injection = 200mcg CJC + 200mcg Ipa.
4. Inject 2-3x daily: morning fasted + pre-bed for optimal GH windows.
5. Wait 30 min post-injection before any caloric intake.
6. Rotate sites through 8-point abdomen map. Swirl -- never shake.
7. Cycle 12 weeks on, 4 weeks off. Store 2-8 deg C. Use within 28 days.

DISCLAIMER: SaiyanMed research use only. Not for human therapeutic use. Consult a qualified professional. SaiyanMed assumes no liability.

SaiyanMed | saiyanmed.com | Research Use Only