

GHRP-6

GH Secretagogue | 10mg x 10 Vials


Potent GHS -- Strong Hunger Effect | Fasted 2-3x Daily

Purity: 99.8% | Endotoxin-Free | Research Use Only

GHRP-6 (Growth Hormone Releasing Peptide-6) is a hexapeptide GH secretagogue and ghrelin receptor agonist. It produces strong GH pulses with notable hunger stimulation (strongest of the GHRP family) -- making it useful for research subjects needing appetite increase alongside GH elevation. Must be fasted.


FASTED REQUIRED: Inject at least 1-2 hours after last meal. Insulin blunts GH release. Wait 30 min before eating post-injection.

GHRP-6 Vial Comparison



Parameter	Specification
Class	GH Secretagogue (GHS) -- ghrelin receptor agonist, potent hunger stimulant
Vial Size	10mg x 10 Purity 99.8%
Reconstitution	1 mL BAC Water: 10mg->10mg/mL (10,000 mcg/mL)
Standard Dose	100-300 mcg per injection -- always fasted
Frequency	2-3x daily fasted expect significant hunger post-injection
vs GHRP-2	Similar GH pulse strength; stronger hunger effect; comparable cortisol/prolactin
Cycle	12 weeks on, 4 weeks off
Storage	2-8 deg C Reconstituted: use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
GHSR-1a	Potent ghrelin receptor agonist; activates Gq/phospholipase C; strong GH granule exocytosis; also activates hunger centres in hypothalamus	Strong GH pulse; significant appetite stimulation
Hunger Stimulation	Ghrelin receptor activation in hypothalamic arcuate nucleus; NPY/AgRP neuron stimulation; potent orexigenic effect; useful in research requiring caloric surplus alongside GH	Appetite increase; useful for bulking/mass protocols
GHRH Synergy	Stacked with CJC-1295 or Sermorelin; dual pathway activation; GHRH primes + GHRP-6 triggers release; amplified GH 3-10x vs either alone	Maximum GH output when stacked with GHRH

Benefit	Context
Potent GH Output	GHRP-6 produces strong GH pulses via ghrelin receptor activation, comparable to GHRP-2 in magnitude.
Appetite Stimulation	The strongest appetite stimulator in the GHRP family -- useful for protocols requiring caloric surplus alongside GH elevation.
Stacking	Combines powerfully with GHRH analogues (CJC-1295, Sermorelin) for 3-10x amplified GH pulses.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.




1mL Insulin Syringe | 29-31G | SubQ

Formula: $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$ | **Units:** $\text{Vol} \times 100$


10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	100mcg	150mcg	200mcg	250mcg	300mcg
Vol(mL)	0.0100	0.0150	0.0200	0.0250	0.0300
Units	1.0U	1.5U	2.0U	2.5U	3.0U
Doses/Vial	100	66	50	40	33

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Confirm fasted	1-2 hr after last meal. Mandatory.
2	Draw dose	Draw volume per dose table. Can combine with GHRH in one syringe.
3	Inject SubQ	Abdomen. Pinch 2-3cm. 45 deg. Rotate sites.
4	Expect hunger	Significant hunger is expected 20-30 min post-injection. Plan meals.

Protocol	Dose	Frequency	Duration	Notes
Standard	200 mcg	2-3x daily fasted	12wk on 4wk off	Expect strong hunger. Time injection before planned meal.
+GHRH	200mcg GHRP-6 +CJC-1295	2x daily fasted	12wk on 4wk off	Combine in syringe. Maximum GH output.

Side Effect	Likelihood	Management
Intense hunger 20-30 min post	Certain	Expected ghrelin effect. Time injection near meal if desired.
Cortisol/prolactin elevation	Mild	Minor at standard doses. Use Ipamorelin if sensitivity.
Water retention	Occasional	GH-mediated. Reduce dose if significant.

Contraindications: Cancer history, pregnancy, under 18.

1. Always inject fasted (1-2hr after last meal). Insulin blunts GH release.
2. 1 mL BAC Water: 10mg->10mg/mL (10,000 mcg/mL).
3. Strong hunger is expected 20-30 min post-injection -- time accordingly.
4. Stack with CJC-1295 for 3-10x amplified GH pulse.
5. If hunger side effect is unwanted, switch to Ipamorelin (no hunger).
6. Inject 2-3x daily. Wait 30 min before eating post-injection.
7. Cycle 12 weeks on, 4 weeks off. Store 2-8 deg C. 28 days max.

DISCLAIMER: SaiyanMed research use only. Not for human therapeutic use. Consult a qualified professional. SaiyanMed assumes no liability.

SaiyanMed | saiyanmed.com | Research Use Only