

KLOW

BPC-157 10mg + TB-500 10mg + GHK-Cu 50mg + KPV 10mg | 80mg x 10 Vials

Complete 4-Peptide Repair, Regeneration & Anti-Inflammatory Stack

1mL BAC Water -> 80mg/mL total | SubQ or IM | No Fasting Required

Purity: 99.8% | Endotoxin-Free | Research Use Only

KLOW is the most comprehensive pre-mixed peptide repair stack in the SaiyanMed range -- combining four synergistic peptides in one 80mg vial. KLOW adds KPV (Lys-Pro-Val, the anti-inflammatory C-terminal fragment of alpha-MSH) to the GLOW formula, providing potent cytokine suppression and MC-receptor-mediated anti-inflammatory resolution alongside the repair, cell migration, and collagen synthesis of the other three components.


KLOW -- Active Components per Vial

BPC-157 10mg #065f46	TB-500 10mg #0f766e	GHK-Cu 50mg #92400e	KPV 10mg #9f1239
---	--	--	-------------------------------------

All components pre-mixed in one vial | One injection delivers full stack

Parameter	Specification
BPC-157	10mg per vial -- GI protection, angiogenesis, tendon repair, eNOS/NO pathway
TB-500	10mg per vial -- actin regulation, cell migration, anti-inflammatory, cardiac/neural repair
GHK-Cu	50mg per vial -- collagen I/III synthesis, gene modulation, antioxidant, wound healing
KPV	10mg per vial -- alpha-MSH C-terminal tripeptide; MC receptor anti-inflammatory; cytokine suppression
Total	80mg per vial x 10 vials per kit 800mg total
Reconstitution	1 mL BAC Water -> 80mg/mL total (1mL = full 80mg dose)
Standard Dose	0.5-1.0 mL per injection (40-80mg) SubQ or IM
Frequency & Storage	2-3x/week no fasting 2-8 deg C use within 28 days

KLOW | MECHANISM OF ACTION & SYNERGY



Complementary, non-overlapping pathways -- synergistic tissue repair

Component	Mechanism	Research Effect
BPC-157	GI mucosal protection; eNOS/NO upregulation; VEGF angiogenesis; tendon fibroblast activation; no known LD50	GI repair; vascular protection; tendon/ligament
TB-500	Thymosin Beta-4 fragment; G-actin sequestration for cell migration; NF-kB inhibition; VEGF; cardiomyocyte survival	Cell migration; anti-inflammatory; cardiac/neural
GHK-Cu	Copper tripeptide; TGF-beta/collagen I/III; 4,000+ gene modulation toward younger phenotype; Cu/Zn-SOD antioxidant	Collagen synthesis; gene regulation; anti-aging
KPV	Alpha-MSH C-terminal (Lys-Pro-Val); binds MC1R/MC3R/MC5R on immune cells; suppresses NF-kB, TNF-alpha, IL-6, IL-1beta; promotes Treg differentiation	Cytokine suppression; anti-inflammatory resolution; immune balance

Benefit	Research Context
Four-Pathway Synergy	BPC-157 (vascular/GI) + TB-500 (cell migration) + GHK-Cu (collagen) + KPV (cytokine suppression) -- four complementary, non-overlapping mechanisms in one injection.
KPV Anti-Inflammatory Amplification	KPV suppresses the inflammatory cytokine cascade (NF-kB, TNF-alpha, IL-6) that can limit tissue repair -- creating a more favourable environment for BPC/TB/GHKCu to work.
Most Complete Pre-Mixed Stack	KLOW is the most comprehensive pre-mixed tissue repair and anti-inflammatory stack available -- replaces 4 separate reconstitution and dosing steps.


KLOW | RECONSTITUTION & DOSE GUIDE

80mg vial (80mg total) -- Add 1 mL BAC Water -> 80mg/mL total. One injection of 1 mL delivers the full dose of all components.



1 mL BAC Water per vial. Roll gently -- NEVER shake.


Step	Action	Detail
1	Gather supplies	Vial, BAC Water, 29-31G insulin syringe, alcohol swabs, sharps container.
2	Clean stoppers	Wipe both vial stoppers with alcohol swab. Air-dry 10-15 seconds.
3	Draw 1 mL BAC Water	Exactly 1 mL BAC Water. Expel all air bubbles.
4	Inject into vial	Aim stream at GLASS WALL -- never onto powder. Inject slowly.
5	Swirl gently & label	Roll between palms 20-30 sec. Clear within 60 sec. Never shake. Label with date. Store 2-8 deg C. Use within 28 days.




1mL Insulin Syringe | 29-31G | SubQ

At 80mg/mL: 1 mL = full 80mg dose | 0.5 mL = half dose | Each vial contains 10 doses at 0.1 mL (1/10 dose per injection)

Injection Volume	0.1 mL (1/10 dose)	0.25 mL (1/4 dose)	0.5 mL (1/2 dose)	1.0 mL (full dose)
Syringe Units	10U	25U	50U	100U
Total Peptide	8mg	20mg	40mg	80mg
Doses/Vial	10	4	2	1



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Reconstitute	Add 1 mL BAC Water. Swirl gently 20-30 sec. Never shake. Clear solution required.
2	Draw dose	0.5 mL (half dose / 40mg) or 1.0 mL (full dose / 80mg). Expel air bubbles.
3	Select site	Inject near target tissue for localised repair, or abdomen for systemic benefit.
4	Inject SubQ/IM	SubQ: pinch 2-3cm, 45 deg, inject slowly. IM: 23-25G, 90 deg, aspirate.
5	Rotate & dispose	Record site. 8-point rotation. Press 5 sec. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Acute Repair	1.0 mL (full dose)	2-3x/week	4-8 wk	Near injury. Maximises all 4 pathways.
Inflammatory Conditions	0.5-1.0 mL	Daily or 2-3x/week	4-12 wk	KPV anti-inflammatory + repair stack.
Maintenance	0.5 mL (half dose)	2x/week	Ongoing	Lower ongoing dose after acute phase.

Side Effect	Likelihood	Management
Injection site reaction	Common	Mild redness. Rotate sites. Fresh needle.
Temporary skin darkening	Occasional	GHK-Cu copper effect. Not harmful. Fades on cessation.
Mild fatigue	Occasional	Systemic repair and immune modulation response. Resolves.

No KPV melanocortin side effects	Confirmed	KPV is C-terminal fragment only -- no tanning, appetite, or libido effects.
----------------------------------	-----------	---

Contraindications: Active cancer history, pregnancy, lactation, under 18.

1. KLOW = BPC-157 10mg + TB-500 10mg + GHK-Cu 50mg + KPV 10mg in one vial.
2. 1 mL BAC Water -> 80mg/mL. Full dose = 1 mL (80mg). Half dose = 0.5 mL (40mg).
3. No fasting required. Inject any time of day.
4. KLOW adds KPV anti-inflammatory to the GLOW formula -- ideal for inflammatory conditions.
5. Inject near target tissue for localised benefit, or abdomen for systemic effect.
6. KPV has no melanocortin side effects (no tanning, appetite, or libido effects).
7. Each vial: 10 full doses (1 mL) or 20 half doses (0.5 mL).
8. Rotate sites through 8-point map. Swirl gently. Store 2-8 deg C, 28 days.
9. Contraindicated in active cancer history. Not for pregnancy or under 18.

DISCLAIMER: This document is produced by SaiyanMed for informational and research reference purposes only. KLOW is a research product not approved for human therapeutic use. This guide does not constitute medical advice. SaiyanMed assumes no liability for misuse.

SaiyanMed | saiyanmed.com | Research Use Only