



LEMON BOTTLE

INJECTABLE FAT DISSOLVING SOLUTION | 10mL | PRE-MIXED

Riboflavin (B2) + Bromelain + Deoxycholic Acid

Ready to Use | Subdermal / IM Injection | Research Use Only

Lemon Bottle is a pre-mixed, ready-to-use injectable lipolytic solution. No reconstitution required. It is administered directly into localised subcutaneous fat deposits using a fine needle. It works via deoxycholic acid-mediated fat cell membrane disruption, bromelain enzymatic action, and riboflavin metabolic support to break down and eliminate stubborn fat deposits.

IMPORTANT: Lemon Bottle is a localised lipolytic compound, NOT a systemic weight-loss injection. It must be injected directly into the target fat deposit at the correct subdermal depth. This is not a SubQ peptide injection — technique and site selection are critical.

Parameter	Specification
Product Name	Lemon Bottle Injectable Fat Dissolving Solution
Formula	Riboflavin (Vitamin B2) + Bromelain (Pineapple Enzyme) + Deoxycholic Acid (bile acid derivative)
Format	Pre-mixed sterile liquid solution — NO reconstitution required
Vial Volume	10 mL per vial 10 vials per kit
Dose per Site	0.1-0.2 mL per injection point (spaced 1-2 cm apart)
Total per Session	Max 10 mL per session across all treatment areas
Route	Subdermal / Subcutaneous into fat deposits ONLY — NOT intramuscular
Needle	30-32G x 4-6 mm (subdermal needle) — short and fine
Injection Depth	4-6 mm — into subcutaneous fat layer only. Never into muscle.
Session Interval	Minimum 4 weeks between sessions on the same area
Sessions Required	Typically 2-4 sessions per treatment area
Appearance	Clear to pale yellow liquid — normal. Discard if cloudy or has particles.
Storage	2-8 deg C refrigerated Protect from light Do not freeze

ACTIVE INGREDIENTS & MECHANISM OF ACTION

Lemon Bottle's three active ingredients work synergistically to disrupt fat cell membranes, enzymatically process released lipids, and support mitochondrial fat metabolism — resulting in permanent localised fat reduction when correctly administered.

Lemon Bottle Mechanism of Action -- Localised Lipolysis



Ingredient	Class	Mechanism	Benefit
Deoxycholic Acid	Bile acid derivative (naturally occurring)	Disrupts phospholipid bilayer of fat cell membranes; causes adipocytolysis (fat cell destruction); triggers local inflammatory response that drives macrophage-mediated fat clearance	Permanent localised fat reduction; FDA-approved as Kybella for submental fat. Primary active lipolytic agent.
Bromelain	Proteolytic enzyme from pineapple stem	Breaks down proteins and lipid complexes released after fat cell disruption; anti-inflammatory activity; reduces post-treatment swelling and bruising; improves lymphatic drainage of debris	Faster resolution of post-injection swelling; reduces downtime; supports clearance of disrupted fat cell components
Riboflavin (Vitamin B2)	Water-soluble B-vitamin (coenzyme)	Cofactor in FAD/FMN electron carriers; essential for beta-oxidation of fatty acids; supports mitochondrial energy production from liberated fatty acids; also serves as a visible yellow marker	Metabolic processing of released fatty acids; energy generation; characteristic yellow colour of the solution confirms riboflavin presence


Key Research Benefits

Benefit	Context
Permanent Fat Cell Destruction	Deoxycholic acid-mediated adipocytolysis is irreversible — destroyed fat cells do not regenerate. Results are long-lasting when combined with stable body weight.
Localised Precision	Direct injection into the target fat deposit allows precise contouring of specific areas that are resistant to diet and exercise.
Minimal Systemic Exposure	Unlike oral or systemic fat-loss agents, Lemon Bottle acts locally. Deoxycholic acid is naturally produced by the body and is rapidly metabolised.
Low Downtime vs Surgery	Non-surgical alternative to liposuction for small, localised fat deposits. Typical swelling resolves within 3-7 days; most resume normal activity immediately.
Bromelain Anti-Inflammatory Advantage	The addition of bromelain differentiates Lemon Bottle from deoxycholic acid alone (Kybella) by actively reducing post-injection inflammatory response and swelling.

INJECTION SITES, DEPTH & DOSING


Critical: Lemon Bottle must be injected into subcutaneous fat tissue at 4-6 mm depth using a 30-32G short needle. Incorrect depth risks muscle injection (pain, damage) or intradermal injection (skin necrosis). Space each injection point 1-2 cm apart in a grid pattern over the target area.

Lemon Bottle Treatment Areas -- Localised Fat Deposits Only



INJECT INTO FAT DEPOSITS ONLY -- Never into muscle, vessel or nerve

Correct Injection Depth -- Subcutaneous Fat Layer Only



Treatment Zone Volumes -- Per Session Guidelines

Abdomen 2-4 mL per session 1-2 cm grid	Flanks 1-2 mL per session 1-2 cm grid	Upper Arms 1-2 mL per session 1-2 cm grid	Inner Thighs 2-4 mL per session 1-2 cm grid	Outer Thighs 2-4 mL per session 1-2 cm grid	Double Chin 0.5-1 mL per session 0.5 cm grid
--	---	---	---	---	--

Do not exceed 10 mL total per session. Space injections 1-2 cm apart.

Area	Needle Size	Depth	Volume per Point	Max per Session
Abdomen	30-32G x 6mm	5-6 mm	0.1-0.2 mL	4 mL
Flanks	30-32G x 6mm	5-6 mm	0.1-0.2 mL	2 mL
Upper Arms	30-32G x 4mm	4-5 mm	0.1 mL	2 mL
Inner/Outer Thighs	30-32G x 6mm	5-6 mm	0.1-0.2 mL	4 mL
Submental (chin)	30G x 4mm	4 mm	0.1 mL	1 mL

INJECTION PROCEDURE & TREATMENT TIMELINE

Step-by-Step Injection Procedure

Step	Action	Detail
1	Consultation & mapping	Mark treatment area with a skin marker in a 1-2 cm grid pattern. Photograph before treatment for comparison.
2	Clean and prepare area	Cleanse the entire treatment area with an antiseptic solution. Apply topical anaesthetic cream 20-30 minutes before treatment if required for comfort.
3	Prepare syringe	Draw Lemon Bottle directly from the vial into a 1 mL or 3 mL syringe. Attach a 30-32G x 4-6 mm needle. Expel air bubbles.
4	Pinch and insert	Pinch the fat tissue between thumb and forefinger to elevate the fat layer. Insert needle at 45 degrees, 4-6 mm depth into the fat — not skin, not muscle.
5	Inject per point	Inject 0.1-0.2 mL per injection point. Move to next grid point 1-2 cm away. Do not exceed 10 mL total per session.
6	Apply pressure	After completing the grid, press the area gently with sterile gauze. Do NOT rub — massage can disperse solution beyond target zone.
7	Post-treatment care	Apply ice pack (wrapped, never directly on skin) for 10-15 minutes to reduce swelling. Inform client of expected reactions (see side effects).
8	Documentation	Record volume used, areas treated, and client response. Schedule follow-up at minimum 4 weeks for the same area.

Treatment Course Timeline

Treatment Course -- Minimum 4 Weeks Between Sessions



Most clients require 2-4 sessions per treatment area

Timeframe	Expected Reactions	Care Instructions
0-24 hours	Redness, swelling, warmth, tenderness at injection sites	Ice pack 10-15 min intervals. Avoid heat, exercise, alcohol.
1-3 days	Peak swelling and firmness ("orange peel" texture normal)	Hydrate well. Gentle lymphatic massage if recommended. Avoid tight clothing over treated area.
3-7 days	Swelling subsides. Possible bruising. Area feels lumpy.	Resume normal activity. Continue hydration. No saunas/hot baths.
2-4 weeks	Visible fat reduction begins. Skin may feel tighter.	Assess results. Schedule next session if additional treatment needed.
4-8 weeks	Full results of session visible. Area noticeably reduced.	Maintain stable body weight for lasting results.

SIDE EFFECTS, SAFETY, STORAGE & GOLDEN RULES

Side Effect	Likelihood	Management
Swelling at treatment site	Certain	Expected and necessary part of the process. Ice 10-15 min post-treatment. Resolves 3-7 days.
Redness and warmth	Certain	Normal inflammatory response. Resolves within 24-48 hours.
Tenderness / pain at site	Very Common	Expected. Topical anaesthetic pre-treatment. Paracetamol if needed. Avoid NSAIDs first 24h (may blunt inflammatory response needed for results).
Bruising	Common	Use fine 30-32G needle. Avoid blood thinners 5 days before treatment. Resolves 5-10 days.
Lumpy / firm texture	Common	Normal during healing phase (1-3 weeks). Gentle massage after 72 hours if recommended. Resolves fully.
Numbness at site	Occasional	Temporary. Resolves within days to weeks. If persistent, review technique.
Skin discolouration	Occasional	Yellow tint from riboflavin — harmless, fades within hours. Post-inflammatory hyperpigmentation possible in darker skin tones — use SPF 50+ on treated area.
Necrosis / ulceration	Rare (technique error)	Risk if injected too superficially (intradermal) or into vascular structures. Use correct needle and depth. Seek medical attention immediately.

STOP & SEEK MEDICAL ADVICE: Signs of infection: increasing redness, spreading warmth, pus, fever above 38 deg C

STOP & SEEK MEDICAL ADVICE: Skin breakdown, blistering, or open wounds at injection site (possible necrosis)

STOP & SEEK MEDICAL ADVICE: Severe difficulty swallowing if treating neck/chin area (nerve or muscle involvement)

STOP & SEEK MEDICAL ADVICE: Severe allergic reaction: hives, facial swelling, difficulty breathing (anaphylaxis)

STOP & SEEK MEDICAL ADVICE: Persistent numbness or paralysis beyond 4 weeks

Absolute Contraindications: Pregnancy or breastfeeding, active infection or open wounds in treatment area, allergy to any component (deoxycholic acid, bromelain, riboflavin), autoimmune conditions affecting skin, bleeding disorders or anticoagulant therapy, body dysmorphic disorder. Not for use in individuals under 18. Do not treat over bony prominences, tendons, nerves, or vascular structures.

Storage Guidelines

State	Temperature	Duration	Notes
Unopened vial	2-8 deg C	Per expiry	Keep upright. Away from light. Do not freeze.
Opened / in-use	2-8 deg C	Use same session	Discard any unused solution from an opened vial after the session.
Prepared syringe	Room temp	Use within 30 min	Do not pre-draw syringes for later use.
Transport	Below 25 deg C	Max 48 hr	Insulated cooler bag. Protect from light and temperature extremes.

Golden Rules — Lemon Bottle

1. Inject into subcutaneous fat ONLY -- correct depth is 4-6 mm. Never into skin or muscle.
2. Use a 30-32G x 4-6 mm needle. Never use a standard SubQ peptide needle for this treatment.
3. Space each injection point exactly 1-2 cm apart in a grid pattern over the target area.
4. Do not exceed 10 mL total Lemon Bottle per session across all treatment areas.
5. Allow a minimum of 4 weeks between sessions on the same treatment area.
6. Swelling, redness, and firmness are EXPECTED and part of the lipolytic process -- not a complication.
7. Do NOT rub or massage the area immediately post-injection -- press gently with gauze only.
8. Apply ice (not direct contact) for 10-15 minutes post-treatment to manage swelling.
9. Discard any unused solution from an opened vial after the session -- do not save for later.
10. This is a localised contouring treatment, not a systemic weight-loss solution. Diet and exercise remain essential.

DISCLAIMER: This document is produced by SaiyanMed for informational and research reference purposes only. Lemon Bottle injectable solution must only be administered by a trained and qualified practitioner. This guide does not constitute medical advice. Self-administration is not recommended. SaiyanMed assumes no liability for adverse outcomes arising from improper use of this product.

SaiyanMed | saiyanmed.com | Research Use Only | For Educational Purposes