

LIPO-C

Lipotropic Cocktail | MIC + L-Carnitine + B12 + B5 + B6

Complete Injection & Usage Guide -- 10mL x 10 Vials


Pharmaceutical Grade | Pre-Mixed Solution | Ready to Use | Research Use Only

Key Product Information

Lipo-C is a pre-mixed, ready-to-use liquid solution. No reconstitution is required. Draw directly from the vial using a clean syringe. Each 10mL vial provides multiple doses depending on your protocol (typical dose: 0.5-1.0 mL per injection).

Parameter	Details
Format	Pre-mixed sterile liquid solution — ready to use, NO reconstitution required
Vial Size	10 mL per vial 10 vials per kit 100 mL total
Typical Dose	0.5 mL to 1.0 mL per injection
Doses per Vial	10 doses at 1.0 mL 20 doses at 0.5 mL
Doses per Kit	100 doses at 1.0 mL 200 doses at 0.5 mL
Route	Intramuscular (IM) preferred Subcutaneous (SubQ) alternative
Frequency	1-3x per week (research protocol)
Storage	2-8 deg C refrigerated Protect from light Do not freeze
Appearance	Clear to slightly yellow/pink liquid — normal; discard if cloudy or particulate

Active Ingredient Formula




ACTIVE INGREDIENTS & MECHANISM OF ACTION

Lipo-C combines six synergistic lipotropic and B-vitamin compounds that work together to support hepatic fat metabolism, mitochondrial energy production, and cellular methylation. Each ingredient contributes a distinct and complementary mechanism.

Ingredient	Class	Mechanism	Research Benefit
Methionine	Essential amino acid (sulphur-containing)	Precursor to SAM-e (S-adenosyl-methionine); drives hepatic methylation cycles; promotes liver fat export as VLDL	Liver detoxification, fat mobilisation, homocysteine regulation
Inositol	Carbocyclic polyol (B-vitamin family)	Regulates insulin receptor signalling; component of cell membrane phospholipids; reduces hepatic fat accumulation	Insulin sensitisation, liver fat reduction, mood regulation
Choline	Essential nutrient (quaternary ammonium)	Required for VLDL synthesis and hepatic fat export; precursor to acetylcholine and phosphatidylcholine	Prevents fatty liver; cognitive support; muscle function
L-Carnitine	Amino acid derivative	Transports long-chain fatty acids across inner mitochondrial membrane for beta-oxidation (fat burning)	Enhanced fat oxidation, energy production, exercise performance
B12 (Cyanocobalamin)	Water-soluble vitamin (cobalamin)	Cofactor in methylmalonyl-CoA mutase; myelin synthesis; red blood cell formation; homocysteine conversion to methionine	Energy metabolism, neurological health, red blood cell production
B5 (Dexpanthenol)	Provitamin B5 (pantothenol)	Converted to pantothenic acid and then to Coenzyme A; essential for fatty acid metabolism and citric acid cycle	Energy production, fat metabolism, adrenal hormone synthesis
B6 (Pyridoxine)	Water-soluble vitamin (pyridoxal family)	Cofactor in over 100 enzymatic reactions; amino acid metabolism; neurotransmitter synthesis; glycogen phosphorylase	Protein metabolism, mood regulation, immune function

DOSE GUIDE & INJECTION TECHNIQUE

No reconstitution needed. Draw Lipo-C directly from the vial. Clean the vial rubber stopper with an alcohol swab before each draw.



Standard 1mL Syringe -- Draw Lipo-C directly (no reconstitution)

Protocol	Dose per Injection	Frequency	Route	Notes
Standard	1.0 mL (1cc)	3x per week	IM preferred	Mon / Wed / Fri — most common research protocol
Moderate	0.5 mL	3x per week	IM or SubQ	Suitable for smaller individuals or first-time users
Maintenance	1.0 mL	1-2x per week	IM or SubQ	Once weight loss goal reached; ongoing metabolic support
Daily (Research)	0.5 mL	Daily (5 days on, 2 days off)	SubQ	Higher-frequency research protocol; rotate sites daily

Step-by-Step Injection Procedure


Step	Action	Detail
1	Refrigerator to room temp	Remove vial from fridge 15-20 minutes before injection. Cold solution causes unnecessary discomfort at the injection site.
2	Wash hands	Wash thoroughly with soap and warm water for 20 seconds. Dry completely.
3	Gather materials	Lipo-C vial, 1mL syringe with needle (23-25G x 1-1.5" for IM; 29-31G x 5/16" for SubQ), alcohol swabs, sharps container.
4	Clean vial stopper	Wipe the rubber stopper with a fresh alcohol swab. Allow to air-dry for 10 seconds.
5	Draw the dose	Insert needle into vial. Invert vial. Draw prescribed volume (0.5 or 1.0 mL). Remove air bubbles by tapping and expelling.
6	Clean injection site	For IM: wipe deltoid or thigh with alcohol swab, dry 10-15 sec. For SubQ: wipe abdomen or love-handle area.
7	Inject	IM: Insert needle at 90 degrees with a dart-like motion; inject slowly. SubQ: Pinch skin, insert at 45 degrees, release pinch, inject slowly.
8	Withdraw and press	Remove needle. Press site with dry swab 5-10 seconds. Do NOT rub — some localised burning/warmth is normal with Lipo-C.
9	Safe disposal	Place needle immediately into sharps container. Never recap, reuse, or leave exposed.

INJECTION SITES, ROTATION & SCHEDULE

Preferred route: Intramuscular (IM) into the deltoid muscle (upper arm) for fastest absorption and minimal discomfort. **Subcutaneous (SubQ)** into the abdomen is a suitable alternative. Always rotate injection sites systematically to prevent localised irritation, tissue hardening, or lipodystrophy at any single site.

Site	Route	Needle	Notes
Deltoid (upper arm)	IM (Preferred)	23-25G 1 to 1.5 inch	Fastest absorption. Alternate left and right arm each session.
Vastus Lateralis (outer thigh)	IM (Alternative)	23-25G 1 to 1.5 inch	Good alternative for higher volumes. Use outer mid-thigh only.
Abdomen (2 inches from navel)	SubQ (Alternative)	29-31G 5/16 inch	Pinch skin. 45-degree angle. Slower absorption than IM.
Upper arm (back)	SubQ (Alternative)	29-31G 5/16 inch	Suitable for SubQ if abdomen is irritated. Requires assistance.

Lipo-C Injection Sites -- IM Preferred | SubQ Alternative



IM (90 deg) vs SubQ (45 deg) Injection Angles



Recommended Schedule -- 3x Per Week (Mon / Wed / Fri)



Alternative: 1-2x per week for maintenance. Daily use also researched.

SIDE EFFECTS, STORAGE & GOLDEN RULES

Side Effect	Likelihood	Management
Localised burning / stinging at injection site	Very Common	Normal with Lipo-C. Warm the vial to room temperature before injecting. Inject slowly. Do not rub site after injection.
Redness or warmth at injection site	Common	Expected local reaction. Rotate sites to prevent irritation. Resolves within 1-2 hours.
Nausea (especially at higher doses)	Occasional	Inject with a light meal. Reduce to 0.5 mL if persistent.
Urinary odour change	Occasional	Normal sulphur by-product of methionine metabolism. Not harmful.
Fatigue or mild headache	Occasional	Usually transient. Stay hydrated. Ensure adequate B-vitamin baseline.
Increased energy / mild stimulant effect	Common	Expected effect of B12 + Carnitine. Inject morning or midday to avoid sleep disruption.
Allergic reaction at site	Rare	Stop use. Seek medical advice. Do not rechallenge without assessment.

STOP & SEEK MEDICAL ADVICE: Severe or spreading redness, swelling, heat at injection site (possible infection)

STOP & SEEK MEDICAL ADVICE: Difficulty breathing or swallowing after injection (anaphylaxis)

STOP & SEEK MEDICAL ADVICE: Severe chest pain or palpitations

STOP & SEEK MEDICAL ADVICE: Jaundice or dark urine (hepatic concern)

STOP & SEEK MEDICAL ADVICE: Numbness or tingling in extremities (high-dose B6 neuropathy — reduce dose)

Contraindications: Known allergy to any Lipo-C component, active liver disease or hepatic impairment, kidney disease (methionine caution), pregnancy or lactation, trimethylaminuria (fish odour syndrome — choline contraindicated). Not for use under 18.

Storage Guidelines

State	Temperature	Duration	Notes
Unopened vial	2-8 deg C	Per expiry	Keep upright. Protect from light. Do not freeze.
Opened/in-use vial	2-8 deg C	28 days	Cap securely after each draw. Inspect for cloudiness before use.
Short-term transport	Below 25 deg C	Max 72 hr	Insulated bag with ice packs. Avoid direct ice contact.
Never freeze	Do NOT freeze	N/A	Freezing degrades B12 and disrupts the solution. Discard if frozen.

Golden Rules — Lipo-C

1. Warm the vial to room temperature for 15-20 minutes before injecting — cold solution stings.
2. Draw Lipo-C directly from the vial — NO reconstitution is required.
3. IM injection into the deltoid is the preferred route for fastest absorption.
4. Inject slowly over 5-10 seconds to minimise localised burning.

5. Do NOT rub the injection site after withdrawal — mild warmth and redness are normal.

6. Rotate injection sites every session. Allow at least 1 week rest per site.

7. Inject morning or midday — B12 and carnitine can cause energy stimulation.

8. Inspect the vial before each draw — discard if cloudy, discoloured, or particulate.

9. Store at 2-8 deg C. Never freeze. Opened vial is good for 28 days.

10. Contraindicated in liver disease, kidney disease, and known component allergy.

DISCLAIMER: This document is produced by SaiyanMed for informational and research reference purposes only. This guide does not constitute medical advice and must not be used to guide self-administration or therapeutic use without qualified medical supervision. Consult a licensed medical professional before beginning any injection protocol. SaiyanMed assumes no liability for misuse of this product or information herein.

SaiyanMed | saiyanmed.com | Research Use Only | For Educational Purposes