

Selank

Anxiolytic Nootropic Peptide | 5mg & 10mg x 10 Vials

Anxiety | Cognition | BDNF Upregulation | SubQ / Intranasal


Purity: 99.8% | Endotoxin-Free | Research Use Only

Selank is a synthetic heptapeptide analogue of tuftsin (Thr-Lys-Pro-Arg-Pro-Gly-Pro) developed by the Institute of Molecular Genetics (Russian Academy of Sciences). It exhibits anxiolytic, nootropic, and immunomodulatory effects with a unique safety profile -- no dependence, sedation, or withdrawal. Also available as an intranasal formulation for direct CNS delivery.



Parameter	Specification
Class	Synthetic heptapeptide anxiolytic/nootropic -- tuftsin analogue
Vial Sizes	5mg x 10 10mg x 10 Purity 99.8%
Reconstitution	1 mL BAC Water: 5mg->5mg/mL 10mg->10mg/mL
Standard Dose	250-1,000 mcg per injection SubQ intranasal also effective
Routes	SubQ Intranasal (dilute, 1-2 drops/nostril) no fasting
Frequency	1-3x daily acute or ongoing protocols
Cycle & Storage	Ongoing (low dependence risk) 2-8 deg C use within 28 days

MECHANISM OF ACTION & RESEARCH BENEFITS



Pathway	Mechanism	Benefit
Anxiolytic (GABAergic)	Modulates GABA-A receptor sensitivity; anxiolytic without sedation or dependence; reduces anxiety comparable to benzodiazepines in animal models but without tolerance; normalises HPA axis stress response	Anxiety; stress; PTSD; anxiolytic research without dependence
Cognitive & BDNF	Upregulates BDNF (brain-derived neurotrophic factor); enhances memory consolidation; improves learning and attention; nootropic effects independent of anxiolytic mechanism	Cognitive enhancement; memory; BDNF research
Immunomodulation & Enkephalins	Tuftsins-derived immunostimulant activity; modulates enkephalin metabolism (endogenous opioid regulation); mood-stabilising; antidepressant effects in research	Immune support; mood regulation; endogenous opioid research

Benefit	Context
No Dependence	Selank does not cause tolerance, dependence, or withdrawal -- a key advantage over benzodiazepines for anxiety research.
BDNF Upregulation	Unique dual mechanism: anxiolytic plus nootropic via BDNF. Most anxiolytics impair cognition -- Selank improves it.
Intranasal Option	Intranasal delivery provides rapid CNS-targeted effect for acute anxiety protocols.


RECONSTITUTION & DOSE CALCULATION

All vials use 1 mL BAC Water. Room temperature. Swirl gently -- never shake.



1 mL BAC Water per vial. Roll gently -- NEVER shake.

Step	Action	Detail
1	Gather	Vial, BAC Water, insulin syringe (29-31G), swabs, sharps.
2	Clean stoppers	Wipe both stoppers. Air-dry 10-15 sec.
3	Draw 1mL BAC	Exactly 1 mL. Expel air bubbles.
4	Inject into vial	Aim at GLASS WALL. Slowly. Never onto powder.
5	Swirl & label	Roll 20-30 sec. Clear within 60 sec. Never shake. Label date. 2-8 deg C. Use within 28 days.



1mL Insulin Syringe | 29-31G | SubQ

Formula: $\text{Vol(mL)} = \frac{\text{Dose(mcg)}}{\text{Conc(mcg/mL)}}$ | **Units:** $\text{Vol} \times 100$


5mg x 10 -- 1mL BAC -> 5.0mg/mL (5,000mcg/mL)

Dose(mcg)	250mcg	500mcg	750mcg	1000mcg
Vol(mL)	0.0500	0.1000	0.1500	0.2000
Units	5.0U	10.0U	15.0U	20.0U
Doses/Vial	20	10	6	5


10mg x 10 -- 1mL BAC -> 10.0mg/mL (10,000mcg/mL)

Dose(mcg)	250mcg	500mcg	750mcg	1000mcg
Vol(mL)	0.0250	0.0500	0.0750	0.1000
Units	2.5U	5.0U	7.5U	10.0U
Doses/Vial	40	20	13	10

INJECTION TECHNIQUE, PROTOCOLS & GOLDEN RULES



8-Point Rotation Map -- Rotate Every Injection



Step	Action	Detail
1	Select route	SubQ (29-31G, abdomen) or intranasal (dilute, nasal spray).
2	Draw/prepare	SubQ: draw dose. Intranasal: dilute to 0.15% solution in saline.
3	Administer	SubQ: 45 deg, pinch skin. Intranasal: tilt head, spray, sniff.
4	Rotate & dispose	8-point rotation for SubQ. Sharps container.

Protocol	Dose	Frequency	Duration	Notes
Anxiety Protocol	250-500 mcg	1-3x daily	Ongoing (low risk)	No dependence or tolerance. Safe ongoing use.
Cognitive Enhancement	500-1,000 mcg	1-2x daily	8-12 wk	BDNF upregulation and memory consolidation.

Side Effect	Likelihood	Management
Mild sedation at high dose	Rare	GABA modulation. Reduce dose.
Mood improvement	Expected effect	Anxiolytic and mood-normalising effect.

Contraindications: Cancer history, pregnancy, under 18.

1. 5mg: 1mL BAC->5mg/mL | 10mg: 1mL->10mg/mL.

2. No fasting required. Inject any time -- morning and afternoon preferred.
3. SubQ injection or intranasal delivery -- both effective.
4. Intranasal: dilute to 0.15% (1.5mg/mL) in saline. 1-2 drops per nostril.
5. No tolerance, dependence, or withdrawal -- safe for ongoing use.
6. Upregulates BDNF: cognitive enhancement alongside anxiolytic effects.
7. Rotate SubQ sites through 8-point abdomen map.
8. Store 2-8 deg C. Use within 28 days.
9. Not for use during pregnancy, lactation, or in individuals under 18.

DISCLAIMER: SaiyanMed research use only. Not for human therapeutic use. Consult a qualified professional. SaiyanMed assumes no liability.

SaiyanMed | saiyanmed.com | Research Use Only