

✔ Congratulations! You passed!

Grade received 100% To pass 80% or higher

Go to next item

1. Assume that there are 2 happy people and 2 unhappy people in a room. Concretely, persons A and B are happy and persons C and D are unhappy. If you were to randomly pick a person from the room, what is the probability that the person is happy.

1 / 1 point

- ☒ 1/2
☐ 1/4
☐ 3/4
☐ 0

✔ Correct

2. Assume that there are 2 happy people and 2 unhappy people in a room. Concretely, persons A and B are happy and persons C and D are unhappy. If a friend showed you the part of the room where the two happy people are, what is the probability that you choose person B?

1 / 1 point

- ☒ 1/2
☐ 1/4
☐ 3/4
☐ 1

✔ Correct

3. From the equations presented below, express the probability of a tweet being positive given that it contains the word happy in terms of the probability of a tweet containing the word happy given that it is positive

1 / 1 point

$$P(\text{Positive} \mid \text{"happy"}) = \frac{P(\text{Positive} \cap \text{"happy"})}{P(\text{"happy"})}$$

$$P(\text{"happy"} \mid \text{Positive}) = \frac{P(\text{"happy"} \cap \text{Positive})}{P(\text{Positive})}$$

- ☒ $P(\text{Positive} \mid \text{"happy"}) = P(\text{"happy"} \mid \text{Positive}) \times \frac{P(\text{Positive})}{P(\text{"happy"})}$
☐ $P(\text{Positive} \mid \text{"happy"}) = P(\text{"happy"} \mid \text{Positive}) \times \frac{P(\text{"happy"})}{P(\text{Positive})}$
☐ $P(\text{Positive} \cap \text{"happy"}) = P(\text{"happy"} \mid \text{Positive}) \times \frac{P(\text{Positive})}{P(\text{"happy"})}$
☐ $P(\text{Positive} \cap \text{"happy"}) = P(\text{"happy"} \mid \text{Positive}) \times \frac{P(\text{"happy"})}{P(\text{Positive})}$

✔ Correct

Yes, that is the correct answer.

4. Bayes rule is defined as

1 / 1 point

- ☒ $P(X \mid Y) = P(Y \mid X) \times \frac{P(X)}{P(Y)}$
☐ $P(X \mid Y) = P(Y \mid X) \times \frac{P(Y)}{P(X)}$
☐ $P(X \mid Y) = P(X \mid Y) \times \frac{P(X)}{P(Y)}$
☐ $P(X \mid Y) = P(Y \mid X) \times \frac{P(X)}{P(Y \mid X)}$

✔ Correct

Yes.

5. Suppose that in your dataset, 25% of the positive tweets contain the word 'happy'. You also know that a total of 13% of the tweets in your dataset contain the word 'happy', and that 40% of the total number of tweets are positive. You observe the tweet: 'happy to learn NLP'. What is the probability that this tweet is positive? (Please, round your answer up to two decimal places. Remember that $0.578 = 0.58$ and $0.572 = 0.57$)

1 / 1 point

0.77

✔ Correct

That's right. You just applied Bayes' rule.

6. The log likelihood for a certain word w_i is defined as:

1 / 1 point

$$\log\left(\frac{P(w_i|pos)}{P(w_i|neg)}\right).$$

- ☒ Positive numbers imply that the word is positive.

✔ Correct

- ☐ Positive numbers imply that the word is negative.

- ☒ Negative numbers imply that the word is negative.

✔ Correct

- ☐ Negative numbers imply that the word is positive.

7. The log likelihood mentioned in lecture, which is the log of the ratio between two probabilities is bounded between

1 / 1 point


- ☐ -1 and 1
☒ $-\infty$ and ∞
☐ 0 and ∞
☐ 0 and 1

 **Correct**
 Yes!

8. When implementing naive Bayes, in which order should the following steps be implemented.

1 / 1 point


- ☒
 1. Get or annotate a dataset with positive and negative tweets
 2. Preprocess the tweets: `process_tweet(tweet)` →
 3. Compute `freq(w, class)`
 4. Get $P(w \mid \text{pos})$, $P(w \mid \text{neg})$
 5. Get $\lambda(w)$
 6. Compute $\text{logprior} = \log(P(\text{pos}) / P(\text{neg}))$
- ☐
 1. Get or annotate a dataset with positive and negative tweets
 2. Preprocess the tweets: `process_tweet(tweet)` →
 3. Compute `freq(w, class)`
 4. Get $\lambda(w)$
 5. Get $P(w \mid \text{pos})$, $P(w \mid \text{neg})$
 6. Compute $\text{logprior} = \log(P(\text{pos}) / P(\text{neg}))$
- ☐
 1. Get or annotate a dataset with positive and negative tweets
 2. Compute `freq(w, class)`
 3. Preprocess the tweets: `process_tweet(tweet)` →
 4. Get $P(w \mid \text{pos})$, $P(w \mid \text{neg})$
 5. Get $\lambda(w)$
 6. Compute $\text{logprior} = \log(P(\text{pos}) / P(\text{neg}))$
- ☐
 1. Get or annotate a dataset with positive and negative tweets
 2. Compute `freq(w, class)`
 3. Preprocess the tweets: `process_tweet(tweet)` →
 4. Compute $\text{logprior} = \log(P(\text{pos}) / P(\text{neg}))$
 5. Get $P(w \mid \text{pos})$, $P(w \mid \text{neg})$
 6. Get $\lambda(w)$

 **Correct**
 Yes, that is correct.

9. To test naive bayes model, which of the following are required?

1 / 1 point


- ☒ $X_{\text{val}}, Y_{\text{val}}, \lambda, \text{logprior}$
☐ $X_{\text{val}}, Y_{\text{val}}, \text{logprior}$
☐ $X_{\text{val}}, \lambda, \text{logprior}$
☐ $Y_{\text{val}}, \lambda, \text{logprior}$

 **Correct**
 This is correct.

10. Which of the following is NOT an application of naive Bayes?

1 / 1 point

- ☐ Sentiment Analysis
☐ Author identification
☐ Information retrieval
☐ Word disambiguation
☒ Numerical predictions

 **Correct**
 This is correct.