


# MATH 3341: Introduction to Scientific Computing Lab

Melissa Butler

University of Wyoming


November 15, 2021



The background of the slide features a large, faint watermark of the University of Wyoming seal. The seal is circular with a rope-like border. Inside the border, the words "UNIVERSITY OF WY" are at the top, "EQUALITY" is in the center, and "1886" is at the bottom. There are also some stars and a small figure in the center.

## Lab 12: Romberg Integration



The background of the slide features a large, faint watermark of the University of Wyoming seal. The seal is circular with a rope-like border. Inside the border, the words "UNIVERSITY OF WYOMING" are at the top, "EQUALITY" is in the center, and "1886" is at the bottom. A book is depicted in the center of the seal.

## Romberg Integration



# Composite Trapezoidal Rule

- Composite Trapezoidal rule for approximating the integral of a function  $f(x)$  on an interval  $[a, b]$  using  $m$  subintervals:

$$I = \int_a^b f(x) dx \approx \frac{1}{2} \sum_{j=0}^{m-1} (x_{j+1} - x_j)(y_{j+1} + y_j),$$

where  $a = x_0 < x_1 < \cdots < x_m = b$ ,  $y_j = f(x_j)$ ,  $j = 0, \dots, m$ .



# Composite Trapezoidal Rule

- Composite Trapezoidal rule for approximating the integral of a function  $f(x)$  on an interval  $[a, b]$  using  $m$  subintervals:

$$I = \int_a^b f(x) dx \approx \frac{1}{2} \sum_{j=0}^{m-1} (x_{j+1} - x_j)(y_{j+1} + y_j),$$

where  $a = x_0 < x_1 < \dots < x_m = b$ ,  $y_j = f(x_j)$ ,  $j = 0, \dots, m$ .

- Let  $x_{j+1} - x_j = h = (b - a)/m$ ,  $j = 0, 1, \dots, m - 1$ .

$$I \approx \frac{1}{2} \sum_{j=0}^{m-1} h[f(x_{j+1}) + f(x_j)] = \frac{h}{2} \left[ f(a) + f(b) + 2 \sum_{j=1}^{m-1} f(x_j) \right],$$


where  $x_j = x_0 + jh = a + jh$  for each  $j = 0, 1, \dots, m$ .



# Composite Trapezoidal Rule

- Composite Trapezoidal Rule using  $m$  subintervals:

$$I \approx \frac{h}{2} \left[ f(a) + f(b) + 2 \sum_{j=1}^{m-1} f(a + jh) \right].$$



# Composite Trapezoidal Rule

- Composite Trapezoidal Rule using  $m$  subintervals:

$$I \approx \frac{h}{2} \left[ f(a) + f(b) + 2 \sum_{j=1}^{m-1} f(a + jh) \right].$$

- Finding approximation for

$m_1 = 1, m_2 = 2, m_3 = 4, \dots, m_n = 2^{n-1}$  for  $n \in \mathbb{N}$ . The corresponding step size  $h_k$  for each  $m_k$  is then given by  $h_k = (b - a)/m_k = (b - a)/2^{k-1}$ . The composite trapezoidal rule then becomes

$$I = \int_a^b f(x) dx \approx \frac{h_k}{2} \left[ f(a) + f(b) + 2 \sum_{j=1}^{2^{k-1}-1} f(a + jh_k) \right].$$



Here we'll use the notation  $R_{k,1}$  to denote the portion used for the trapezoidal approximation. In other words,

$$\begin{aligned}R_{1,1} &= \frac{h_1}{2} [f(a) + f(b)] = \frac{b-a}{2} [f(a) + f(b)], \\R_{2,1} &= \frac{h_2}{2} [f(a) + f(b) + 2f(a+h_2)] \\&= \frac{1}{2} \frac{h_1}{2} [f(a) + f(b) + 2f(a+h_2)] \\&= \frac{1}{2} \left\{ \frac{h_1}{2} [f(a) + f(b)] + 2 \frac{h_1}{2} f(a+h_2) \right\} \\&= \frac{1}{2} [R_{1,1} + h_1 f(a+h_2)], \\R_{3,1} &= \frac{1}{2} \{ R_{2,1} + h_2 [f(a+h_3) + f(a+3h_3)] \}.\end{aligned}$$





This leads to the Trapezoidal rule in the general form

$$R_{k,1} = \frac{1}{2} \left[ R_{k-1,1} + h_{k-1} \sum_{j=1}^{2^{k-2}} f(a + (2j-1)h_k) \right] \quad \text{for } k = 2, 3, \dots, n.$$

This method converges very slowly on its own. A technique called Richardson's Extrapolation is applied to speed convergence. Essentially, this performs a method of averaging previously calculated entries to obtain the next entry in the table. This is given in general form

$$R_{k,j} = R_{k,j-1} + \frac{R_{k,j-1} - R_{k-1,j-1}}{4^{j-1} - 1}.$$



This method will give us the following entries of  $R$  in a tabular format. The number of rows is determined by the value that we desire.

$$\begin{bmatrix} R_{1,1} & & & & \\ R_{2,1} & R_{2,2} & & & \\ R_{3,1} & R_{3,2} & R_{3,3} & & \\ R_{4,1} & R_{4,2} & R_{4,3} & R_{4,4} & \\ \vdots & \vdots & \vdots & \vdots & \ddots \\ R_{n,1} & R_{n,2} & R_{n,3} & R_{n,4} & \cdots & R_{n,n} \end{bmatrix}$$



# Algorithm

---

## Algorithm 1: Romberg Integration

---

**Function** romberg( $f, a, b, n$ ):

$h \leftarrow b - a$ ;

$R_{1,1} \leftarrow [f(a) + f(b)] \cdot h/2$ ;

**for**  $k \leftarrow 2$  **to**  $n$  **do**

$R_{k,1} \leftarrow \frac{1}{2} \left[ R_{k-1,1} + h \sum_{j=1}^{2^{k-2}} f(a + (2j-1) \cdot h/2) \right]$ ;

**for**  $j \leftarrow 2$  **to**  $k$  **do**

$R_{k,j} \leftarrow R_{k,j-1} + \frac{R_{k,j-1} - R_{k-1,j-1}}{4^{j-1} - 1}$ ;

**end**

$h \leftarrow h/2$ ;

**end**

**return**  $[R_{1,1}, R_{2,2}, R_{3,3}, \dots, R_{n,n}]$ ;

**end**

