

## 0.1 Pair Selection


It is important to obtain true particle pairs in the analysis. In particular, contamination from pairs constructed with split or merged tracks, and pairs sharing daughters, can introduce an artificial signal into the correlation function, obscuring the actual physics.

### 1. Shared Daughter Cut for Pairs


- (a) V0-V0 Pairs (i.e.  $\Lambda(\bar{\Lambda})K_S^0$  analyses)
  - Remove all pairs which share a daughter
    - Ex.  $\Lambda$  and  $K_S^0$  particles which share a  $\pi^-$  daughter are not included
- (b) V0-Track Pairs (i.e.  $\Lambda(\bar{\Lambda})K^\pm$  analyses)
  - Remove pairs if Track is also used as a daughter of the V0
    - In these analyses, this could only occur if, for instance, a  $K$  is misidentified as a  $\pi$  or  $p$  in the V0 reconstruction
- (c)  $\Xi$ -Track Pairs
  - Remove pairs if Track is also used as a daughter of the  $\Xi$ 
    - In these analyses, this could only occur if, for instance, a  $K$  is misidentified as a  $\pi$  or  $p$  in the V0 reconstruction, or misidentified as bachelor  $\pi$ .
  - Remove pair if bachelor  $\pi$  is also a daughter of the  $\Lambda$ 
    - This is not a pair cut, but is included here because this cut occurs in the AliFemtoXiTrackPairCut class

### 2. Average Separation Cuts


- Used to cut out splitting and merging effects
- The motivation for these cuts can be seen in Figures 1, 2, and 3, in which average separation correlation functions are presented
- (a)  $\Lambda(\bar{\Lambda})K_S^0$  Analyses
  - Average separation  $> 6.0$  cm for like charge sign daughters
    - ex.  $p$  daughter of  $\Lambda$  and  $\pi^+$  daughter of  $K_S^0$
  - No cut for unlike-sign daughters
- (b)  $\Lambda(\bar{\Lambda})K^\pm$  Analyses
  - Average Separation  $> 8.0$  cm for daughter of  $\Lambda(\bar{\Lambda})$  sharing charge sign of  $K^\pm$ 
    - ex. in  $\Lambda K^+$  analysis,  $p$  daughter of  $\Lambda$  with  $K^+$
  - No cut for unlike signs
- (c)  $\Xi(\bar{\Xi})K^\pm$  Analyses
  - Average Separation  $> 8.0$  cm for any daughter of  $\Xi$  sharing charge sign of  $K^\pm$ 
    - ex. in  $\Xi^- K^-$  analysis,  $\pi^-$  daughter of  $\Lambda$  daughter with  $K^-$ , and bachelor  $\pi^-$  daughter with  $K^-$
  - No cut for unlike signs



**Fig. 1:** Average separation (cm) correlation functions of  $\Lambda(\bar{\Lambda})$  and  $K_S^0$  Daughters. Only like-sign daughter pairs are shown (the distributions for unlike-signs were found to be flat). The title of each subfigure shows the daughter pair, as well as the mother of each daughter (in “()”), ex. top left is p from  $\Lambda$  with  $\pi^+$  from  $K_S^0$ .



**Fig. 2:** Average separation (cm) correlation functions of  $\Lambda(\bar{\Lambda})$  Daughter and  $K^\pm$ . Only like-sign pairs are shown (unlike-signs were flat). In the subfigure titles, the particles in “()” represent the mothers, ex. top left is p from  $\Lambda$  with  $K^+$ .



**Fig. 3:** Average separation (cm) correlation functions of  $\Xi$  Daughter and  $K^\pm$ . In the subfigure titles, the particles in “()” represent the mothers, ex. top left is p from  $\Lambda$  from  $\Xi^-$  with  $K^+$ .