

Contents

1	Introduction	3
2	Data Sample and Software	3
2.1	Data Sample	3
2.2	Software	3
3	Data Selection	3
3.1	Event Selection and Mixing	3
3.2	K^\pm Track Selection	4
3.3	V0 Selection	4
3.3.1	Λ Reconstruction	4
3.3.2	K_S^0 Reconstruction	6
3.4	Cascade Reconstruction	9
3.5	Pair Selection	9
4	Correlation Functions	10
5	Fitting	10
5.1	Model: Lambda-Kaon	10
5.2	Model: Cascade-Kaon	11
5.3	Momentum Resolution Corrections	11
5.4	Residual Correlations	11
6	Systematic Errors	11
7	Results and Discussion	11
8	To Do	11

List of Figures

1	Λ Reconstruction	4
2	K_S^0 Reconstruction	5
3	Short Caption	6
4	K_S^0 contamination in $\Lambda(\bar{\Lambda})$ collection	6
5	K_S^0 contamination in Λ collection	7
6	K_S^0 contamination in $\bar{\Lambda}$ collection	7
7	Λ Purity	8
8	Λ contamination in K_S^0 collection	8
9	Λ contamination in K_S^0 collection	9
10	$\bar{\Lambda}$ contamination in K_S^0 collection	9
11	$\bar{\Lambda}$ contamination in K_S^0 collection	10
12	K_S^0 Purity	10
13	Ξ Reconstruction	11
14	Average Separation $\Lambda(\bar{\Lambda})K_S^0$	14
15	Average Separation $\Lambda(\bar{\Lambda})K^\pm$	14

1 Introduction

This will be my introduction. Remember, Jai suggested to make each sentence a separate line to make changes easier to track in git. Otherwise, git will treat an entire paragraph as a single line!

And a new paragraph begins with an empty line.

2 Data Sample and Software

2.1 Data Sample

Talk about the data sample

Run list: 170593, 170572, 170388, 170387, 170315, 170313, 170312, 170311, 170309, 170308, 170306, 170270, 170269, 170268, 170230, 170228, 170207, 170204, 170203, 170193, 170163, 170159, 170155, 170091, 170089, 170088, 170085, 170084, 170083, 170081, 170040, 170027, 169965, 169923, 169859, 169858, 169855, 169846, 169838, 169837, 169835, 169591, 169590, 169588, 169587, 169586, 169557, 169555, 169554, 169553, 169550, 169515, 169512, 169506, 169504, 169498, 169475, 169420, 169419, 169418, 169417, 169415, 169411, 169238, 169167, 169160, 169156, 169148, 169145, 169144, 169138, 169099, 169094, 169091, 169045, 169044, 169040, 169035, 168992, 168988, 168826, 168777, 168514, 168512, 168511, 168467, 168464, 168460, 168458, 168362, 168361, 168342, 168341, 168325, 168322, 168311, 168310, 168115, 168108, 168107, 168105, 168076, 168069, 167988, 167987, 167985, 167920, 167915

2.2 Software

Talk about the software

3 Data Selection

3.1 Event Selection and Mixing

The events used in this study were selected with the following criteria:

- Triggers
 - minimum bias (kMB)
 - central (kCentral)
 - semi-central (kSemiCentral)
- z-position of reconstructed event vertex must be within 10 cm of the center of the ALICE detector
- the event must contain at least one particle of each type from the pair of interest

The event mixing was handled by the AliFemtoVertexMultAnalysis class, which only mixes events with like vertex position and centrality. The following criteria were used for event mixing:

- Number of events to mix = 5
- Vertex position bin width = 2 cm
- Centrality bin width = 5

The AliFemtoEventReaderAODChain class is used to read the events. Event flattening is not currently used. FilterBit(7). The centrality is determined by the "V0M" method of AliCentrality, set by calling AliFemtoEventReaderAOD::SetUseMultiplicity(kCentrality). I utilize the SetPrimaryVertexCorrectionTPCPoints switch, which causes the reader to shift all TPC points to be relative to the event vertex.

3.2 K^\pm Track Selection

Talk about how charged kaons are identified

3.3 V0 Selection

Λ ($\bar{\Lambda}$) and K_S^0 are neutral particles which cannot be directly detected, but must instead be reconstructed through detection of their decay products, or daughters. In general, particles which are topologically reconstructed in this fashion are called V0 particles. The class AliFemtoV0TrackCutNSigmaFilter (which is an extension of AliFemtoV0TrackCut) is used to reconstruct the V0s.




Fig. 1: Λ Reconstruction

3.3.1 Λ Reconstruction

The following cuts were used to select good Λ ($\bar{\Lambda}$) candidates:

1. Cuts Common to Both Daughters




Fig. 2: K_S^0 Reconstruction

- (a) $|\eta| < 0.8$
- (b) `SetTPCnclsDaughters(80)`
- (c) `SetStatusDaughters(AliESDtrack::kTPCrefic)`
- (d) `SetMaxDcaV0Daughters(0.4)`

2. Pion Specific Daughter Cuts

- (a) $p_T > 0.16$
- (b) DCA to prim vertex > 0.3

3. Proton Specific Daughter Cuts

- (a) $p_T >$
 - 0.5 (p)
 - 0.3 (\bar{p})
- (b) DCA to prim vertex > 0.1

4. Lambda Cuts

- (a) $|\eta| < 0.8$




Fig. 3: Long caption

- (b) $p_T > 0.4$
- (c) $|m_{inv} - m_{PDG}| < 3.8 \text{ MeV}$
- (d) Cosine of pointing angle > 0.9993
- (e) OnFlyStatus = false
- (f) Decay Length $< 60 \text{ cm}$




Fig. 4: Mass assuming K_S^0 -hypothesis for V0 candidates passing all Λ ($\bar{\Lambda}$) cuts, i.e. assume the daughters are $\pi^+\pi^-$ instead of $p^+\pi^-$ ($\pi^+\bar{p}^-$). The slight peak around $m_{inv} = 0.5 \text{ GeV}/c^2$ likely contains misidentified K_S^0 particles in our Λ collection. If one simply cuts out the entire peak, good Λ particles will be lost. Ideally, the Λ selection and K_S^0 misidentification cuts are selected such that the peak is removed from this plot while leaving the distribution continuous.

3.3.2 K_S^0 Reconstruction

The following cuts were used to select good K_S^0 candidates:




Fig. 5: Mass assuming K_S^0 -hypothesis for V0 candidates passing all Λ cuts, i.e. assume the daughters are $\pi^+\pi^-$ instead of $p^+\pi^-$. The slight peak around $m_{inv} = 0.5 \text{ GeV}/c^2$ likely contains misidentified K_S^0 particles in our Λ collection. If one simply cuts out the entire peak, good Λ particles will be lost. Ideally, the Λ selection and K_S^0 misidentification cuts are selected such that the peak is removed from this plot while leaving the distribution continuous.




Fig. 6: Mass assuming K_S^0 -hypothesis for V0 candidates passing all $\bar{\Lambda}$ cuts, i.e. assume the daughters are $\pi^+\pi^-$ instead of $\pi^+\bar{p}^-$. Similar to Figure 5

1. Pion Daughter Cuts

- (a) $|\eta| < 0.8$
- (b) SetTPCncIsDaughters(80)
- (c) SetStatusDaughters(AliESDtrack::kTPCrefic)
- (d) SetMaxDcaV0Daughters(0.3)
- (e) $p_T > 0.15$
- (f) DCA to prim vertex > 0.3

2. K_S^0 Cuts

- (a) $|\eta| < 0.8$




Fig. 7: Λ Purity

- (b) $p_T > 0.2$
- (c) $m_{PDG} - 13.677 \text{ MeV} < m_{inv} < m_{PDG} + 2.0323 \text{ MeV}$
- (d) Cosine of pointing angle > 0.9993
- (e) OnFlyStatus = false
- (f) Decay Length $< 30 \text{ cm}$




Fig. 8: Mass assuming Λ -hypothesis for V0 candidates passing all K_S^0 cuts, i.e. assume the daughters are $p^+\pi^-$ instead of $\pi^+\pi^-$. The peak around $m_{inv} = 1.115 \text{ GeV}/c^2$ likely contains misidentified Λ particles in our K_S^0 collection. If one simply cuts out the entire peak, some good K_S^0 particles will be lost. Ideally, the K_S^0 selection and $\Lambda(\bar{\Lambda})$ misidentification cuts can be selected such that the peak is removed from this plot while leaving the distribution continuous.

As can be seen in Figures 9 and 11, some misidentified Λ and $\bar{\Lambda}$ particles contaminate our K_S^0 sample. To attempt to remove these contaminations without throwing away good K_S^0 particles, the following misidentification cuts are imposed; a K_S^0 candidate is rejected if all of the following criteria are satisfied:

$$- |m_{inv, \Lambda(\bar{\Lambda}) \text{ Hypothesis}} - m_{PDG, \Lambda(\bar{\Lambda})}| < 9.0 \text{ MeV}/c^2$$




Fig. 9: Mass assuming Λ -hypothesis for V0 candidates passing all K_S^0 cuts, i.e. assume the daughters are $p^+\pi^-$ instead of $\pi^+\pi^-$. The peak around $m_{inv} = 1.115 \text{ GeV}/c^2$ likely contains misidentified Λ particles in our K_S^0 collection. If one simply cuts out the entire peak, some good K_S^0 particles will be lost. Ideally, the K_S^0 selection and $\Lambda(\bar{\Lambda})$ misidentification cuts can be selected such that the peak is removed from this plot while leaving the distribution continuous.




Fig. 10: Mass assuming $\bar{\Lambda}$ -hypothesis for V0 candidates passing all K_S^0 cuts, i.e. assume the daughters are $\pi^+\bar{p}^-$ instead of $\pi^+\pi^-$. Similar to Figure 9

- Positive daughter passes $p^+(\pi^+)$ daughter cut implemented for $\Lambda(\bar{\Lambda})$ reconstruction
- Negative daughter passes $\pi^-(\bar{p}^-)$ daughter cut implemented by $\Lambda(\bar{\Lambda})$ reconstruction

3.4 Cascade Reconstruction

Talk about reconstruction cascades

3.5 Pair Selection

Some general remarks on forming pairs

It is important to obtain true particle pairs in the analysis. In particular, contamination from pairs con-




Fig. 11: Mass assuming $\bar{\Lambda}$ -hypothesis for V0 candidates passing all K_S^0 cuts, i.e. assume the daughters are $\pi^+\bar{p}^-$ instead of $\pi^+\pi^-$. Similar to Figure 9




Fig. 12: K_S^0 Purity

structured with split or merged tracks can introduce an artificial signal into the correlation function, obscuring the actual physics.

4 Correlation Functions

General remarks about formaton of correlation functions and what information they provide.

5 Fitting

This section will include the Lednicky model and the method used to fit the Cascade study. It will also include momentum resolution, residual correlations, and any other aspects to obtain a good fit

5.1 Model: Lambda-Kaon

Talk about Lednicky model




Fig. 13: Ξ Reconstruction

5.2 Model: Cascade-Kaon

Talk about model

5.3 Momentum Resolution Corrections

Talk about Momentum resolution corrections

$$C_{fit}(k_{Rec}^*) = \frac{\sum_{k_{True}^*} M_{k_{Rec}^*, k_{True}^*} C_{fit}(k_{True}^*)}{\sum_{k_{True}^*} M_{k_{Rec}^*, k_{True}^*}} \quad (1)$$

5.4 Residual Correlations

Talk about Lednický model

6 Systematic Errors

This study is currently ongoing. See Table 1.

7 Results and Discussion

8 To Do

DCA $\Lambda(\bar{\Lambda})$

Pair Type	Centrality	p-value	
		0.4 vs 0.5 mm	0.5 vs 0.6 mm
ΛK_S^0	0-10%	0.36	0.05
	10-30%	0.10	0.37
	30-50%	0.27	6.7e-8
$\bar{\Lambda} K_S^0$	0-10%	0.08	3.2e-4
	10-30%	0.15	0.31
	30-50%	3.7e-3	7.1e-3

Table 1: DCA V0 LamK0 captionDCA K_S^0

Pair Type	Centrality	p-value	
		0.2 vs 0.3 mm	0.3 vs 0.4 mm
ΛK_S^0	0-10%	0.32	0.76
	10-30%	2.1e-3	0.13
	30-50%	0.04	0.06
$\bar{\Lambda} K_S^0$	0-10%	2.8e-7	1.3e-4
	10-30%	0.22	0.62
	30-50%	0.76	0.02

Table 2: DCA V0 LamK0 captionDCA $\Lambda(\bar{\Lambda})$

Pair Type	Centrality	p-value	
		0.4 vs 0.5 mm	0.5 vs 0.6 mm
ΛK^+	0-10%	0.01	3.2e-5
	10-30%	5.9e-3	0.22
	30-50%	0.85	0.84
$\bar{\Lambda} K^-$	0-10%	0.15	0.03
	10-30%	3.1e-4	0.42
	30-50%	7.2e-3	0.42
ΛK^-	0-10%	0.35	0.05
	10-30%	1.4e-5	5.6e-3
	30-50%	0.05	0.70
$\bar{\Lambda} K^+$	0-10%	0.84	0.16
	10-30%	0.16	3.3e-3
	30-50%	2.5e-4	0.20

Table 3: DCA V0 LamKch captionDCA $\Lambda(\bar{\Lambda})$ Daughters

Pair Type	Centrality	p-value	
		0.3 vs 0.4 mm	0.4 vs 0.5 mm
ΛK_S^0	0-10%	0.39	0.51
	10-30%	0.30	0.84
	30-50%	1.3e-38	8.7e-3
$\bar{\Lambda} K_S^0$	0-10%	0.35	0.07
	10-30%	0.07	0.13
	30-50%	0.44	0.01

Table 4: DCA $\Lambda(\bar{\Lambda})$ Daughters LamK0 caption

DCA K_S^0 Daughters

Pair Type	Centrality	p-value	
		0.2 vs 0.3 mm	0.3 vs 0.4 mm
ΛK_S^0	0-10%	0.08	0.29
	10-30%	0.01	0.47
	30-50%	6.6e-3	0.82
$\bar{\Lambda} K_S^0$	0-10%	0.38	0.44
	10-30%	0.13	0.25
	30-50%	0.06	0.53

Table 5: DCA K_S^0 Daughters LamK0 captionDCA $\Lambda(\bar{\Lambda})$ Daughters

Pair Type	Centrality	p-value	
		0.3 vs 0.4 mm	0.4 vs 0.5 mm
ΛK^+	0-10%	0.79	0.06
	10-30%	0.10	0.60
	30-50%	8.4e-3	0.25
$\bar{\Lambda} K^-$	0-10%	2.4e-4	0.63
	10-30%	0.06	3.3e-4
	30-50%	0.03	0.04
ΛK^-	0-10%	0.70	0.40
	10-30%	0.94	0.04
	30-50%	0.05	9.5e-5
$\bar{\Lambda} K^+$	0-10%	0.09	0.04
	10-30%	0.10	0.17
	30-50%	0.10	0.43

Table 6: DCA $\Lambda(\bar{\Lambda})$ Daughters LamKch caption




Fig. 14: Average Separation $\Lambda(\bar{\Lambda})K_S^0$




Fig. 15: Average Separation $\Lambda(\bar{\Lambda})K^\pm$