

Contents

1	Introduction	4
2	Data Sample and Software	4
2.1	Data Sample	4
2.2	Software	4
3	Data Selection	4
3.1	Event Selection and Mixing	4
3.2	K^\pm Track Selection	5
3.3	V0 Selection	6
3.3.1	Λ Reconstruction	7
3.3.2	K_S^0 Reconstruction	9
3.4	Cascade Reconstruction	10
3.5	Pair Selection	13
4	Correlation Functions	14
5	Fitting	15
5.1	Model: ΛK_S^0 , ΛK^\pm , $\Xi^{ch} K_S^0$	15
5.2	Model: $\Xi^{ch} K^{ch}$	18
5.3	Momentum Resolution Corrections	21
5.4	Residual Correlations	23
6	Systematic Errors	26
6.1	Systematic Errors: ΛK_S^0	27
6.1.1	Particle and Pair Cuts	27
6.1.2	Non-Flat Background	28
6.1.3	Fit Range	28
6.2	Systematic Errors: ΛK^\pm	28
6.2.1	Particle and Pair Cuts	28
6.2.2	Non-Flat Background	28
6.2.3	Fit Range	28
6.3	Systematic Errors: ΞK^\pm	28
6.3.1	Particle and Pair Cuts	28

7 Results and Discussion	29
7.1 Results: ΛK_S^0 and ΛK^\pm	29
7.2 Results: ΞK^\pm	38
8 To Do	40
9 Additional Figures	40
9.1 Residuals	40

List of Figures

1	V0 Reconstruction	7
2	K_S^0 contamination in $\Lambda(\bar{\Lambda})$ collection	8
3	Λ and $\bar{\Lambda}$ Purity	9
4	$\Lambda(\bar{\Lambda})$ contamination in K_S^0 collection	11
5	K_S^0 Purity	12
6	Ξ Reconstruction	13
7	$\Xi^-(\bar{\Xi}^+)$ Purity	13
8	Average Separation of $\Lambda(\bar{\Lambda})$ and K_S^0 Daughters	15
9	Average Separation of $\Lambda(\bar{\Lambda})$ Daughter and K^\pm	16
10	Average Separation of Ξ Daughters and K^\pm	17
11	$\Lambda(\bar{\Lambda})K_S^0$ Correlation Functions	18
12	ΛK^+ and $\bar{\Lambda} K^-$ Correlation Functions	19
13	ΛK^- and $\bar{\Lambda} K^+$ Correlation Functions	20
14	Correlation Functions: ΛK^+ vs ΛK^- for 0-10% Centrality	21
15	Momentum Resolution: Sample k_{True}^* vs. k_{Rec}^*	24
16	Particle Contaminations Visible in k_{True}^* vs. k_{Rec}^*	25
17	Transform Matrices for ΛK^+ Analysis	26
18	Transform Matrices for $\bar{\Lambda} K^+$ Analysis	27
19	$\Lambda K_S^0(\bar{\Lambda} K_S^0)$ Fits	30
20	$\Lambda K_S^0(\bar{\Lambda} K_S^0)$ Fits (Wide Range)	31
21	$\Lambda K^+(\bar{\Lambda} K^-)$ Fits	32
22	$\Lambda K^+(\bar{\Lambda} K^-)$ Fits (Wide Range)	33
23	$\Lambda K^-(\bar{\Lambda} K^+)$ Fits	34
24	$\Lambda K^-(\bar{\Lambda} K^+)$ Fits (Wide Range)	35
25	m_T Scaling of Radii	38
26	ΞK^\pm Results	38
27	ΞK^\pm Data with Coulomb-Only Bands, 0-10% Centrality	39
28	Effect of Strong Force Inclusion on Coulomb-Only Curve for ΞK^\pm systems	41
29	ΞK^\pm Global Coulomb-Only Fit (Set 1)	42
30	ΞK^\pm Global Coulomb-Only Fit (Set 2)	43
31	$\Xi^- K^+$ Coulomb-Only Fit	43
32	$\Xi^- K^-$ Coulomb-Only Fit	44

33	Residuals: $\Sigma^0 K^+$ to ΛK^+ (0-10% Centrality)	44
34	Residuals: $\Xi^0 K^+$ to ΛK^+ (0-10% Centrality)	45
35	Residuals: $\Xi^- K^+$ to ΛK^+ (0-10% Centrality)	45
36	Residuals: $\Sigma^{*+} K^+$ to ΛK^+ (0-10% Centrality)	46
37	Residuals: $\Sigma^{*-} K^+$ to ΛK^+ (0-10% Centrality)	46
38	Residuals: $\Sigma^{*0} K^+$ to ΛK^+ (0-10% Centrality)	47
39	Residuals: ΛK^{*0} to ΛK^+ (0-10% Centrality)	47
40	Residuals: $\Sigma^0 K^{*0}$ to ΛK^+ (0-10% Centrality)	48
41	Residuals: $\Xi^0 K^{*0}$ to ΛK^+ (0-10% Centrality)	48
42	Residuals: $\Xi^- K^{*0}$ to ΛK^+ (0-10% Centrality)	49

1 1 Introduction

2 We present results from a femtoscopic analysis of Lambda-Kaon correlations in Pb-Pb collisions at $\sqrt{s_{NN}}$
 3 = 2.76 TeV by the ALICE experiment at the LHC. All pair combinations of Λ and $\bar{\Lambda}$ with K^+ , K^- and
 4 K_S^0 are analyzed. The femtoscopic correlations are the result of strong final-state interactions, and are
 5 fit with a parametrization based on a model by R. Lednický and V. L. Lyuboshitz [?]. This allows us to
 6 both characterize the emission source and measure the scattering parameters for the particle pairs. We
 7 observe a large difference in the Λ - K^+ ($\bar{\Lambda}$ - K^-) and Λ - K^- ($\bar{\Lambda}$ - K^+) correlations in pairs with low relative
 8 momenta ($k^* \lesssim 100$ MeV). Additionally, the average of the Λ - K^+ ($\bar{\Lambda}$ - K^-) and Λ - K^- ($\bar{\Lambda}$ - K^+) correlation
 9 functions is consistent with our Λ - K_S^0 ($\bar{\Lambda}$ - K_S^0) measurement. The results suggest an effect arising from
 10 different quark-antiquark interactions in the pairs, i.e. $s\bar{s}$ in Λ - K^+ ($\bar{\Lambda}$ - K^-) and $u\bar{u}$ in Λ - K^- ($\bar{\Lambda}$ - K^+). To
 11 gain further insight into this hypothesis, we currently are conducting a Ξ -K femtoscopic analysis.

12 2 Data Sample and Software

13 2.1 Data Sample

14 The analysis used “pass 2” reconstructed Pb-Pb data from LHC11h (AOD145). The runlist was selected
 15 from runs with global quality tag “1” in the ALICE Run Condition Table. Approximately 40 million
 16 combined central, semi-central, and minimum bias events were analyzed. Runs from both positive (++)
 17 and negative (--) magnetic field polarity settings were used.

18 Run list: 170593, 170572, 170388, 170387, 170315, 170313, 170312, 170311, 170309, 170308, 170306,
 19 170270, 170269, 170268, 170230, 170228, 170207, 170204, 170203, 170193, 170163, 170159, 170155,
 20 170091, 170089, 170088, 170085, 170084, 170083, 170081, 170040, 170027, 169965, 169923, 169859,
 21 169858, 169855, 169846, 169838, 169837, 169835, 169591, 169590, 169588, 169587, 169586, 169557,
 22 169555, 169554, 169553, 169550, 169515, 169512, 169506, 169504, 169498, 169475, 169420, 169419,
 23 169418, 169417, 169415, 169411, 169238, 169167, 169160, 169156, 169148, 169145, 169144, 169138,
 24 169099, 169094, 169091, 169045, 169044, 169040, 169035, 168992, 168988, 168826, 168777, 168514,
 25 168512, 168511, 168467, 168464, 168460, 168458, 168362, 168361, 168342, 168341, 168325, 168322,
 26 168311, 168310, 168315, 168310, 168310, 168310, 168310, 168310, 168310, 168310, 168310, 168310,
 27 167915

28 Analysis was also performed on the LHC12a17a_fix (AOD149) Monte Carlo HIJING events for certain
 29 checks. THERMINATOR2 was also used for certain aspects, such as transform matrices described feed-
 30 down contributions.

31 2.2 Software

32 The analysis was performed on the PWGCF analysis train using AliRoot v5-08-18-1 and AliPhysics
 33 vAN-20161027-1.

34 The main classes utilized include: AliFemtoVertexMultAnalysis, AliFemtoEventCutEstimators, AliFem-
 35 toESDTrackCutNSigmaFilter, AliFemtoV0TrackCutNSigmaFilter, AliFemtoXiTrackCut, AliFemtoV0PairCut,
 36 AliFemtoV0TrackPairCut, AliFemtoXiTrackPairCut, and AliFemtoAnalysisLambdaKaon. All of these
 37 classes are contained in /AliPhysics/PWGCF/FEMTOSCOPY/AliFemto and .../AliFemtoUser.

38 3 Data Selection

39 3.1 Event Selection and Mixing

40 The events used in this study were selected with the class AliFemtoEventCutEstimators according to the
 41 following criteria:

- 42 – Triggers
- 43 – minimum bias (kMB)
- 44 – central (kCentral)
- 45 – semi-central (kSemiCentral)
- 46 – z-position of reconstructed event vertex must be within 10 cm of the center of the ALICE detector
- 47 – the event must contain at least one particle of each type from the pair of interest

48 The event mixing was handled by the AliFemtoVertexMultAnalysis class, which only mixes events with
49 like vertex position and centrality. The following criteria were used for event mixing:

- 50 – Number of events to mix = 5
- 51 – Vertex position bin width = 2 cm
- 52 – Centrality bin width = 5

53 The AliFemtoEventReaderAODChain class is used to read the events. Event flattening is not currently
54 used. FilterBit(7). The centrality is determined by the “V0M” method of AliCentrality, set by calling Al-
55 iFemtoEventReaderAOD::SetUseMultiplicity(kCentrality). I utilize the SetPrimaryVertexCorrectionT-
56 PCPoints switch, which causes the reader to shift all TPC points to be relative to the event vertex.

57 3.2 K^\pm Track Selection

58 Charged kaons are identified using the AliFemtoESDTrackCutNSigmaFilter class. The specific cuts used
59 in this analysis are as follows:

60 Track Selection:

- 61 – Kinematic range:
- 62 – $0.14 < p_T < 1.5$
- 63 – $|\eta| < 0.8$
- 64 – FilterBit(7)
- 65 – TPC tracks
- 66 – Track Quality
- 67 – Minimum number of clusters in the TPC (fminTPCncls) = 80
- 68 – Maximum allowed χ^2/N_{DOF} for ITS clusters = 3.0
- 69 – Maximum allowed χ^2/N_{DOF} for TPC clusters = 4.0
- 70 – Primary Particle Selection:
- 71 – Maximum XY impact parameter = 2.4
- 72 – Maximum Z impact parameter = 3.0
- 73 – Remove particles with any kink labels (fRemoveKinks = true)
- 74 – Maximum allowed sigma to primary vertex (fMaxSigmaToVertex) = 3.0

75 K $^\pm$ Identification:

76 – PID Probabilities:

- 77 – K: > 0.2
- 78 – π : < 0.1
- 79 – μ : < 0.8
- 80 – p: < 0.1

81 – Most probable particle type must be Kaon (fMostProbable=3)

82 – TPC and TOF N $_\sigma$ cuts:

- 83 – p < 0.4 GeV/c: N $_{\sigma K,TPC}$ < 2
- 84 – 0.4 < p < 0.45 GeV/c: N $_{\sigma K,TPC}$ < 1
- 85 – 0.45 < p < 0.8 GeV/c: N $_{\sigma K,TPC}$ < 3 & N $_{\sigma K,TOF}$ < 2
- 86 – 0.8 < p < 1.0 GeV/c: N $_{\sigma K,TPC}$ < 3 & N $_{\sigma K,TOF}$ < 1.5
- 87 – p > 1.0 GeV/c: N $_{\sigma K,TPC}$ < 3 & N $_{\sigma K,TOF}$ < 1

88 – Electron Rejection: Reject if N $_{\sigma e^-,TPC}$ < 3

89 – Pion Rejection: Reject if:

- 90 – p < 0.65
 - 91 * if TOF and TPC available: N $_{\sigma \pi,TPC}$ < 3 & N $_{\sigma \pi,TOF}$ < 3
 - 92 * else
 - 93 · p < 0.5: N $_{\sigma \pi,TPC}$ < 3
 - 94 · 0.5 < p < 0.65: N $_{\sigma \pi,TPC}$ < 2
- 95 – 0.65 < p < 1.5: N $_{\sigma \pi,TPC}$ < 5 & N $_{\sigma \pi,TOF}$ < 3
- 96 – p > 1.5: N $_{\sigma \pi,TPC}$ < 5 & N $_{\sigma \pi,TOF}$ < 2

97 The purity of the K $^\pm$ collections was estimated using the MC data, for which the true identity of each
98 reconstructed K $^\pm$ particle is known. Therefore, the purity may be estimated as:

$$Purity(K^\pm) = \frac{N_{true}}{N_{reconstructed}} \quad (1)$$

99 Purity(K $^+$) \approx Purity(K $^-$) \approx 97%

100 3.3 V0 Selection

101 Λ ($\bar{\Lambda}$) and K $_S^0$ are neutral particles which cannot be directly detected, but must instead be reconstructed
102 through detection of their decay products, or daughters. This process is illustrated in Figure 1. In
103 general, particles which are topologically reconstructed in this fashion are called V0 particles. The
104 class AliFemtoV0TrackCutNSigmaFilter (which is an extension of AliFemtoV0TrackCut) is used to
105 reconstruct the V0s.

106 In order to obtain a true and reliable signal, one must ensure good purity of the V0 collection. The purity
107 of the collection is calculated as:

$$Purity = \frac{Signal}{Signal + Background} \quad (2)$$

To obtain both the signal and background, the invariant mass distribution (m_{inv}) of all V0 candidates must be constructed immediately before the final invariant mass cut. Examples of such distributions can be found in Figures 3 and 5. It is vital that this distribution be constructed immediately before the final m_{inv} cut, otherwise it would be impossible to estimate the background. As shown in Figures 3 and 5, the background is fit (with a polynomial) outside of the peak region of interest to obtain an estimate for the background within the region. Within the m_{inv} cut limits, the background is the region below the fit while the signal is the region above the fit.




Fig. 1: V0 Reconstruction

3.3.1 Λ Reconstruction

The following cuts were used to select good Λ ($\bar{\Lambda}$) candidates:

1. Daughter Particle Cuts

(a) Cuts Common to Both Daughters

- i. $|\eta| < 0.8$
- ii. SetTPCnclsDaughters(80)
- iii. SetStatusDaughters(AliESDtrack::kTPCrefic)
- iv. SetMaxDcaV0Daughters(0.4)

(b) Pion Specific Daughter Cuts

- i. $p_T > 0.16$
- ii. DCA to prim vertex > 0.3

(c) Proton Specific Daughter Cuts

- i. $p_T > 0.5(p) [0.3(\bar{p})]$ GeV/c
- ii. DCA to prim vertex > 0.1

129 2. V0 Cuts

- 130 (a) $|\eta| < 0.8$
 131 (b) $p_T > 0.4$
 132 (c) $|m_{inv} - m_{PDG}| < 3.8 \text{ MeV}$
 133 (d) DCA to prim. vertex $< 0.5 \text{ cm}$
 134 (e) Cosine of pointing angle > 0.9993
 135 (f) OnFlyStatus = false
 136 (g) Decay Length $< 60 \text{ cm}$

137 3. Shared Daughter Cut for V0 Collection

- 138 – Iterate through V0 collection to ensure that no daughter is used in more than one V0 candidate




Fig. 2: Mass assuming K_S^0 -hypothesis for V0 candidates passing all Λ (2a) and $\bar{\Lambda}$ (2b) cuts. The “NoMisID” distribution (black triangles) uses the V0 finder without any attempt to remove misidentified K_S^0 . The slight peak in the “NoMisID” distribution around $m_{inv} = 0.5 \text{ GeV}/c^2$ contains misidentified K_S^0 particles in our $\Lambda(\bar{\Lambda})$ collection. “SimpleMisID” (pink squares) simply cuts out the entire peak, which throws away some good Λ and $\bar{\Lambda}$ particles. “MisID_NoM_{inv}Comp” (green squares) uses the misidentification cut outlined in the text, but does not utilize the invariant mass comparison method. “MisID_M_{inv}Comp” (red circles) utilizes the full misidentification methods, and is currently used for this analysis. “N_{pass}/N_{ev}” is the total number of $\Lambda(\bar{\Lambda})$ particles found, normalized by the total number of events. The purity of the collection is also listed.

139 Figure 2a shows the mass assuming K_S^0 hypothesis for the Λ collection, i.e. assume the daughters are
 140 $\pi^+\pi^-$ instead of $\pi^+\bar{p}^-$. Figure 2b is a similar plot, but is for the $\bar{\Lambda}$ collection, i.e. assume the daughters
 141 are $\pi^+\pi^-$ instead of $\pi^+\bar{p}^-$. The K_S^0 contamination is visible, although not profound, in both in the slight
 142 peaks around $m_{inv} = 0.497 \text{ GeV}/c^2$. If one simply cuts out the entire peak, good Λ particles will be
 143 lost. Ideally, the Λ selection and K_S^0 misidentification cuts are selected such that the peak is removed
 144 from this plot while leaving the distribution continuous. To attempt to remove these K_S^0 contaminations
 145 without throwing away good Λ and $\bar{\Lambda}$ particles, the following misidentification cuts are imposed; a $\Lambda(\bar{\Lambda})$
 146 candidate is rejected if all of the following criteria are satisfied:

- 147 – $|m_{inv, K_S^0 \text{ Hypothesis}} - m_{PDG, K_S^0}| < 9.0 \text{ MeV}/c^2$
 148 – Positive and negative daughters pass π daughter cut implemented for K_S^0 reconstruction

$$- \left| m_{inv, K_S^0} Hypothesis - m_{PDG, K_S^0} \right| < \left| m_{inv, \Lambda(\bar{\Lambda})} Hypothesis - m_{PDG, \Lambda(\bar{\Lambda})} \right|$$

Figure 3 shows the invariant mass (M_{inv}) distribution of all $\Lambda(\bar{\Lambda})$ candidates immediately before the final invariant mass cut. These distributions are used to calculate the collection purities. The Λ and $\bar{\Lambda}$ purities are found to be: Purity(Λ) \approx Purity($\bar{\Lambda}$) \approx 95%.




Fig. 3: Invariant mass (M_{inv}) distribution of all Λ (a) and $\bar{\Lambda}$ (b) candidates immediately before the final invariant mass cut. The bottom figures are zoomed to show the background with fit. The vertical green lines represent the M_{inv} cuts used in the analyses, the red vertical lines delineate the region over which the background was fit, and the blue line shows the background fit. These distributions are used to calculate the collection purities, Purity(Λ) \approx Purity($\bar{\Lambda}$) \approx 95%.

3.3.2 K_S^0 Reconstruction

The following cuts were used to select good K_S^0 candidates:

1. Pion Daughter Cuts

- (a) $|\eta| < 0.8$
- (b) SetTPCnclsDaughters(80)
- (c) SetStatusDaughters(AliESDtrack::kTPCrefic)
- (d) SetMaxDcaV0Daughters(0.3)
- (e) $p_T > 0.15$
- (f) DCA to prim vertex > 0.3

2. K_S^0 Cuts

- (a) $|\eta| < 0.8$
- (b) $p_T > 0.2$
- (c) $m_{PDG} - 13.677 \text{ MeV} < m_{inv} < m_{PDG} + 2.0323 \text{ MeV}$
- (d) DCA to prim. vertex $< 0.3 \text{ cm}$
- (e) Cosine of pointing angle > 0.9993
- (f) OnFlyStatus = false
- (g) Decay Length $< 30 \text{ cm}$

170 3. Shared Daughter Cut for V0 Collection

- 171 – Iterate through V0 collection to ensure that no daughter is used in more than one V0 candidate

172 As can be seen in Figure 4, some misidentified Λ and $\bar{\Lambda}$ particles contaminate our K_S^0 sample. Figure
 173 4a shows the mass assuming Λ -hypothesis for the K_S^0 collection, i.e. assume the daughters are $p^+\pi^-$
 174 instead of $\pi^+\pi^-$. Figure 4b is similar, but shows the mass assuming $\bar{\Lambda}$ hypothesis for the collection,
 175 i.e. assume the daughters are $\pi^+\bar{p}^-$ instead of $\pi^+\pi^-$. The Λ contamination can be seen in 4a, and the
 176 $\bar{\Lambda}$ contamination in 4b, in the peaks around $m_{inv} = 1.115 \text{ GeV}/c^2$. Additionally, the $\bar{\Lambda}$ contamination is
 177 visible in Figure 4a, and the Λ contamination visible in Figure 4b, in the region of excess around 1.65
 178 $< m_{inv} < 2.1 \text{ GeV}/c^2$. This is confirmed as the number of misidentified Λ particles in the sharp peak
 179 of Figure 4a (misidentified $\bar{\Lambda}$ particles in the sharp peak of Figure 4b) approximately equals the excess
 180 found in the $1.65 < m_{inv} < 2.1 \text{ GeV}/c^2$ region of Figure 4a (Figure 4b).

181 The peaks around $m_{inv} = 1.115 \text{ GeV}/c^2$ in Figure 4 contain both misidentified Λ ($\bar{\Lambda}$) particles and good
 182 K_S^0 . If one simply cuts out the entire peak, some good K_S^0 particles will be lost. Ideally, the K_S^0 selection
 183 and $\Lambda(\bar{\Lambda})$ misidentification cuts can be selected such that the peak is removed from this plot while leaving
 184 the distribution continuous. To attempt to remove these Λ and $\bar{\Lambda}$ contaminations without throwing away
 185 good K_S^0 particles, the following misidentification cuts are imposed; a K_S^0 candidate is rejected if all of
 186 the following criteria are satisfied (for either Λ or $\bar{\Lambda}$ hypothesis):

- 187 – $|m_{inv, \Lambda(\bar{\Lambda}) \text{ Hypothesis}} - m_{PDG, \Lambda(\bar{\Lambda})}| < 9.0 \text{ MeV}/c^2$
- 188 – Positive daughter passes $p^+(\pi^+)$ daughter cut implemented for $\Lambda(\bar{\Lambda})$ reconstruction
- 189 – Negative daughter passes $\pi^-(\bar{p}^-)$ daughter cut implemented by $\Lambda(\bar{\Lambda})$ reconstruction
- 190 – $|m_{inv, \Lambda(\bar{\Lambda}) \text{ Hypothesis}} - m_{PDG, \Lambda(\bar{\Lambda})}| < |m_{inv, K_S^0 \text{ Hypothesis}} - m_{PDG, K_S^0}|$

191 3.4 Cascade Reconstruction

192 Our motivation for studying ΞK^\pm systems is to hopefully better understand the striking difference in the
 193 ΛK^+ and ΛK^- data at low k^* (Figure 14).


194 The reconstruction of Ξ particles is one step above V0 reconstruction. V0 particles are topologically
 195 reconstructed by searching for the charged daughters' tracks into which they decay. With Ξ particles, we
 196 search for the V0 particle and charged daughter into which the Ξ decays. In the case of Ξ^- , we search
 197 for the Λ (V0) and π^- (track) daughters. We will refer to this π as the “bachelor π ”.

198 The following cuts were used to select good Ξ^- ($\bar{\Xi}^+$) candidates:


199 1. V0 Daughter Reconstruction

200 (a) V0 Daughter Particle Cuts

- 201 i. Cuts Common to Both Daughters
 - 202 A. $|\eta| < 0.8$
 - 203 B. SetTPCnclsDaughters(80)
 - 204 C. SetStatusDaughters(AliESDtrack::kTPCrefic)
 - 205 D. SetMaxDcaV0Daughters(0.4)
- 206 ii. Pion Specific Daughter Cuts
 - 207 A. $p_T > 0.16$



(a) Mass assuming Λ -hypothesis for K_S^0 collection, i.e. assume the daughters are $p^+\pi^-$ instead of $\pi^+\pi^-$.



(b) Mass assuming $\bar{\Lambda}$ -hypothesis for K_S^0 collection, i.e. assume the daughters are $\pi^+\bar{p}^-$ instead of $\pi^+\pi^-$.

Fig. 4: Mass assuming Λ -hypothesis (4a) and $\bar{\Lambda}$ -hypothesis (4b) for K_S^0 collection. The “NoMisID” distribution (black triangles) uses the V0 finder without any attempt to remove misidentified Λ and $\bar{\Lambda}$. The peak in the “NoMisID” distribution around $m_{inv} = 1.115$ GeV/c^2 contains misidentified Λ (4a) and $\bar{\Lambda}$ (4b) particles in our K_S^0 collection. “SimpleMisID” (pink squares) simply cuts out the entire peak, which throws away some good K_S^0 particles. “MisID_NoM_{inv}Comp” (green squares) uses the misidentification cut outlined in the text, but does not utilize the invariant mass comparison method. “MisID_M_{inv}Comp” (red circles) utilizes the full misidentification methods, and is currently used for this analysis. “N_{pass}/N_{ev}” is the total number of K_S^0 particles found, normalized by the total number of events. The purity of the collection is also listed. Also note, the relative excess of the “NoMisID” distribution around $1.65 < m_{inv} < 2.1$ GeV/c^2 shows misidentified $\bar{\Lambda}$ (4a) and Λ (4b) particles in our K_S^0 collection.




Fig. 5: Invariant mass (M_{inv}) distribution of all K_S^0 candidates immediately before the final invariant mass cut. The bottom figure is zoomed to show the background with fit. The vertical green lines represent the M_{inv} cut used in the analyses, the red vertical lines delineate the region over which the background was fit, and the blue line shows the background fit. This distribution is used to calculate the collection purity, $\text{Purity}(K_S^0) \approx 98\%$.

- 210 A. $p_T > 0.5(p) [0.3(\bar{p})] \text{ GeV}/c$
 211 B. DCA to prim vertex > 0.1
- 212 (b) V0 Cuts
 213 i. $|\eta| < 0.8$
 214 ii. $p_T > 0.4 \text{ GeV}/c$
 215 iii. $|m_{inv} - m_{PDG}| < 3.8 \text{ MeV}$
 216 iv. DCA to prim. vertex $> 0.2 \text{ cm}$
 217 v. Cosine of pointing angle to Ξ decay vertex > 0.9993
 218 vi. OnFlyStatus = false
 219 vii. Decay Length $< 60 \text{ cm}$
 220 viii. The misidentification cuts described in Section 3.3.1 are utilized
- 221 2. Bachelor π Cuts
 222 (a) $|\eta| < 0.8$
 223 (b) $p_T < 100 \text{ GeV}/c$
 224 (c) DCA to prim vertex $> 0.1 \text{ cm}$
 225 (d) SetTPCnclsDaughters(70)
 226 (e) SetStatusDaughters(AliESDtrack::kTPCrefic)
- 227 3. Ξ Cuts
 228 (a) $|\eta| < 0.8$
 229 (b) $0.8 < p_T < 100 \text{ GeV}/c$
 230 (c) $|m_{inv} - m_{PDG}| < 3.0 \text{ MeV}$
 231 (d) DCA to prim. vertex $< 0.3 \text{ cm}$
 232 (e) Cosine of pointing angle > 0.9992
- 233 4. Shared Daughter Cut for Ξ Collection
 234 – Iterate through Ξ collection to ensure that no daughter is used in more than one Ξ candidate




Fig. 6: Ξ Reconstruction

235 The purity of our Ξ and $\bar{\Xi}$ collections are calculated just as those of our V0 collections 3.3. Figure 7,
236 which is used to calculate the purity, shows the m_{inv} distribution of our $\Xi(\bar{\Xi})$ candidates just before the
237 final m_{inv} cut. Currently, we have Purity(Ξ^-) \approx 90% and Purity($\bar{\Xi}^+$) \approx 92%.




Fig. 7: $\Xi^-(\bar{\Xi}^+)$ Purity 0-10%: Purity(Ξ^-) \approx 90% and Purity($\bar{\Xi}^+$) \approx 92%.

238 3.5 Pair Selection

239 It is important to obtain true particle pairs in the analysis. In particular, contamination from pairs con-
240 structed with split or merged tracks, and pairs sharing daughters, can introduce an artificial signal into
241 the correlation function, obscuring the actual physics.

242 1. Shared Daughter Cut for Pairs

243 (a) V0-V0 Pairs (i.e. $\Lambda(\bar{\Lambda})K_S^0$ analyses)

- 244 – Remove all pairs which share a daughter
 245 – Ex. Λ and K_S^0 particles which share a π^- daughter are not included
- 246 (b) V0-Track Pairs (i.e. $\Lambda(\bar{\Lambda})K^\pm$ analyses)
 247 – Remove pairs if Track is also used as a daughter of the V0
 248 – In these analyses, this could only occur if, for instance, a K is misidentified as a π
 249 or p in the V0 reconstruction
- 250 (c) Ξ -Track Pairs
 251 – Remove pairs if Track is also used as a daughter of the Ξ
 252 – In these analyses, this could only occur if, for instance, a K is misidentified as a π
 253 or p in the V0 reconstruction, or misidentified as bachelor π .
 254 – Remove pair if bachelor π is also a daughter of the Λ
 255 – This is not a pair cut, but is included here because this cut occurs in the
 256 AliFemto XiTrackPairCut class

257 2. Average Separation Cuts

- 258 – Used to cut out splitting and merging effects
 259 – The motivation for these cuts can be seen in Figures 8, 9, and 10, in which average separation
 260 correlation functions are presented

- 261 (a) $\Lambda(\bar{\Lambda})K_S^0$ Analyses
 262 – Average separation > 6.0 cm for like charge sign daughters
 263 – ex. p daughter of Λ and π^+ daughter of K_S^0
 264 – No cut for unlike-sign daughters
- 265 (b) $\Lambda(\bar{\Lambda})K^\pm$ Analyses
 266 – Average Separation > 8.0 cm for daughter of $\Lambda(\bar{\Lambda})$ sharing charge sign of K^\pm
 267 – ex. in ΛK^+ analysis, p daughter of Λ with K^+
 268 – No cut for unlike signs
- 269 (c) $\Xi(\bar{\Xi})K^\pm$ Analyses
 270 – Average Separation > 8.0 cm for any daughter of Ξ sharing charge sign of K^\pm
 271 – ex. in $\Xi^- K^-$ analysis, π^- daughter of Λ daughter with K^- , and bachelor π^- daugh-
 272 ter with K^-
 273 – No cut for unlike signs

274 4 Correlation Functions

275 This analysis studies the momentum correlations of both Λ -K and Ξ -K pairs using the two-particle cor-
 276 relation function, defined as $C(k^*) = A(k^*)/B(k^*)$, where $A(k^*)$ is the signal distribution, $B(k^*)$ is the
 277 reference (or background) distribution, and k^* is the momentum of one of the particles in the pair rest
 278 frame. In practice, $A(k^*)$ is constructed by binning in k^* pairs from the same event. Ideally, $B(k^*)$ is sim-
 279 ilar to $A(k^*)$ in all respects excluding the presence of femtoscopic correlations [?]; as such, $B(k^*)$ is used
 280 to divide out the phase-space effects, leaving only the femtoscopic effects in the correlation function.

281 In practice, $B(k^*)$ is obtained by forming mixed-event pairs, i.e. particles from a given event are paired
 282 with particles from $N_{mix} (= 5)$ other events, and these pairs are then binned in k^* . In forming the back-
 283 ground distribution, it is important to mix only similar events; mixing events with different phase-spaces




Fig. 8: Average separation (cm) correlation functions of $\Lambda(\bar{\Lambda})$ and K_S^0 Daughters. Only like-sign daughter pairs are shown (the distributions for unlike-signs were found to be flat). The title of each subfigure shows the daughter pair, as well as the mother of each daughter (in “()”), ex. top left is p from Λ with π^+ from K_S^0 .

284 can lead to artificial signals in the correlaton function. Therefore, in this analysis, we mix events with
 285 primary vertices within 2 cm and centralities within 5% of each other. Also note, a vertex correction is
 286 also applied to each event, which essentially recenters the the primary vertices to $z = 0$.

287 This analysis presents correlation functions for three centrality bins (0-10%, 10-30%, and 30-50%),
 288 and is currently pair transverse momentum ($k_T = 0.5|\mathbf{p}_{T,1} + \mathbf{p}_{T,2}|$) integrated (i.e. not binned in k_T).
 289 The correlation functions are constructed separately for the two magnetic field configurations, and are
 290 combined using a weighted average:

$$C_{combined}(k^*) = \frac{\sum_i w_i C_i(k^*)}{\sum_i w_i} \quad (3)$$

291 where the sum runs over the correlation functions to be combined, and the weight, w_i , is the number of
 292 numerator pairs in $C_i(k^*)$. Here, the sum is over the two field configurations.

293 Figures 11, 12, and 13 show the correlation functions for all centalities studied for $\Lambda K_S^0(\bar{\Lambda} K_S^0)$, $\Lambda K^\pm(\bar{\Lambda} K^\mp)$,
 294 and $\Lambda K^\pm(\bar{\Lambda} K^\pm)$, respectively. All were normalized in the range $0.32 < k^* < 0.4$ GeV/c.

295 5 Fitting

296 5.1 Model: ΛK_S^0 , ΛK^\pm , $\Xi^{ch} K_S^0$

297 The two-particle relative momentum correlation function may be written theoretically by the Koonin-
 298 Pratt equation [?, ?]:

$$C(\mathbf{k}^*) = \int S(\mathbf{r}^*) |\Psi_{\mathbf{k}^*}(\mathbf{r}^*)|^2 d^3 \mathbf{r}^* \quad (4)$$




Fig. 9: Average separation (cm) correlation functions of $\Lambda(\bar{\Lambda})$ Daughter and K^\pm . Only like-sign pairs are shown (unlike-signs were flat). In the subfigure titles, the particles in “()” represent the mothers, ex. top left is p from Λ with K^+ .

299 In the absence of Coulomb effects, and assuming a spherically gaussian source of width R , the 1D
 300 femtoscopic correlation function can be calculated analytically using:

$$C(k^*) = 1 + \lambda [C_{QI}(k^*) + C_{FSI}(k^*)] \quad (5)$$

301 C_{QI} describes plane-wave quantum interference:

$$C_{QI}(k^*) = \alpha \exp(-4k^{*2}R^2) \quad (6)$$

302 where $\alpha = (-1)^{2j}/(2j+1)$ for identical particles with spin j , and $\alpha = 0$ for non-identical particles.
 303 Obviously, $\alpha = 0$ for all analyses presented in this note. C_{FSI} describes the s-wave strong final state
 304 interaction between the particles:

$$\begin{aligned} C_{FSI}(k^*) &= (1 + \alpha) \left[\frac{1}{2} \left| \frac{f(k^*)}{R} \right|^2 \left(1 - \frac{d_0}{2\sqrt{\pi}R} \right) + \frac{2\Re f(k^*)}{\sqrt{\pi}R} F_1(2k^*R) - \frac{\Im f(k^*)}{R} F_2(2k^*R) \right] \\ f(k^*) &= \left(\frac{1}{f_0} + \frac{1}{2} d_0 k^{*2} - ik^* \right)^{-1}; \quad F_1(z) = \int_0^z \frac{e^{x^2-z^2}}{z} dx; \quad F_2(z) = \frac{1-e^{-z^2}}{z} \end{aligned} \quad (7)$$

305 where R is the source size, $f(k^*)$ is the s-wave scattering amplitude, f_0 is the complex scattering length,
 306 and d_0 is the effective range of the interaction.

307 The code developed to fit the data is called “LednickýFitter”, and utilizes the ROOT TMinuit implemen-
 308 tation of the MINUIT fitting package. In short, given a function with a number of parameters, the fitter
 309 explores the parameter space searching for the minimum of the equation. In this implementation, the
 310 function to be minimized should represent the difference between the measure and theoretical corre-
 311 lation functions. However, a simple χ^2 test is inappropriate for fitting correlation functions, as the ratio
 312 two Poisson distributions does not result in a Poisson distribution. Instead, a log-likelihood fit function
 313 of the following form is used [?]:




Fig. 10: Average separation (cm) correlation functions of Ξ Daughter and K^\pm . In the subfigure titles, the particles in “()” represent the mothers, ex. top left is p from Λ from Ξ^- with K^+ .

$$\chi_{PML}^2 = -2 \left[A \ln \left(\frac{C(A+B)}{A(C+1)} \right) + B \ln \left(\frac{A+B}{B(C+1)} \right) \right] \quad (8)$$

where A is the experimental signal distribution (numerator), B is the experimental background distribution (denominator), and C is the theoretical fit correlation function.

The LednickyFitter uses Equations 5 – 7 to build the theoretical fit, and Equation 8 as the statistic quantifying the quality of the fit. The parameters to be varied by MINUIT are: λ , R , f_0 ($\mathbb{R}f_0$ and $\mathbb{I}f_0$ separately), d_0 , and normalization N . The fitter currently includes methods to correct for momentum resolution and a non-flat background. These corrections are applied to the fit function, the data is never touched. The




Fig. 11: ΛK_S^0 (left) and $\bar{\Lambda} K_S^0$ (right) correlation functions for 0-10% (top), 10-30% (middle), and 30-50% (bottom) centralities. The lines represent the statistical errors, while the boxes represent the systematic errors.

fitter is able to share parameters between different analyses and fit all simultaneously.

In a typical fit, a given pair is fit with its conjugate (ex. ΛK^+ with $\bar{\Lambda} K^-$) across all centralities (0-10%, 10-30%, 30-50%), for a total of 6 simultaneous analyses. Each analysis has a unique λ and normalization parameter. The radii are shared between analyses of like centrality, as these should have similar source sizes. The scattering parameters ($\mathbb{R}f_0, \mathbb{I}f_0, d_0$) are shared amongst all.

Figures 19, 21, and 23 (Section 7) show experimental data with fits for all studied centralities for ΛK_S^0 with $\bar{\Lambda} K_S^0$, ΛK^+ with $\bar{\Lambda} K^-$, and ΛK^- with $\bar{\Lambda} K^+$, respectively. In the figures, the black solid line represents the “raw” fit, i.e. not corrected for momentum resolution effects nor non-flat background. The green line shows the fit to the non-flat background. The purple points show the fit after momentum resolution and non-flat background corrections have been applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties.

5.2 Model: $\Xi^{ch}K^{ch}$

When fitting the $\Xi^-(\bar{\Xi}^+)K^\pm$ results, it is necessary to include both strong and Coulomb effects. In this case, Equation 5 is no longer valid, and, in fact, there is no analytical form with which to fit. Therefore, we must begin with the wave function describing the pair interaction, and simulate many particle pairs to obtain a theoretical fit correlation function. The code developed to achieve this functionality is called “CoulombFitter”. Currently, in order to generate the statistics needed for a stable fit, we find that $\sim 10^4$ simulated pairs per 10 MeV bin are necessary. Unfortunately, the nature of this process means that the “CoulombFitter” takes much longer to run than the “LednickyFitter” of Section 5.1.

The two-particle correlation function may be written as:




Fig. 12: ΛK^+ (left) and $\bar{\Lambda} K^-$ (right) correlation functions for 0-10% (top), 10-30% (middle), and 30-50% (bottom) centralities. The lines represent the statistical errors, while the boxes represent the systematic errors.

$$C(\mathbf{k}^*) = \sum_S \rho_S \int S(\mathbf{r}^*) |\Psi_{\mathbf{k}^*}^S(\mathbf{r}^*)|^2 d^3 \mathbf{r}^* \quad (9)$$

where ρ_S is the normalized emission probability of particles in a state with spin S , $S(\mathbf{r}^*)$ is the pair emission source distribution (assumed to be Gaussian), and $\Psi_{\mathbf{k}^*}^S(\mathbf{r}^*)$ is the two-particle wave-function including both strong and Coulomb interactions [?]:

$$\Psi_{\mathbf{k}^*}(\mathbf{r}^*) = e^{i\delta_c} \sqrt{A_c(\eta)} [e^{i\mathbf{k}^* \cdot \mathbf{r}^*} F(-i\eta, 1, i\xi) + f_c(k^*) \frac{\tilde{G}(\rho, \eta)}{r^*}] \quad (10)$$

where $\rho = k^* r^*$, $\eta = (k^* a_c)^{-1}$, $\xi = \mathbf{k}^* \cdot \mathbf{r}^* + k^* r^* \equiv \rho(1 + \cos \theta^*)$, and $a_c = (\mu z_1 z_2 e^2)^{-1}$ is the two-particle Bohr radius (including the sign of the interaction). δ_c is the Coulomb s-wave phase shift, $A_c(\eta)$ is the Coulomb penetration factor, $\tilde{G} = \sqrt{A_c}(G_0 + iF_0)$ is a combination of the regular (F_0) and singular (G_0) s-wave Coulomb functions. $f_c(k^*)$ is the s-wave scattering amplitude:

$$f_c(k^*) = [\frac{1}{f_0} + \frac{1}{2} d_0 k^{*2} - \frac{2}{a_c} h(\eta) - ik^* A_c(\eta)]^{-1} \quad (11)$$

where, the “h-function”, $h(\eta)$, is expressed through the digamma function, $\psi(z) = \Gamma'(z)/\Gamma(z)$ as:

$$h(\eta) = 0.5[\psi(i\eta) + \psi(-i\eta) - \ln(\eta^2)] \quad (12)$$




Fig. 13: ΛK^- (left) and $\bar{\Lambda} K^+$ (right) correlation functions for 0-10% (top), 10-30% (middle), and 30-50% (bottom) centralities. The lines represent the statistical errors, while the boxes represent the systematic errors. The peak at $k^* \approx 0.2$ GeV/c is due to the Ω^- resonance.

348 Unfortunately, with this analysis, we are not sensitive to, and therefore not able to distinguish between,
 349 the iso-spin singlet and triplet states. We proceed with our analysis, but the results must be interpreted
 350 as iso-spin averaged scattering parameters.

351 As stated before, to generate a fit correlation function, we must simulate a large number of pairs, calculate
 352 the wave-function, and average Ψ^2 over all pairs in a given k^* bin. Essentially, we calculate Equation 9
 353 by hand:

$$\begin{aligned}
 C(\mathbf{k}^*) &= \sum_S \rho_S \int S(\mathbf{r}^*) |\Psi_{\mathbf{k}^*}^S(\mathbf{r}^*)|^2 d^3 \mathbf{r}^* \\
 &\longrightarrow C(|\mathbf{k}^*|) \equiv C(k^*) = \sum_S \rho_S \langle |\Psi^S(\mathbf{k}_i^*, \mathbf{r}_i^*)|^2 \rangle_i \\
 &\longrightarrow C(k^*) = \lambda \sum_S \rho_S \langle |\Psi^S(\mathbf{k}_i^*, \mathbf{r}_i^*)|^2 \rangle_i + (1 - \lambda)
 \end{aligned} \tag{13}$$

354 where $\langle \rangle_i$ represents an average over all pairs in a given k^* bin.

355 In summary, for a given k^* bin, we must draw $N_{\text{pairs}} \sim 10^4$ pairs, and for each pair:

- 356 1. Draw a random \mathbf{r}^* vector according to our Gaussian source distribution $S(\mathbf{r}^*)$
- 357 2. Draw a random \mathbf{k}^* vector satisfying the $|\mathbf{k}^*|$ restriction of the bin
 - 358 – We draw from real k^* vectors obtained from the data




Fig. 14: Correlation Functions: ΛK^+ vs ΛK^- ($\bar{\Lambda} K^+$ vs $\bar{\Lambda} K^-$) for 0-10% centrality. The peak in ΛK^- ($\bar{\Lambda} K^+$) at $k^* \approx 0.2 \text{ GeV}/c$ is due to the Ω^- resonance. The lines represent the statistical errors. (NOTE: This figure is slightly dated, and a new one will be generated which includes both statistical and systematic uncertainties)

359 – However, we find that drawing from a distribution flat in k^* gives similar results

360 3. Construct the wave-function Ψ

361 After all pairs for a given k^* bin are simulated and wave-functions obtained, the results are averaged to
362 give the fit result.

363 Construction of the wave-functions, Equation 10, involves a number of complex functions not included
364 in standard C++ or ROOT libraries (namely, $h(\eta)$, $\tilde{G}(\rho, \eta)$), and $F(-i\eta, 1, i\xi)$. These functions were
365 even difficult to find and implement from elsewhere. Our solution was to embed a Mathematica kernel
366 into our C++ code to evaluate these functions. However, having Mathematica work on-the-fly with the
367 fitter was far too time consuming (fitter would have taken day, maybe weeks to finish). Our solution
368 was to use Mathematica to create matrices representing these functions for different parameter values.
369 During fitting, these matrices were then interpolated and the results used to build the wave-functions.
370 This method decreased the running time dramatically, and we are not able to generate results in under \sim
371 1 hour. This process will be explained in more detail in future versions of the note.

372 **5.3 Momentum Resolution Corrections**

373 Finite track momentum resolution causes the reconstructed momentum of a particle to smear around the
374 true value. This, of course, also holds true for V0 particles. The effect is propagated up to the pairs
375 of interest, which causes the reconstructed relative momentum (k_{Rec}^*) to differ from the true momentum
376 (k_{True}^*). Smearing of the momentum typically will result in a suppression of the signal.

377 The effect of finite momentum resolution can be investigated using the MC data, for which both the true

378 and reconstructed momenta are available. Figure 15 shows sample k_{True}^* vs. k_{Rec}^* plots for $\Lambda(\bar{\Lambda})K^\pm$ 0-
 379 10% analyses; Figure 15a was generated using same-event pairs, while Figure 15b was generated using
 380 mixed-event pairs (with $N_{mix} = 5$).

381 If there are no contaminations in our particle collection, the plots in Figure 15 should be smeared around
 382 $k_{True}^* = k_{Rec}^*$; this is mostly true in our analyses. However, there are some interesting features of our results
 383 which demonstrate a small (notice the log-scale on the z-axis) contamination in our particle collection.
 384 The structure around $k_{Rec}^* = k_{True}^* - 0.15$ is mainly caused by K_S^0 contamination in our $\Lambda(\bar{\Lambda})$ sample. The
 385 remaining structure not distributed about $k_{Rec}^* = k_{True}^*$ is due to π and e contamination in our K^\pm sample.
 386 These contaminations are more visible in Figure 16, which show k_{Rec}^* vs. k_{True}^* plots (for a small sample
 387 of the ΛK^+ 0-10% central analysis), for which the MC truth (i.e. true, known identity of the particle)
 388 was used to eliminate misidentified particles in the $K^+(a)$ and $\Lambda(b)$ collections. (NOTE: This is an old
 389 figure and is for a small sample of the data. A new version will be generated shortly. It, nonetheless,
 390 demonstrates the point well).

391 Information gained from looking at k_{Rec}^* vs k_{True}^* can be used to apply corrections to account for the
 392 effects of finite momentum resolution on the correlation functions. A typical method involves using the
 393 MC HIJING data to build two correlation functions, $C_{Rec}(k^*)$ and $C_{True}(k^*)$, using the generator-level
 394 momentum (k_{True}^*) and the measured detector-level momentum (k_{Rec}^*). The data is then corrected by
 395 multiplying by the ratio, C_{True}/C_{Rec} , before fitting. This essentially unsmears the data, which that can
 396 be compared directly to theoretical predictions and fits. Although this is conceptually simple, there are
 397 a couple of big disadvantages to this method. First, HIJING does not incorporate final-state interactions,
 398 so weights must be used when building same-event (numerator) distributions. These weights account for
 399 the interactions, and, in the absence of Coulomb interactions, can be calculated using Eq. 5. Of course,
 400 these weights are valid only for a particular set of fit parameters. Therefore, in the fitting process, during
 401 which the fitter explores a large parameter set, the corrections will not remain valid. As such, applying
 402 the momentum resolution correction and fitting becomes a long and drawn out iterative process. An initial
 403 parameter set is obtained (through fitting without momentum resolution corrections, theoretical models,
 404 or a good guess), then the MC data is run over to obtain the correction factor, the data is fit using the
 405 correction factor, a refined parameter set is extracted, the MC data is run over again to obtain the new
 406 correction factor, etc. This process continues until the parameter set stabilizes. The second issue concerns
 407 statistics. With the MC data available on the grid, we were not able to generate the statistics necessary
 408 to use the raw C_{True}/C_{Rec} ratio. The ratio was not stable, and when applied to the data, obscured the
 409 signal. Attempting to fit the ratio to use to generate the corrections also proved problematic. However, as
 410 HIJING does not include final-state interactions, the same-event and mixed-event pairs are very similar
 411 (with the exception of things like energy and momentum conservation, etc). Therefore, one may build
 412 the numerator distribution using mixed-event pairs. This corresponds, more or less, to simply running a
 413 the weight generator through the detector framework.

414 A second approach is to use information gained from plots like those in Figure 15, which can be considered
 415 response matrices. The response matrix describes quantitatively how each k_{Rec}^* bin receives contributions
 416 from multiple k_{True}^* bins, and can be used to account for the effects of finite momentum resolution.
 417 With this approach, the resolution correction is applied on-the-fly during the fitting process by propagating
 418 the theoretical (fit) correlation function through the response matrix, according to:

$$C_{fit}(k_{Rec}^*) = \frac{\sum_{k_{True}^*} M_{k_{Rec}^*, k_{True}^*} C_{fit}(k_{True}^*)}{\sum_{k_{True}^*} M_{k_{Rec}^*, k_{True}^*}} \quad (14)$$

419 where $M_{k_{Rec}^*, k_{True}^*}$ is the response matrix (Figure 15), $C_{fit}(k_{True}^*)$ is the fit binned in k_{True}^* , and the denominator normalizes the result.

421 Equation 14 describes that, for a given k_{Rec}^* bin, the observed value of $C(k_{Rec}^*)$ is a weighted average of
 422 all $C(k_{True}^*)$ values, where the weights are the normalized number of counts in the $[k_{Rec}^*, k_{True}^*]$ bin. As
 423 seen in Figure 15, overwhelmingly the main contributions comes from the $k_{Rec}^* = k_{True}^*$ bins. Although
 424 the correction is small, it is non-negligible for the low- k^* region of the correlation function.

425 Here, the momentum resolution correction is applied to the fit, not the data. In other words, during
 426 fitting, the theoretical correlation function is smeared just as real data would be, instead of unsmearing
 427 the data. This may not be ideal for the theorist attempting to compare a model to experimental data, but
 428 it leaves the experimental data unadulterated. The current analyses use this second approach to applying
 429 momentum resolution corrections because of two major advantages. First, the MC data must be analyzed
 430 only once, and no assumptions about the fit are needed. Secondly, the momentum resolution correction
 431 is applied on-the-fly by the fitter, delegating the iterative process to a computer instead of the user.

432 5.4 Residual Correlations

433 The purpose of this analysis is study the interaction and scale of the emitting source of the pairs. In order
 434 to obtain correct results, it is important for our particle collections to consist of primary particles. In
 435 practice, this is difficult to achieve for our Λ and $\bar{\Lambda}$ collections. Many of our Λ particles are not primary,
 436 but originate as decay products from other hyperons, including Σ^0 , Ξ^- , Ξ^0 and Ω^- . In these decays,
 437 the Λ carries away a momentum very similar to that of its parent. As a result, the correlation function
 438 between a secondary Λ and, for instance, a K^+ will be sensitive to, and dependent upon, the interaction
 439 between the parent of the Λ and the K^+ . In effect, the correlation between the parent of the Λ and the K^+
 440 (ex. $\Sigma^0 K^+$) will be visible, although smeared out, in the ΛK^+ data. We call this a residual correlation
 441 resulting from feed-down.

442 As it is difficult for us to eliminate these residual correlations in our analyses, we must attempt to account
 443 for them in our fitter. To achieve this, we will simultaneously fit the data for both the primary correlation
 444 function and the residual correlations. For example, in the simple case of a ΛK^+ analysis with residuals
 445 arising soley from $\Sigma^0 K^+$ feed-down:

$$C_{measured}(k_{\Lambda K^+}^*) = 1 + \lambda_{\Lambda K^+}[C_{\Lambda K^+}(k_{\Lambda K^+}^*) - 1] + \lambda_{\Sigma^0 K^+}[C_{\Sigma^0 K^+}(k_{\Lambda K^+}^*) - 1]$$

$$C_{\Sigma^0 K^+}(k_{\Lambda K^+}^*) \equiv \frac{\sum_{k_{\Sigma^0 K^+}^*} C_{\Sigma^0 K^+}(k_{\Sigma^0 K^+}^*) T(k_{\Sigma^0 K^+}^*, k_{\Lambda K^+}^*)}{\sum_{k_{\Sigma^0 K^+}^*} T(k_{\Sigma^0 K^+}^*, k_{\Lambda K^+}^*)} \quad (15)$$

446 $C_{\Sigma^0 K^+}(k_{\Sigma^0 K^+}^*)$ is the $\Sigma^0 K^+$ correlation function from, for instance, Equation 5, and T is the transform
 447 matrix generated with THERMINATOR.

448 This equation can be easily extended to include feed-down from more sources:

$$C_{measured}(k_{\Lambda K}^*) = 1 + \lambda_{\Lambda K}[C_{\Lambda K}(k_{\Lambda K}^*) - 1] + \lambda_{\Sigma^0 K}[C_{\Sigma^0 K}(k_{\Lambda K}^*) - 1] + \dots$$

$$+ \lambda_{P_1 P_2}[C_{P_1 P_2}(k_{\Lambda K}^*) - 1] + \lambda_{other}[C_{other}(k_{\Lambda K}^*) - 1]$$

$$C_{P_1 P_2}(k_{\Lambda K}^*) \equiv \frac{\sum_{k_{P_1 P_2}^*} C_{P_1 P_2}(k_{P_1 P_2}^*) T(k_{P_1 P_2}^*, k_{\Lambda K}^*)}{\sum_{k_{P_1 P_2}^*} T(k_{P_1 P_2}^*, k_{\Lambda K}^*)} \quad (16)$$

449 Or, more compactly:





Fig. 15: Sample k_{True}^* vs. k_{Rec}^* plot for $\Lambda(\bar{\Lambda})K^\pm$ 0-10% analyses. The structure which appears around $k_{Rec}^* = k_{True}^* - 0.15$ is mainly caused by K_S^0 contamination in our $\Lambda(\bar{\Lambda})$ sample. The remaining structure not distributed about $k_{Rec}^* = k_{True}^*$ is due to π and e contamination in our K^\pm sample. These contaminations are more clearly visible in Figure 16


$$C_{measured}(k_{\Lambda K}^*) = 1 + \sum_i \lambda_i [C_i(k_{\Lambda K}^*) - 1] \quad (17)$$

The framework for extracting the necessary transform matrices from the THERMINATOR data is already in place, and has been used to generate the examples from ΛK^+ and $\bar{\Lambda} K^+$ analyses shown in Figures 17 and 18. However, these residual correlations have not yet been implemented in the fitter.

There is an obvious added complication here, as, for instance, the $\Xi^- K^\pm$ residuals necessitate the inclusion of the CoulombFitter into the process. The complication of combining the two fitters is not as troubling as the increase in fitting time that this is sure to bring. We have two solutions to bypass such a



(a) (Top Left) All misidentified K^+ excluded. (Bottom Left) All misidentified Λ and K^+ excluded. (Right) The difference of (Top Left) - (Bottom Left), which reveals the contamination in our Λ collection. The structure which appears around $k_{Rec}^* = k_{True}^* - 0.15$ is mainly caused by K_S^0 contamination in our $\Lambda(\bar{\Lambda})$ sample.



(b) (Top Left) All misidentified Λ excluded. (Bottom Left) All misidentified Λ and K^+ excluded. (Right) The difference of (Top Left) - (Bottom Left), which reveals the contamination in our K^+ collection. The structure not distributed about $k_{Rec}^* = k_{True}^*$ is due to π and e^- contamination in our K^+ sample.

Fig. 16: Note: This is an old figure and is for a small sample of the data. A new version will be generated shortly.
y-axis = k_{Rec}^* , x-axis = k_{True}^* .

(Left) k_{Rec}^* vs. k_{True}^* plots for a small sample of the ΛK^+ 0-10% central analysis, MC truth was used to eliminate misidentified particles in the K^+ (a) and Λ (b) collections. (Right) The difference of the top left and bottom left plots. Contaminations in our particle collections are clearly visible. Figure (a) demonstrates a K_S^0 contamination in our Λ collection; Figure (b) demonstrates a π and e^- contamination in our K^\pm collection.

456 large increase in run time. First, we can use our experimental $E^{ch}K^{ch}$ data to represent all charged parent
457 pair system. Alternatively, we can assume the strong interaction is negligible in the charged residual,
458 and generate the parent correlation function given radius and λ parameters. We find in our $E^{ch}K^{ch}$ that a

459 Coulomb-only description of the system describes, reasonable well, the broad features of the correlation.
 460 The strong interaction is necessary for the fine details. However, as these correlations are run through
 461 a transform matrix, which largely flattens out and fine details, a Coulomb-only description should be
 462 sufficient.




Fig. 17: Transform Matrices generated with THERMINATOR for ΛK^+ Analysis

463 6 Systematic Errors

464 In order to understand my systematic uncertainties, the analysis code was run many times using slightly
 465 different values for a number of important cuts, and the results were compared.
 466 In order to quantify the systematic errors on the data, all correlation functions built using all varied cut
 467 values were bin-by-bin averaged, and the resulting variance of each bin was taken as the systematic error.
 468 The cuts which were utilized in this study are presented in Sections 6.1.1 (ΛK_S^0) and 6.2.1 (ΛK^\pm).
 469 Similarly, the fit parameters extracted from all of these correlation functions were averaged, and the
 470 resulting variances were taken as the systematic errors for the fit parameters. As with the systematic
 471 errors on the data, this was performed for all varied cut values. Additionally, a systematic analysis was
 472 done on our fit method (which, for now, just includes our choice of fit range). These two sources of
 473 uncertainty were combined in quadrature to obtain the final systematic uncertainties on the extracted fit
 474 parameters.




Fig. 18: Transform Matrices generated with THERMINATOR for $\bar{\Lambda}K^+$ Analysis

475 6.1 Systematic Errors: ΛK_S^0

476 6.1.1 Particle and Pair Cuts

477 The cuts included in the systematic study, as well as the values used in the variations, are listed below.
 478 Note, the central value corresponds to that used in the analysis.

- 479 1. DCA $\Lambda(\bar{\Lambda})$: {4, 5, 6 mm}
- 480 2. DCA K_S^0 : {2, 3, 4 mm}
- 481 3. DCA $\Lambda(\bar{\Lambda})$ Daughters: {3, 4, 5 mm}
- 482 4. DCA K_S^0 Daughters: {2, 3, 4 mm}
- 483 5. $\Lambda(\bar{\Lambda})$ Cosine of Pointing Angle: {0.9992, 0.9993, 0.9994}
- 484 6. K_S^0 Cosine of Pointing Angle: {0.9992, 0.9993, 0.9994}
- 485 7. DCA to Primary Vertex of $p(\bar{p})$ Daughter of $\Lambda(\bar{\Lambda})$: {0.5, 1, 2 mm}
- 486 8. DCA to Primary Vertex of $\pi^-(\pi^+)$ Daughter of $\Lambda(\bar{\Lambda})$: {2, 3, 4 mm}
- 487 9. DCA to Primary Vertex of π^+ Daughter of K_S^0 : {2, 3, 4 mm}
- 488 10. DCA to Primary Vertex of π^- Daughter of K_S^0 : {2, 3, 4 mm}
- 489 11. Average Separation of Like-Charge Daughters: {5, 6, 7 cm}

490 **6.1.2 Non-Flat Background**

491 We fit our non-flat background with a linear function. To study the contribution of this choice to our sys-
 492 tematic errors, we also fit with a quadratic and Gaussian form. The resulting uncertainties are combined
 493 with the uncertainties arising from our particle cuts.

494 **6.1.3 Fit Range**

495 Our choice of k^* fit range was varied by $\pm 25\%$. The resulting uncertainties in the extracted parameter
 496 sets were combined with our uncertainties arising from our particle and pair cuts.

497 **6.2 Systematic Errors: ΛK^\pm**

498 **6.2.1 Particle and Pair Cuts**

499 The cuts included in the systematic study, as well as the values used in the variations, are listed below.
 500 Note, the central value corresponds to that used in the analysis.

- 501 1. DCA $\Lambda(\bar{\Lambda})$: {4, 5, 6 mm}
- 502 2. DCA $\Lambda(\bar{\Lambda})$ Daughters: {3, 4, 5 mm}
- 503 3. $\Lambda(\bar{\Lambda})$ Cosine of Pointing Angle: {0.9992, 0.9993, 0.9994}
- 504 4. DCA to Primary Vertex of $p(\bar{p})$ Daughter of $\Lambda(\bar{\Lambda})$: {0.5, 1, 2 mm}
- 505 5. DCA to Primary Vertex of $\pi^-(\pi^+)$ Daughter of $\Lambda(\bar{\Lambda})$: {2, 3, 4 mm}
- 506 6. Average Separation of $\Lambda(\bar{\Lambda})$ Daughter with Same Charge as K^\pm : {7, 8, 9 cm}
- 507 7. Max. DCA to Primary Vertex in Transverse Plane of K^\pm : {1.92, 2.4, 2.88}
- 508 8. Max. DCA to Primary Vertex in Longitudinal Direction of K^\pm : {2.4, 3.0, 3.6}

509 **6.2.2 Non-Flat Background**

510 We fit our non-flat background with a linear function. To study the contribution of this choice to our sys-
 511 tematic errors, we also fit with a quadratic and Gaussian form. The resulting uncertainties are combined
 512 with the uncertainties arising from our particle cuts.

513 **6.2.3 Fit Range**

514 Our choice of k^* fit range was varied by $\pm 25\%$. The resulting uncertainties in the extracted parameter
 515 sets were combined with our uncertainties arising from our particle and pair cuts.

516 **6.3 Systematic Errors: ΞK^\pm**

517 **6.3.1 Particle and Pair Cuts**

518 The cuts included in the systematic study, as well as the values used in the variations, are listed below.
 519 Note, the central value corresponds to that used in the analysis.

- 520 1. Max. DCA $\Xi(\bar{\Xi})$: {x, y, z mm}
- 521 2. Max. DCA $\Xi(\bar{\Xi})$ Daughters: {x, y, z mm}
- 522 3. Min. $\Xi(\bar{\Xi})$ Cosine of Pointing Angle to Primary Vertex: {0.9991, 0.9992, 0.9993}

- 523 4. Min. $\Lambda(\bar{\Lambda})$ Cosine of Pointing Angle to $\Xi(\bar{\Xi})$ Decay Vertex: {0.9992, 0.9993, 0.9994}
- 524 5. Min. DCA Bachelor π : {0.2, 0.3, 0.4 mm}
- 525 6. Min. DCA $\Lambda(\bar{\Lambda})$: {0.5, 1, 2 mm}
- 526 7. Max. DCA $\Lambda(\bar{\Lambda})$ Daughters: {3, 4, 5 mm}
- 527 8. Min. $\Lambda(\bar{\Lambda})$ Cosine of Pointing Angle To Primary Vertex: {0.9992, 0.9993, 0.9994}
- 528 9. Min. DCA to Primary Vertex of $p(\bar{p})$ Daughter of $\Lambda(\bar{\Lambda})$: {0.5, 1, 2 mm}
- 529 10. Min. DCA to Primary Vertex of $\pi^-(\pi^+)$ Daughter of $\Lambda(\bar{\Lambda})$: {2, 3, 4 mm}
- 530 11. Min. Average Separation of $\Lambda(\bar{\Lambda})$ Daughter and K^\pm with like charge: {7, 8, 9 cm}
- 531 12. Min. Average Separation of Bachelor π and K^\pm with like charge: {x, y, z cm}
- 532 13. Max. DCA to Primary Vertex in Transverse Plane of K^\pm : {1.92, 2.4, 2.88}
- 533 14. Max. DCA to Primary Vertex in Longitudinal Direction of K^\pm : {2.4, 3.0, 3.6}


534 7 Results and Discussion

535 7.1 Results: ΛK_S^0 and ΛK^\pm

536 Figures 19, 21, and 23 (Section 7) show experimental data with fits for all studied centralities for ΛK_S^0
 537 with $\bar{\Lambda} K_S^0$, ΛK^+ with $\bar{\Lambda} K^-$, and ΛK^- with $\bar{\Lambda} K^+$, respectively. The parameter sets extracted from the fits
 538 can be found in Tables 1 and 2. All correlation functions were normalized in the range $0.32 < k^* < 0.40$
 539 GeV/c, and fit in the range $0.0 < k^* < 0.30$ GeV/c. For the ΛK^- and $\bar{\Lambda} K^+$ analyses, the region 0.19
 540 $< k^* < 0.23$ GeV/c was excluded from the fit to exclude the bump caused by the Ω^- resonance. The
 541 non-flat background was fit with a linear form from $0.6 < k^* < 0.9$ GeV/c. The theoretical fit function
 542 was then multiplied by this background during the fitting process.

543 In the figures (19, 21, and 23), the black solid line represents the “raw” fit, i.e. not corrected for momen-
 544 tum resolution effects nor non-flat background. The green line shows the fit to the non-flat background.
 545 The purple points show the fit after momentum resolution and non-flat background corrections have been
 546 applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties.

547 For the ΛK_S^0 fits, R was restricted to [2.0, 10.0 fm] and Λ was restricted to [0.1, 0.8]. This gave the lowest
 548 χ^2 value, but loosening this restriction changes the fit parameters slightly. Notice, the 10-30% radius is
 549 at its limit, as is λ from the 30-50% ΛK_S^0 analysis. This accounts for the 0.000 systematic uncertainty of
 550 the 10-30% R value currently quoted in Table 1. An estimate for this uncertainty should be included in
 551 the next version of this note. In the future, we may need to throw out the 30-50% data from the fit, but
 552 this is not ideal.



ALI-PREL-125950

Fig. 19: Fits to the ΛK_s^0 (left) and $\bar{\Lambda} K_s^0$ (right) data for the centralities 0-10% (top), 10-30% (middle), and 30-50% (bottom). The lines represent the statistical errors, while the boxes represent the systematic errors. Each has unique λ and normalization parameters. The radii are shared amongst like centralities; the scattering parameters ($\mathbb{R}f_0$, $\mathbb{I}f_0$, d_0) are shared amongst all. The black solid line represents the “raw” fit, i.e. not corrected for momentum resolution effects nor non-flat background. The green line shows the fit to the non-flat background. The purple points show the fit after momentum resolution and non-flat background corrections have been applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties. Here, R was restricted to [2.,10.] and Λ was restricted to [0.1,0.8].




Fig. 20: Same as Fig. 19, but with a wider range of view. Fits to the ΛK_S^0 (left) and $\bar{\Lambda} K_S^0$ (right) data for the centralities 0-10% (top), 10-30% (middle), and 30-50% (bottom). The lines represent the statistical errors, while the boxes represent the systematic errors. Each has unique λ and normalization parameters. The radii are shared amongst like centralities; the scattering parameters ($\mathbb{R}f_0$, $\mathbb{I}f_0$, d_0) are shared amongst all. The black solid line represents the “raw” fit, i.e. not corrected for momentum resolution effects nor non-flat background. The green line shows the fit to the non-flat background. The purple points show the fit after momentum resolution and non-flat background corrections have been applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties. Here, R was restricted to [2.,10.] and Λ was restricted to [0.1,0.8].




Fig. 21: Fits to the ΛK^+ (left) and $\bar{\Lambda} K^-$ (right) data for the centralities 0-10% (top), 10-30% (middle), and 30-50% (bottom). The lines represent the statistical errors, while the boxes represent the systematic errors. Each has unique λ and normalization parameters. The radii are shared amongst like centralities; the scattering parameters ($\text{Re}[f_0]$, $\text{Im}[f_0]$, d_0) are shared amongst all. The black solid line represents the “raw” fit, i.e. not corrected for momentum resolution effects nor non-flat background. The green line shows the fit to the non-flat background. The purple points show the fit after momentum resolution and non-flat background corrections have been applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties.





Fig. 22: Same as Fig. 21, but with a wider range of view. Fits to the ΛK^+ (left) and $\bar{\Lambda} K^-$ (right) data for the centralities 0-10% (top), 10-30% (middle), and 30-50% (bottom). The lines represent the statistical errors, while the boxes represent the systematic errors. Each has unique λ and normalization parameters. The radii are shared amongst like centralities; the scattering parameters ($\mathbb{R}f_0$, $\mathbb{I}f_0$, d_0) are shared amongst all. The black solid line represents the “raw” fit, i.e. not corrected for momentum resolution effects nor non-flat background. The green line shows the fit to the non-flat background. The purple points show the fit after momentum resolution and non-flat background corrections have been applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties.



ALI-PREL-125931

Fig. 23: Fits to the ΛK^- (left) with $\bar{\Lambda} K^+$ (right) data for the centralities 0-10% (top), 10-30% (middle), and 30-50% (bottom). The lines represent the statistical errors, while the boxes represent the systematic errors. Each has unique λ and normalization parameters. The radii are shared amongst like centralities; the scattering parameters ($\text{Re}[f_0]$, $\text{Im}[f_0]$, d_0) are shared amongst all. The black solid line represents the “raw” fit, i.e. not corrected for momentum resolution effects nor non-flat background. The green line shows the fit to the non-flat background. The purple points show the fit after momentum resolution and non-flat background corrections have been applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties.




Fig. 24: Same as Fig. 23, but with a wider range of view. Fits to the ΛK^- (left) with $\bar{\Lambda} K^+$ (right) data for the centralities 0-10% (top), 10-30% (middle), and 30-50% (bottom). The lines represent the statistical errors, while the boxes represent the systematic errors. Each has unique λ and normalization parameters. The radii are shared amongst like centralities; the scattering parameters ($\mathbb{R} f_0$, $\mathbb{I} f_0$, d_0) are shared amongst all. The black solid line represents the “raw” fit, i.e. not corrected for momentum resolution effects nor non-flat background. The green line shows the fit to the non-flat background. The purple points show the fit after momentum resolution and non-flat background corrections have been applied. The initial values of the parameters is listed, as well as the final fit values with uncertainties.

Fit Results $\Lambda(\bar{\Lambda})K_S^0$						
Pair Type	Centrality	Fit Parameters				
		λ	R	$\mathbb{R}f_0$	$\mathbb{I}f_0$	d_0
ΛK_S^0	0-10%	0.400 \pm 0.187 (stat.) \pm 0.116 (sys.)	3.024 \pm 0.541 (stat.) \pm 0.329 (sys.)	-0.157 \pm 0.031 (stat.) \pm 0.043 (sys.)	0.176 \pm 0.077 (stat.) \pm 0.059 (sys.)	3.566 \pm 0.947 (stat.) \pm 2.836 (sys.)
	10-30%		2.270 \pm 0.413 (stat.) \pm 0.324 (sys.)			
	30-50%		1.669 \pm 0.307 (stat.) \pm 0.280 (sys.)			
	0-10%	0.400 \pm 0.187 (stat.) \pm 0.116 (sys.)	3.024 \pm 0.541 (stat.) \pm 0.329 (sys.)	-0.157 \pm 0.031 (stat.) \pm 0.043 (sys.)	0.176 \pm 0.077 (stat.) \pm 0.059 (sys.)	3.566 \pm 0.947 (stat.) \pm 2.836 (sys.)
	10-30%		2.270 \pm 0.413 (stat.) \pm 0.324 (sys.)			
	30-50%		1.669 \pm 0.307 (stat.) \pm 0.280 (sys.)			

Table 1: Fit Results $\Lambda(\bar{\Lambda})K_S^0$. Each pair is fit simultaneously with its conjugate (ie. ΛK_S^0 with $\bar{\Lambda} K_S^0$) across all centralities (0-10%, 10-30%, 30-50%), for a total of 6 simultaneous analyses in the fit. Each analysis has a unique λ and normalization parameter. The radii are shared between analyses of like centrality, as these should have similar source sizes. The scattering parameters ($\mathbb{R}f_0$, $\mathbb{I}f_0$, d_0) are shared amongst all. The fit is done on the data with only statistical error bars. The errors marked as “stat.” are those returned by MINUIT. The errors marked as “sys.” are those which result from my systematic analysis (as outlined in Section 6).

Fit Results $\Lambda(\bar{\Lambda})K^\pm$						
Pair Type	Centrality	Fit Parameters				
		λ	R	$\mathbb{R}f_0$	$\mathbb{I}f_0$	d_0
ΛK^+	0-10%	0.379 \pm 0.085 (stat.) \pm 0.220 (sys.)	4.045 \pm 0.381 (stat.) \pm 0.830 (sys.)	-0.687 \pm 0.160 (stat.) \pm 0.223 (sys.)	0.391 \pm 0.143 (stat.) \pm 0.111 (sys.)	0.639 \pm 0.534 (stat.) \pm 1.621 (sys.)
	10-30%	0.485 \pm 0.129 (stat.) \pm 0.241 (sys.)	3.923 \pm 0.454 (stat.) \pm 0.663 (sys.)			
	30-50%	0.639 \pm 0.195 (stat.) \pm 0.204 (sys.)	3.717 \pm 0.554 (stat.) \pm 0.420 (sys.)			
	0-10%	0.371 \pm 0.083 (stat.) \pm 0.217 (sys.)	4.045 \pm 0.381 (stat.) \pm 0.830 (sys.)	0.183 \pm 0.134 (stat.) \pm 0.095 (sys.)	0.453 \pm 0.181 (stat.) \pm 0.184 (sys.)	-5.292 \pm 2.895 (stat.) \pm 7.658 (sys.)
	10-30%	0.411 \pm 0.111 (stat.) \pm 0.201 (sys.)	3.923 \pm 0.454 (stat.) \pm 0.663 (sys.)			
	30-50%	0.616 \pm 0.192 (stat.) \pm 0.203 (sys.)	3.717 \pm 0.554 (stat.) \pm 0.420 (sys.)			
$\bar{\Lambda} K^-$	0-10%	0.453 \pm 0.162 (stat.) \pm 0.186 (sys.)	4.787 \pm 0.788 (stat.) \pm 1.375 (sys.)	0.183 \pm 0.134 (stat.) \pm 0.095 (sys.)	0.453 \pm 0.181 (stat.) \pm 0.184 (sys.)	-5.292 \pm 2.895 (stat.) \pm 7.658 (sys.)
	10-30%	0.395 \pm 0.149 (stat.) \pm 0.198 (sys.)	4.001 \pm 0.719 (stat.) \pm 0.978 (sys.)			
	30-50%	0.199 \pm 0.077 (stat.) \pm 0.132 (sys.)	2.112 \pm 0.517 (stat.) \pm 0.457 (sys.)			
	0-10%	0.479 \pm 0.170 (stat.) \pm 0.152 (sys.)	4.787 \pm 0.788 (stat.) \pm 1.375 (sys.)	-0.157 \pm 0.031 (stat.) \pm 0.043 (sys.)	0.176 \pm 0.077 (stat.) \pm 0.059 (sys.)	3.566 \pm 0.947 (stat.) \pm 2.836 (sys.)
	10-30%	0.491 \pm 0.179 (stat.) \pm 0.148 (sys.)	4.001 \pm 0.719 (stat.) \pm 0.978 (sys.)			
	30-50%	0.224 \pm 0.083 (stat.) \pm 0.106 (sys.)	2.112 \pm 0.517 (stat.) \pm 0.457 (sys.)			

Table 2: Fit Results $\Lambda(\bar{\Lambda})K^\pm$ Each pair is fit simultaneously with its conjugate (ie. ΛK^+ with $\bar{\Lambda} K^-$ and ΛK^- with $\bar{\Lambda} K^+$) across all centralities (0-10%, 10-30%, 30-50%), for a total of 6 simultaneous analyses in the fit. Each analysis has a unique λ and normalization parameter. The radii are shared between analyses of like centrality, as these should have similar source sizes. The scattering parameters ($\mathbb{R}f_0$, $\mathbb{I}f_0$, d_0) are shared amongst all. The fit is done on the data with only statistical error bars. The errors marked as “stat.” are those returned by MINUIT. The errors marked as “sys.” are those which result from my systematic analysis (as outlined in Section 6).

Fit Parameters (value \pm statistical error \pm systematic error)

Pair Type	Centrality	R		
$\Lambda K^+ \& \bar{\Lambda} K^-$	0-10%	$4.04 \pm 0.38 \pm 0.83$		
	10-30%	$3.92 \pm 0.45 \pm 0.66$		
	30-50%	$3.72 \pm 0.55 \pm 0.42$		
		$\Re f_0$	$\Im f_0$	d_0
		$-0.69 \pm 0.16 \pm 0.22$	$0.39 \pm 0.14 \pm 0.11$	$0.64 \pm 0.53 \pm 1.62$
$\Lambda K^- \& \bar{\Lambda} K^+$	0-10%	$4.79 \pm 0.79 \pm 1.38$		
	10-30%	$4.00 \pm 0.72 \pm 0.98$		
	30-50%	$2.11 \pm 0.52 \pm 0.46$		
		$\Re f_0$	$\Im f_0$	d_0
		$0.18 \pm 0.13 \pm 0.10$	$0.45 \pm 0.18 \pm 0.18$	$-5.29 \pm 2.94 \pm 7.66$
$\Lambda K_S^0 \& \bar{\Lambda} K_S^0$	0-10%	$3.02 \pm 0.54 \pm 0.33$		
	10-30%	$2.27 \pm 0.41 \pm 0.32$		
	30-50%	$1.67 \pm 0.30 \pm 0.28$		
		$\Re f_0$	$\Im f_0$	d_0
		$-0.16 \pm 0.03 \pm 0.04$	$0.18 \pm 0.08 \pm 0.06$	$3.57 \pm 0.95 \pm 2.84$

Figure 25 shows extracted R_{inv} parameters as a function of transverse mass (m_T) for various pair systems over several centralities. The published ALICE data [?] is shown with transparent, open symbols. The new ΛK results are shown with opaque, filled symbols. The radii show an increasing size with increasing centrality, as is expected from the simple geometric picture of the collisions. The radii decrease in size with increasing m_T , and we see an approximate scaling of the radii with transverse mass, as is expected in the presence of collective flow in the system.




Fig. 25: Extracted fit R_{inv} parameters as a function of pair transverse mass (m_T) for various pair systems over several centralities. The ALICE published data [?] is shown with transparent, open symbols. The new ΛK results are shown with opaque, filled symbols. In the left, the ΛK^+ (with its conjugate pair) results are shown separately from the ΛK^- (with its conjugate pair) results. In the right, all ΛK^\pm results are averaged.

559 7.2 Results: ΞK^\pm

Even without any fits to the data, the fact that the $\Xi^- K^+$ data dips below unity (Fig. 26) is exciting, as this cannot occur purely from a Coulomb interaction. We hope that this dip signifies that we are able to peer through the overwhelming contribution from the Coulomb interaction to see the effects arising from the strong interaction.




Fig. 26: ΞK^\pm Results for 0-10% Centrality. (Left) $\Xi^- K^+$ and $\Xi^- K^-$ (Right) $\Xi^+ K^+$ and $\Xi^+ K^-$

Figure 27 demonstrates graphically, that the $\Xi^- K^+$ results cannot be described by solely the Coulomb interaction. In this figure, we present the data along with a Coulomb-only band. The Coulomb-only band is spanned by two Coulomb-only curves, whose parameters are given in the figure. The Coulomb-only curves were generated using a technique identical to the generation of the fit function, described in Sec. 5.2, except, of course, with the nuclear scattering parameters all set to zero. The Coulomb-only




Fig. 27: ΞK^\pm data with Coulomb-only bands for the 0-10% centrality bin. The Coulomb-only bands span two sets of Coulomb-only curves: (1) $\lambda = 0.9$, $R = 1.0$ fm and (2) $\lambda = 0.1$, $R = 10.0$ fm. The Coulomb-only curves are simulated correlation functions for the respective pair system assuming only a Coulomb interaction, i.e. ignoring the strong interaction. The Coulomb-only curves change monotonically with varying λ and varying R , therefore, any intermediate parameter set will fall within this Coulomb-only band.

569 curves change monotonically with varying λ or varyin radius parametres, therefore, any curves built with
570 parameter sets intermediate to those use in the Coulomb-only band will be contained in the band.

571 Including the strong interaction into the simulation can dramatically change the resulting correlation
572 function, as shown in Figure 28. In the figure, the solid line represents a Coulomb-only curve, i.e. a
573 simulated correlation function with the strong interaction turned off. The dashed lines represent a full
574 simulation, including both the strong and Coulomb interactions. The two dashed lines differ only in the
575 real part of the assumed scattering length: positive in Set 1, and negative in Set 2. In the top figure,
576 for the $\Xi^- K^+$ simulation, we see that parameter set 2, with a negative real part of the scattering length,
577 causes the simulated curve to dip below unity, as is seen in the data. If there is a parallel to be drawn
578 between this analysis and the ΛK analysis, we expect to see similar effects in the ΛK^+ system and the
579 $\Xi^- K^+$ systems. In these systems, we could have an $s\bar{s}$ annihilation picture. Or, another possible way of
580 thinking about these systems is in terms of net strangeness. The ΛK^+ system has $S=0$, while the ΛK^-
581 has $S=-2$. The $\Xi^- K^+$ has $S=-1$, while the $\Xi^- K^-$ has $S=-3$.

582 The author was asked to perform a global Coulomb-only fit to the data, to ensure that the system truly
583 could not be described simply by the Coulomb interaction. In order words, in the fit, the strong force was
584 turned off, and the $\Xi^- K^+$, $\bar{\Xi}^+ K^-$, $\Xi^- K^-$, $\bar{\Xi}^+ K^+$ systems all share one singlge radius parameter, while the

pair and conjugate pair systems share a λ parameter. The results of this fit are shown in Figures 29 and 30. In Fig. 29, there was a lower limit of 0.1 fm placed on the radius parameter, and the radius parameter was initialized to 3 fm (as seems reasonable, when considering the transverse mass of the system and looking at Fig. 25). As is shown in the results, the radius parameter reached this unrealistic lower bound of 0.1 fm. In Fig. 30, the parameters were all unbounded, and the radius parameter was initialized to 10 fm. In this case, the radius parameters remains high, and ends at an unrealistic value of 10.84 fm. In both cases, the λ parameters are too low. From these figures, we conclude that a global Coulomb-only fit is not suitable for the data.

Although the global Coulomb-only fit failed, it is possible that a Coulomb-only fit performed on $\Xi^- K^+$ and $\bar{\Xi}^+ K^-$ separately from $\Xi^- K^-$ and $\bar{\Xi}^+ K^+$ could be suitable. The result of such fits are shown in Figures 31 and 32. Figure 31, shows that the fit is not able to describe the dip in the $\Xi^- K^+$ data below unity. Of course, this is obviously true for an attractive Coulomb-only fit. The radius parameter of 8.43 fm extracted from this fit is unrealistically large. In Figure 32 shows the Coulomb-only fit can described the $\Xi^- K^-$ data reasonable well; although the extracted radius of 3.73 fm is somewhat larger than expected.

8 To Do



(b) ΞK^- and $\bar{\Xi} K^+$ simulation

Fig. 28: Effect on the Coulomb-only curve of including the strong interaction for ΞK^\pm systems. The solid line represents a Coulomb-only curve, i.e. a simulated correlation function with the strong interaction turned off. The dashed lines represent a full simulation, including both the strong and Coulomb interactions. The two dashed lines differ only in the real part of the assumed scattering length: positive in Set 1, and negative in Set 2.






Fig. 29: ΞK^\pm Global Coulomb-only fit (Set 1) for 0-10% centrality. In this fit, there was a lower limit of 0.1 fm placed on the radius parameter, and the radius parameter was initialized to 3 fm (as seems reasonable, when considering the transverse mass of the system and looking at Fig. 25). As is shown in the results, the radius parameter reached this unrealistic lower bound of 0.1 fm. Also, the extracted λ parameters are too low.






Fig. 32: $\Xi^- \text{K}^-$ Coulomb-only fit for 0-10% centrality

601 **9 Additional Figures**

602 **9.1 Residuals**




Fig. 33: Residuals: $\Sigma^0 \text{K}^+$ to ΛK^+ (0-10% Centrality)




Fig. 34: Residuals: $\Xi^0 K^+$ to ΛK^+ (0-10% Centrality)




Fig. 35: Residuals: $\Xi^- K^+$ to ΛK^+ (0-10% Centrality)




Fig. 36: Residuals: Σ^*+K^+ to ΛK^+ (0-10% Centrality)




Fig. 37: Residuals: Σ^*-K^+ to ΛK^+ (0-10% Centrality)




Fig. 38: Residuals: $\Sigma^{*0} \text{K}^+$ to ΛK^+ (0-10% Centrality)




Fig. 39: Residuals: ΛK^{*0} to ΛK^+ (0-10% Centrality)




Fig. 40: Residuals: $\Sigma^0 K^{*0}$ to ΛK^+ (0-10% Centrality)




Fig. 41: Residuals: $\Xi^0 K^{*0}$ to ΛK^+ (0-10% Centrality)




Fig. 42: Residuals: $\Xi\text{-}K^{*0}$ to ΛK^+ (0-10% Centrality)