

Λ K femtoscopy in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV measured with ALICE

Jesse T. Buxton¹ on behalf of the ALICE Collaboration

The Ohio State University, Columbus OH 43210, USA,
`jesse.thomas.buxton@cern.ch`

Abstract. The first measurements of the scattering parameters of Λ K pairs in all three charge combinations (ΛK^+ , ΛK^- , and ΛK_S^0) are presented. The results are achieved through a femtoscopic analysis of Λ K correlations in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV recorded by ALICE at the LHC. The femtoscopic correlations result from strong final-state interactions, and are fit with a parametrization allowing for both the characterization of the pair emission source and the measurement of the scattering parameters for the particle pairs.

Keywords: femtoscopy, heavy-ion collisions

1 Introduction

Femtoscopy is an experimental method used to study the space–time characteristics of the particle emitting sources in relativistic particle collisions. With this method, two- (or many-) particle relative-momentum correlation functions are used to connect the final-state momentum distributions to the space–time distributions of particle emission at freeze-out [1]. The correlation functions are sensitive to quantum statistics, as well as strong and Coulomb final-state interactions (FSI). Thus, femtoscopy can offer a unique environment in which to measure nuclear scattering parameters, many of which are difficult, if not impossible, to measure otherwise.

2 Data analysis

This work reports on the analysis of Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV produced by the LHC and measured by the ALICE experiment in 2011. Charged particle tracking was performed using the Time Projection Chamber (TPC) and the Inner Tracking System (ITS). Particle identification for reconstructed tracks was carried out using both the TPC and Time-Of-Flight (TOF) detectors in the pseudorapidity range $|\eta| < 0.8$. The purities of the K^\pm collections are estimated to be $P_{K^\pm} \approx 97\%$. Electrically neutral Λ ($\bar{\Lambda}$) and K_S^0 particles were reconstructed through their weak decays: $\Lambda \rightarrow p\pi^-$ and $K_S^0 \rightarrow \pi^+\pi^-$. The Λ and $\bar{\Lambda}$ purities are estimated to be $P_{\Lambda(\bar{\Lambda})} \approx 95\%$, and that of the K_S^0 is $P_{K_S^0} \approx 98\%$. When forming particle pairs, a shared daughter restriction is applied to ensure the first particle in the pair is unique from the second. Furthermore, an average separation constraint is imposed to remove splitting and merging effects.

3 Analysis methods

The two-particle correlation function is defined as the ratio of the covariant two-particle and single-particle spectra. In practice, the correlation function is formed experimentally as $C(k^*) = \mathcal{N} \frac{A(k^*)}{B(k^*)}$, where $A(k^*)$ is the signal distribution, $B(k^*)$ is the reference distribution, and \mathcal{N} is a normalization parameter. The signal distribution is the same-event distribution of particle pairs, and the reference distribution is obtained using mixed-event pairs [2], i.e., particles from a given event are paired with those from another event.

Theoretically, the ΛK correlation function can be described analytically with a model derived by Lednický and Lyuboshitz [3],

$$C(k^*)_{\text{Lednický}} = 1 + \sum_S \rho_S \left[\frac{1}{2} \left| \frac{f^S(k^*)}{R_{\text{inv}}} \right|^2 \left(1 - \frac{d_0^S}{2\sqrt{\pi}R_{\text{inv}}} \right) + \frac{2\Re f^S(k^*)}{\sqrt{\pi}R_{\text{inv}}} F_1(2k^*R_{\text{inv}}) - \frac{\Im f^S(k^*)}{R_{\text{inv}}} F_2(2k^*R_{\text{inv}}) \right], \quad (1)$$

where $f(k^*)$ is the complex scattering amplitude, F_1 and F_2 are analytic functions, and R_{inv} is the radius of the spherically symmetric Gaussian distribution assumed for the pair emission source in the pair rest frame. The complex scattering amplitude is evaluated via the effective range approximation, $f(k^*) = \left(\frac{1}{f_0} + \frac{1}{2}d_0k^{*2} - ik^* \right)^{-1}$, where f_0 is the complex s-wave scattering length and d_0 is the effective range of the interaction.

Residual correlations The finally measured correlation function is a combination of the genuine ΛK correlation with contributions from impurities and residual correlations resulting from resonance feed-down [4],

$$C_{\text{measured}}(k_{\Lambda K}^*) = 1 + \lambda'_{\Lambda K} [C_{\Lambda K}(k_{\Lambda K}^*) - 1] + \sum_{ij} \lambda'_{ij} [C_{ij}(k_{\Lambda K}^*) - 1], \quad (2)$$

where $\lambda'_{ij} = \lambda_{\text{Fit}} \lambda_{ij}$, the ΛK term represents the genuine ΛK correlation, and the ij terms denote the contributions from impurities and residual correlations. The λ_{ij} parameters serve as weights dictating the relative strength of each component's contribution to the observed signal, and are estimated using the THERMINATOR 2 and HIJING simulations [4, 5]. The net contribution from pairs which contain at least one misidentified member are assumed to average to unity.

The main sources of residual correlations in the ΛK systems result from Λ hyperons which have decayed from Σ^0 , Ξ^0 , and Ξ^- parents. When modeling the $\Sigma^0 K$ and $\Xi^0 K$ systems, for which the interactions are not known, the source radii and scattering parameters are assumed equal to those of the daughter ΛK system. For modeling the $\Xi^- K^\pm$ parent contribution, the available experimental $\Xi^- K^\pm$ data are used. Each residual component, $C_{ij}(k_{\Lambda K}^*)$ in Eq. 2, is the parent correlation function expressed in terms of the relative momentum of the daughter ΛK pair, and is obtained using a transform matrix generated with the THERMINATOR 2 [6] simulation (see [4] for more details).

Non-femtoscopic background A significant non-femtoscopic background is observed in all of the studied ΛK correlations, which is primarily due to particle collimation associated with elliptic flow, and results from mixing events with unlike event planes [7]. These backgrounds are modeled using 6th-order polynomial fits to the THERMINATOR 2 simulation, as shown in Fig. 1, which is then applied as a scale factor in the final fit function.




Fig. 1. THERMINATOR 2 simulation (open triangles) with experimental data (closed circles) for the 30–50% centrality range. Results are shown for ΛK^+ (left), ΛK^- (middle), and ΛK_S^0 (right). A 6th-order polynomial fit to the simulation is shown as a dashed curve. This polynomial scaled to match the experimental data is drawn as a solid curve.

4 Results

Figure 2 shows experimental ΛK correlation functions with fits for the 0–10% centrality percentile interval. All six ΛK systems are fit simultaneously across all centralities, with a single radius and λ_{Fit} parameter for each centrality interval. Scattering parameters ($\Re f_0$, $\Im f_0$, d_0) are shared between pair-conjugate systems, but assumed unique among the different ΛK charge combinations. Figure 3 summarizes the extracted ΛK fit parameters. For all ΛK systems, positive imaginary parts of the scattering lengths, $\Im(f_0)$, describing the inelastic scattering channels, are extracted. More interestingly, the results show that the ΛK^+ and ΛK^- systems differ in the sign of the real part, $\Re(f_0)$, of their scattering lengths, with a negative value for ΛK^+ and positive value for ΛK^- (and ΛK_S^0). The real part of the scattering length describes the effect of the strong interaction, with a positive value signifying an attraction and a negative value signifying a repulsion. Therefore, the results from this analysis indicate that the strong force is repulsive in the ΛK^+ interaction and attractive in the ΛK^- and ΛK_S^0 interactions.

5 Summary

Results from a femtoscopic analysis of ΛK correlations in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 2.76$ TeV measured by the ALICE experiment at the LHC have been




Fig. 2. Fit results for the ΛK data in the 0–10% centrality range; $\Lambda K^+ \oplus \bar{\Lambda} K^-$ are shown in the left column, $\Lambda K^- \oplus \bar{\Lambda} K^+$ in the middle, and $\Lambda K_S^0 \oplus \bar{\Lambda} K_S^0$ in the right. The curves show the primary ΛK contribution to the fit, i.e., $1 + \chi'_{\Lambda K} C_{\Lambda K}$ in Eq. 2 (dotted), the fit to the non-femtoscopic background (dashed), and final fit (solid).




Fig. 3. Extracted fit parameters for all of the ΛK systems. The cross ([A] = [8]) and X (B = [9]) points show theoretical predictions made using chiral perturbation theory.

presented. The femtoscopic radii, λ parameters, and scattering parameters were extracted from one-dimensional correlation functions in terms of the invariant momentum difference. Striking differences are observed in the ΛK^+ , ΛK^- , and ΛK_S^0 correlation functions, and the extracted scattering parameters indicate that the strong force is repulsive in the ΛK^+ interaction and attractive in the ΛK^- and ΛK_S^0 interactions. This effect could be due to different quark–antiquark interactions between the pairs, or from different net strangeness for each system.

References

1. M.A. Lisa, S. Pratt, R. Soltz, U. Wiedemann, *Ann. Rev. Nucl. Part. Sci.* **55** (2005)
2. G.I. Kopylov, *Phys. Lett.* **B50**, 472 (1974)
3. R. Lednický, V.L. Lyuboshitz, *Sov. J. Nucl. Phys.* **35**, 770 (1982)
4. A. Kiesel, H. Zbroszczyk, M. Szymaski, *Phys. Rev.* **C89**(5), 054916 (2014)
5. S. Acharya, et al., *Phys. Rev.* **C99**(2), 024001 (2019)
6. M. Chojnacki, A. Kiesel, W. Florkowski, W. Broniowski, *Comput. Phys. Commun.* **183**, 746 (2012)
7. A. Kiesel, *Acta Physica Polonica B* **48**, 717 (2017)
8. Y.R. Liu, S.L. Zhu, *Phys. Rev.* **D75**, 034003 (2007)
9. M. Mai, P.C. Bruns, B. Kubis, U.G. Meissner, *Phys. Rev.* **D80**, 094006 (2009)