

Part III Combinatorics

Based on lectures by Prof B. Bollobás

Michaelmas 2016
University of Cambridge

Contents

1	Introduction	1
---	--------------	---

1 Introduction

Let X, Y, \dots be sets

Definition. We call $\mathcal{A} \subset \mathcal{P}(X)$ a **set system** or **family of sets**. \mathcal{A} is naturally identified with a bipartite graph $G_{\mathcal{A}}(U, W)$ with $U = \mathcal{A}$, $W = \bigcup_{A \in \mathcal{A}} A$ or $W = X$. Indeed, $Ax \in E(G_{\mathcal{A}}) \iff x \in A$.

Definition. Given $\mathcal{A} \in \mathcal{P}(X)$, a **set of distinct representatives** (SDR) is an injection $f: \mathcal{A} \rightarrow X$ s.t. $f(A) \in A \forall A \in \mathcal{A}$. In its bipartite graph, an SDR corresponds to a complete matching $U \rightarrow W$.

Theorem 1 (Hall, 1935). A set system \mathcal{A} has an SDR if $\forall \mathcal{A}' \subset \mathcal{A}$, $|\bigcup_{A \in \mathcal{A}'} A| \geq |\mathcal{A}'|$.

Theorem 1'. A bipartite graph $G(U, W)$ has a complete matching $U \rightarrow W$ if $\forall S \subset U$, $|\Gamma(S)| \geq |S|$

Corollary 2. Suppose $G(U, W)$ bipartite, $d(u) \geq d(w) \forall u \in U, w \in W$. Then \exists a complete matching $U \rightarrow W$.