

305 Lecture 49 - Truth in a Model

Brian Weatherson

August 3, 2020

- To extend our discussion of truth at a world, to discussion of truth in a whole model.

Associated Reading

- Boxes and Diamonds, section 3.5.

Models

Models have three parts:

1. A set of worlds, typically called W .

Models

Models have three parts:

1. A set of worlds, typically called W .
2. A binary accessibility relation on those worlds, typically called R .

Models have three parts:

1. A set of worlds, typically called W .
2. A binary accessibility relation on those worlds, typically called R .
3. A valuation function on those worlds, typically called V .

Models

Models have three parts:

1. A set of worlds, typically called W .
2. A binary accessibility relation on those worlds, typically called R .
3. A valuation function on those worlds, typically called V .

We'll write the models as $\langle W, R, V \rangle$.

Valuations

V is a function from atomic sentence letters to subsets of W .

- It tells you when the atomic sentences are true.
- When an atomic sentence is not true, it is false.

Truth at a Point

The general theory of truth is built up in stages from the basic theory. Assume we have a model $\langle W, R, V \rangle$, and a point $w \in W$, and are asking whether an arbitrary sentence is true at w in $\langle W, R, V \rangle$.

- p is true at w iff $w \in V(p)$.

Truth at a Point

The general theory of truth is built up in stages from the basic theory. Assume we have a model $\langle W, R, V \rangle$, and a point $w \in W$, and are asking whether an arbitrary sentence is true at w in $\langle W, R, V \rangle$.

- p is true at w iff $w \in V(p)$.
- $\neg A$ is true at w iff A is not true at w .
- $A \wedge B$ is true at w iff A is true at w and B is true at w .
- $A \vee B$ is true at w iff A is true at w or B is true at w .
- $A \rightarrow B$ is true at w iff A is false at w or B is true at w .

Truth at a Point

The general theory of truth is built up in stages from the basic theory. Assume we have a model $\langle W, R, V \rangle$, and a point $w \in W$, and are asking whether an arbitrary sentence is true at w in $\langle W, R, V \rangle$.

- p is true at w iff $w \in V(p)$.
- $\neg A$ is true at w iff A is not true at w .
- $A \wedge B$ is true at w iff A is true at w and B is true at w .
- $A \vee B$ is true at w iff A is true at w or B is true at w .
- $A \rightarrow B$ is true at w iff A is false at w or B is true at w .

This just leaves the modal formulae. I'll set out the rules, then do some worked examples.

Necessary Truth at a Point

First we'll do $\Box A$.

- I'll read this as 'Box A'.

Necessary Truth at a Point

First we'll do $\Box A$.

- I'll read this as 'Box A'.
- Intuitively, it means **It must be that A**, where **must** could be a metaphysical necessity, or an epistemic necessity, or a moral necessity, or anything else.

Necessary Truth at a Point

First we'll do $\Box A$.

- I'll read this as 'Box A'.
- Intuitively, it means **It must be that A**, where **must** could be a metaphysical necessity, or an epistemic necessity, or a moral necessity, or anything else.
- And it is true at w just in case A is true at every world y such that wRy .
- Necessary truth is truth at all accessible worlds.

Possible Truth at a Point

Now we'll do $\Diamond A$.

- I'll read this as 'Diamond A'.

Possible Truth at a Point

Now we'll do $\Diamond A$.

- I'll read this as 'Diamond A'.
- Intuitively, it means **It might be that A**, where **might** could be a metaphysical necessity, or an epistemic necessity, or a moral necessity, or anything else.

Possible Truth at a Point

Now we'll do $\Diamond A$.

- I'll read this as 'Diamond A'.
- Intuitively, it means **It might be that A**, where **might** could be a metaphysical necessity, or an epistemic necessity, or a moral necessity, or anything else.
- And it is true at w just in case A is true at some world y such that wRy .
- Possible truth is truth at some accessible world.

We can run these rules in sequence.

We can run these rules in sequence.

- What does it take for $\Box \Box A$ to be true at w ?

We can run these rules in sequence.

- What does it take for $\Box \Box A$ to be true at w ?
- It is for $\Box A$ to be true at every y such that wRy .

We can run these rules in sequence.

- What does it take for $\Box \Box A$ to be true at w ?
- It is for $\Box A$ to be true at every y such that wRy .
- And that means that A has to be true at every world z such that yRz (for any y such that wRy).

We can think, a little picturesquely, as the accessibility relation being a 'step' between worlds.

- If wRy , then you can 'step' from w to y .

We can think, a little picturesquely, as the accessibility relation being a 'step' between worlds.

- If wRy , then you can 'step' from w to y .
- $\Box A$ means that anywhere you can step to from w is a world where A is true.

We can think, a little picturesquely, as the accessibility relation being a 'step' between worlds.

- If wRy , then you can 'step' from w to y .
- $\Box A$ means that anywhere you can step to from w is a world where A is true.
- And $\Box \Box A$ means that anywhere you can get to in two steps from w is a world where A is true.

Iterated Modalities

We can run the rules in sequence.

We can run the rules in sequence.

- What does it take for $\Diamond \Diamond A$ to be true at w ?

We can run the rules in sequence.

- What does it take for $\Diamond \Diamond A$ to be true at w ?
- It is for $\Diamond A$ to be true at some y such that wRy .

We can run the rules in sequence.

- What does it take for $\Diamond \Diamond A$ to be true at w ?
- It is for $\Diamond A$ to be true at some y such that wRy .
- And that means that A has to be true at some world z such that yRz (for some y such that wRy).

We can run the rules in sequence.

- What does it take for $\Diamond \Diamond A$ to be true at w ?
- It is for $\Diamond A$ to be true at some y such that wRy .
- And that means that A has to be true at some world z such that yRz (for some y such that wRy).
- In the picturesque terms, you can get from w to an A -world in two steps.

Mixed Modalities

What does it mean for $\Diamond \Box A$ to be true at w ?

What does it mean for $\Diamond \Box A$ to be true at w ?

- There is some accessible world where $\Box A$ is true.

What does it mean for $\Diamond \Box A$ to be true at w ?

- There is some accessible world where $\Box A$ is true.
- That is, there is some accessible world such that everywhere you can go from there, A is true.

Mixed Modalities

What does it mean for $\Box \Diamond A$ to be true at w ?

What does it mean for $\Box \Diamond A$ to be true at w ?

- At all accessible worlds, $\Diamond A$ is true.

What does it mean for $\Box \Diamond A$ to be true at w ?

- At all accessible worlds, $\Diamond A$ is true.
- That is, wherever you go, you can get to there is some accessible world such that everywhere you can go from there, A is true.

Longer Sentences

What does it mean for $\Box(p \vee (q \rightarrow \Diamond r))$ to be true at w ?

Longer Sentences

What does it mean for $\Box(p \vee (q \rightarrow \Diamond r))$ to be true at w ?

- It's for $p \vee (q \rightarrow \Diamond r)$ to be true everywhere you can get to (in one step) from w .
- That is, at every one of those worlds, either p is true or $q \rightarrow \Diamond r$ is true.

Longer Sentences

What does it mean for $\Box(p \vee (q \rightarrow \Diamond r))$ to be true at w ?

- It's for $p \vee (q \rightarrow \Diamond r)$ to be true everywhere you can get to (in one step) from w .
- That is, at every one of those worlds, either p is true or $q \rightarrow \Diamond r$ is true.
- That is, at every one of those worlds, either p is true, or q is false, or $\Diamond r$ is true.

Longer Sentences

What does it mean for $\Box(p \vee (q \rightarrow \Diamond r))$ to be true at w ?

- It's for $p \vee (q \rightarrow \Diamond r)$ to be true everywhere you can get to (in one step) from w .
- That is, at every one of those worlds, either p is true or $q \rightarrow \Diamond r$ is true.
- That is, at every one of those worlds, either p is true, or q is false, or $\Diamond r$ is true.
- That is, at every one of those worlds, either p is true, or q is false, or there is some world you can get to where r is true.

Box and connectives

The general rule is just to apply the rules for sentences inside the brackets at each world in W , and then apply the rule for \Box or \Diamond . But there are three special cases worth thinking about.

- $\Box(A \wedge B)$ means that all accessible worlds are A and B worlds.

Box and connectives

The general rule is just to apply the rules for sentences inside the brackets at each world in W , and then apply the rule for \Box or \Diamond . But there are three special cases worth thinking about.

- $\Box(A \wedge B)$ means that all accessible worlds are A and B worlds.
- $\Box(A \vee B)$ means that all accessible worlds make at least one of A and B true.

Box and connectives

The general rule is just to apply the rules for sentences inside the brackets at each world in W , and then apply the rule for \Box or \Diamond . But there are three special cases worth thinking about.

- $\Box(A \wedge B)$ means that all accessible worlds are A and B worlds.
- $\Box(A \vee B)$ means that all accessible worlds make at least one of A and B true.
- $\Box(A \rightarrow B)$ means that all accessible A -worlds are B -worlds.

We'll use that last one a lot.

For Next Time

We'll discuss of examples of truth (and non-truth) in models to explain this material.