MULTIPARTITION SPACES, KIRWAN'S EQUIVARIANT PERFECTION, AND HAUSEL'S FORMULA

ABSTRACT. These notes give a self-contained account of the geometric part of Section 6 of [Hik]. For a quiver with a loop at every vertex and a multipartition m, we define the multipartition quiver Γ_m, the multipartition space E_m and its open subset U_m, the torus T_m, the toric hyperKähler variety Y_m, and the action of the group W_m. We then prove [Hik, Proposition 6.2] as an instance of Kirwan's equivariant perfection argument: the decomposition of E_m by the set of unreachable slots is exactly Hesselink's stratification for the determinant character, in the terminology of Hoskins [Hos]. Finally we recall Hausel's formula and explain, following [Hik, Corollary 6.3], how Proposition 6.2 computes the ratio of its two sums.

This note was written by Claude Opus 5.5 (Anthropic) at the request of Hiraku Nakajima, based on a discussion with him in October 2026. A record of the discussion is available on request.

CONTENTS

Conventions 1
1. Multipartition spaces 1
2. Proposition 6.2 via Kirwan's equivariant perfection 8
3. Hausel's formula and Corollary 6.3 13
References 17

CONVENTIONS

Equation, proposition and corollary numbers of the form (6.1), Proposition 6.2, Corollary 6.3 refer to [Hik]. All other numbers are cross-references within this note and are hyperlinks to the corresponding statement or equation.

Cohomology has complex coefficients, and equivariant cohomology is Borel equivariant cohomology. All graded vector spaces below are concentrated in even degrees, finite-dimensional in each degree and bounded below; for such a space we put

$$\mathrm{Hilb}(M;q) = \sum_d \dim M^{2d}\, q^d.$$

Thus q records *half* the cohomological degree. If X has no odd cohomology, its Poincaré polynomial is $P_X(q) = \mathrm{Hilb}(H^*(X);q)$, and $P_\emptyset = 0$.

For geometric invariant theory of a linear action on an affine space with respect to a character we follow the terminology and notation of Hoskins [Hos, §2]. In §1.7 and Section 2, λ denotes a one-parameter subgroup; elsewhere $\lambda = (\lambda^i)$ denotes a tuple of partitions.

In [Hik] the polynomials h_m defined below bound the graded pieces of a filtration of the Coulomb-side ring B_0 [Hik, Proposition 5.2]; Hausel's formula shows that the bound is sharp [Hik, Proposition 6.4, Corollary 6.6]. We do not revisit that part of the argument.

Date: October 2026.

1. Multipartition spaces

1.1. **The quiver.** Let $Q = (Q_0, Q_1)$ be a finite quiver with at least one loop at every vertex. For an arrow $a \in Q_1$ we write $t(a)$ and $h(a)$ for its tail and head. Fix one loop ℓ_i at each vertex i, and let Q' be the quiver obtained from Q by deleting the chosen loops. Put

$$n_{ij} = \#\{a \in Q_1' : t(a) = i,\ h(a) = j\}, \qquad c_{ij} = n_{ij} + n_{ji}. \tag{1.1}$$

The matrix (c_{ij}) is symmetric with nonnegative integer entries, and $c_{ii} = 2n_{ii}$ is even. Fix a framing vector $w \in \mathbb{Z}_{\geq 0}^{Q_0}$. For $v \in \mathbb{Z}_{\geq 0}^{Q_0}$ put

$$C(v) = \tfrac{1}{2}\sum_{i,j} c_{ij} v_i v_j = \sum_{a \in Q_1'} v_{t(a)} v_{h(a)}. \tag{1.2}$$

1.2. **Multipartitions.**

Definition 1.1. A *multipartition* is a family $m = (m_{i,s})_{i \in Q_0,\, s \geq 1}$ of nonnegative integers, only finitely many of which are nonzero. Its *dimension vector* $\|m\| \in \mathbb{Z}_{\geq 0}^{Q_0}$ is $\|m\|_i = \sum_s s\, m_{i,s}$, and m is called a multipartition of v if $\|m\| = v$. Let $\mathcal{M}$ be the set of all multipartitions (of all dimension vectors) and $\mathcal{M}(v) \subset \mathcal{M}$ the finite subset of multipartitions of v. The empty multipartition $\emptyset$ is the unique multipartition of 0.

Equivalently, m is a Q_0-tuple $\lambda = (\lambda^i)_{i \in Q_0}$ of partitions, where $m_{i,s}$ is the number of parts of λ^i equal to s; we write $m = m(\lambda)$ and use both descriptions.

Definition 1.2. For $k, m \in \mathcal{M}$ put

$$l_i(m) = \textstyle\sum_s m_{i,s} \text{ (the number of parts at } i), \qquad l(m) = (l_i(m))_i, \qquad |l(m)| = \textstyle\sum_i l_i(m), \tag{1.3}$$

$$w(m) = w \cdot l(m) = \textstyle\sum_i w_i\, l_i(m), \qquad (q)_m = \prod_{i,s} \prod_{j=1}^{m_{i,s}} (1 - q^j), \tag{1.4}$$

$$b(k,m) = \sum_{i,j \in Q_0} \sum_{s,t \geq 1} c_{ij} \min(s,t)\, k_{i,s}\, m_{j,t}, \qquad D_m = \tfrac{1}{2}\, b(m,m). \tag{1.5}$$

The form b is bilinear and symmetric. For partitions put $\langle \mu, \nu \rangle = \sum_{a,b} \min(\mu_a, \nu_b)$; then

$$D_{m(\lambda)} = \tfrac{1}{2}\sum_{i,j} c_{ij} \langle \lambda^i, \lambda^j \rangle = \sum_{a \in Q_1'} \langle \lambda^{t(a)}, \lambda^{h(a)} \rangle \in \mathbb{Z}_{\geq 0}. \tag{1.6}$$

Multipartitions are added entrywise, and $\|k + m\| = \|k\| + \|m\|$. We write $k \leq m$ if $k_{i,s} \leq m_{i,s}$ for all (i,s); then $m - k \in \mathcal{M}$. Bilinearity gives

$$D_{k+m'} = D_k + D_{m'} + b(k, m'), \qquad w(k + m') = w(k) + w(m'). \tag{1.7}$$

1.3. **The multipartition quiver and slots.**

Definition 1.3. Let $m \in \mathcal{M}$ and put $\Omega_m = \{(i,s,\alpha) : i \in Q_0,\ s \geq 1,\ 1 \leq \alpha \leq m_{i,s}\}$. The *multipartition quiver* Γ_m has vertex set $\Omega_m \sqcup \{\infty\}$. The vertex ∞ is the *framing vertex*; the other vertices, i.e. the elements of Ω_m, are called *slots*. A slot $d = (i,s,\alpha)$ has *color* $i(d) = i$, *size* $s(d) = s$ and *type* (i,s). For distinct slots $d \neq e$ put

$$b_{de} = c_{i(d)i(e)} \min(s(d), s(e)) = b_{ed}. \tag{1.8}$$

The arrows of Γ_m are:

- for every ordered pair (d,e) of distinct slots, b_{de} arrows $d \to e$, labeled $(d \to e, p)$ with $1 \leq p \leq b_{de}$;
- for every slot d, $w_{i(d)}$ *framing arrows* $\infty \to d$, labeled $(\infty \to d, p)$ with $1 \leq p \leq w_{i(d)}$.

There are $m_{i,s}$ slots of type (i,s) and $|l(m)|$ slots in all; the sizes of the slots of color i are the parts of λ^i. By (1.8), the full subquiver on the slots is *symmetric*: it has as many arrows $d \to e$ as $e \to d$. There are no arrows into ∞ and no loops.

The group

$$W_m = \prod_{i,s} \mathfrak{S}_{m_{i,s}} \tag{1.9}$$

acts on Ω_m by $\sigma(i,s,\alpha) = (i,s,\sigma_{i,s}(\alpha))$. Thus W_m is the group of permutations of the slots preserving types, and $b_{\sigma d,\sigma e} = b_{de}$.

For $S \subseteq \Omega_m$ its *type* is the multipartition $\kappa(S) = k \le m$ with $k_{i,s} = \#\{d \in S \text{ of type } (i,s)\}$; the complement $\Omega_m \setminus S$ has type $m-k$. Two subsets lie in the same W_m-orbit if and only if they have the same type, and the stabilizer of S is $W_k \times W_{m-k}$. We identify

$$\Omega_m \setminus S \;\cong\; \Omega_{m-k} \tag{1.10}$$

by the type-preserving, order-preserving relabeling of the index α; this identification preserves the numbers b_{de}. By definition of b in (1.5),

$$b(\kappa(S),\, m-\kappa(S)) = \sum_{d\in S} \sum_{e\in\Omega_m\setminus S} b_{de}, \qquad D_m = \sum_{\{d,e\},\, d\neq e} b_{de} + \sum_{d\in\Omega_m} n_{i(d)i(d)}\, s(d). \tag{1.11}$$

In the second formula the sum is over unordered pairs; the last term comes from the pairs $d = e$ in $\frac{1}{2}b(m,m)$ and vanishes when Q' has no loops.

1.4. **The multipartition space.**

Definition 1.4. The *multipartition space* E_m is the space of representations of the multipartition quiver Γ_m (Definition 1.3) with a one-dimensional space $\mathbb{C}$ at every vertex:

$$E_m = E_m^{\mathrm{int}} \oplus F_m, \qquad E_m^{\mathrm{int}} = \bigoplus_{d\neq e} \mathbb{C}^{b_{de}}, \qquad F_m = \bigoplus_{d\in\Omega_m} \mathbb{C}^{w_{i(d)}}.$$

We write a point as $B = (B', a)$, where $B' = (B^{(p)}_{d\to e}) \in E_m^{\mathrm{int}}$ consists of the components $B^{(p)}_{d\to e}$, $1 \le p \le b_{de}$, for the arrows between slots, and $a = (a_d^{(p)}) \in F_m$ consists of the components $a_d^{(p)}$, $1 \le p \le w_{i(d)}$, for the framing arrows.

Thus $\dim E_m = 2\sum_{\{d,e\}} b_{de} + w(m)$, and $E_\emptyset$ is a point.

1.5. **The torus and the cut ideal.**

The torus $T_m = (\mathbb{C}^\times)^{\Omega_m}$ acts on E_m by change of basis at the slots, trivially at ∞:

$$t\cdot(B',a): \quad B^{(p)}_{d\to e} \mapsto t_e\, t_d^{-1}\, B^{(p)}_{d\to e}, \qquad a_d^{(p)} \mapsto t_d\, a_d^{(p)}. \tag{1.12}$$

Let $x_d \in H^2_{T_m}(\mathrm{pt})$ be the first Chern class of the character $t \mapsto t_d$, so that

$$P_m := H^*_{T_m}(\mathrm{pt}) = \mathbb{C}[x_d : d \in \Omega_m], \qquad \deg x_d = 2.$$

By (1.12) the weights of E_m are

$$x_e - x_d \;\text{ on each coordinate } B^{(p)}_{d\to e}, \qquad x_d \;\text{ on each coordinate } a_d^{(p)}. \tag{1.13}$$

Since E_m is contractible, $H^*_{T_m}(E_m) = P_m$.

Definition 1.5. For a nonempty subset $S \subseteq \Omega_m$, the *cut polynomial* [Hik, (5.3)] is

$$q_S = \prod_{d\in S} x_d^{\,w_{i(d)}} \prod_{d\in S,\; e\notin S} (x_d - x_e)^{b_{de}} \in P_m.$$

The *cut ideal* $J_m \subseteq P_m$ is the ideal generated by all q_S, and

$$h_m(q) = \mathrm{Hilb}\left((P_m/J_m)^{W_m}; q\right),$$

where W_m acts on P_m/J_m as in §1.6.

We keep the notation of [Hik]; q_S has nothing to do with the variable q. By [Hik, Lemma 6.1], P_m/J_m is finite-dimensional, so h_m is a polynomial. This also follows from Remark 1.12(iv): there P_m/J_m is identified, by Konno's theorem [Kon00, Theorem 3.1], with the cohomology of the toric hyperKähler manifold Y_m, which is finite-dimensional. Moreover the Betti numbers b_{2k} of Y_m are the h-numbers h_k, $0 \le k \le g_m$, of a matroid of rank g_m [HS, Corollaries 4.6 and 6.6], where g_m is as in Remark 1.12(ii). Hence P_m/J_m vanishes in degrees above $2g_m$, and h_m has degree at most g_m. For $m = \emptyset$ we have $P_\emptyset = \mathbb{C}$, $J_\emptyset = 0$ and $h_\emptyset = 1$.

1.6. **The action of W_m.** The arrow labels in Definition 1.3 depend only on the types of the endpoints, so the group W_m of (1.9) acts on Γ_m, and hence on E_m and on T_m:

$$(\sigma B)^{(p)}_{\sigma d \to \sigma e} = B^{(p)}_{d\to e}, \qquad (\sigma a)^{(p)}_{\sigma d} = a^{(p)}_d, \qquad (\sigma t)_{\sigma d} = t_d.$$

These satisfy $\sigma(t\cdot B) = (\sigma t)\cdot(\sigma B)$, so the semidirect product $T_m \rtimes W_m$ acts on E_m. (It is the normalizer of the maximal torus in $\prod_{i,s} GL_{m_{i,s}}$.) If $\mathcal{X} \subseteq E_m$ is stable under $T_m \rtimes W_m$, each σ gives an equivariant automorphism of $(\mathcal{X}, T_m)$; letting σ act by $(\sigma^{-1})^*$ defines a left action of W_m on $H^*_{T_m}(\mathcal{X})$ by graded algebra automorphisms. On $H^*_{T_m}(\mathrm{pt}) = P_m$ it is $\sigma \cdot x_d = x_{\sigma d}$. We record three facts.

(a) Since W_m is finite and coefficients are complex, taking W_m-invariants is exact, and

$$H^*_{T_m\rtimes W_m}(\mathcal{X}) = H^*_{T_m}(\mathcal{X})^{W_m}.$$

(b) $\sigma \cdot q_S = q_{\sigma S}$, hence J_m is W_m-stable and W_m acts on P_m/J_m.

(c) $P_m^{W_m}$ is the tensor product over (i,s) of rings of symmetric polynomials in $m_{i,s}$ variables, so $\mathrm{Hilb}(P_m^{W_m}; q) = 1/(q)_m$, with $(q)_m$ as in (1.4).

1.7. **Stability for the torus action.** We use the notation of [Hos, §2], specialized to the torus T_m acting on E_m. Characters and one-parameter subgroups (1-PSs) of T_m are identified with vectors in $\mathbb{Z}^{\Omega_m}$: a character χ is $t \mapsto \prod_d t_d^{\chi_d}$, a 1-PS λ is $\tau \mapsto (\tau^{\lambda_d})_d$, and the pairing is $(\chi,\lambda) = \sum_d \chi_d\lambda_d$. We write characters additively, so x_d is the character $t \mapsto t_d$, and we use the standard norm $\|\lambda\|^2 = \sum_d \lambda_d^2$, which is invariant under W_m and satisfies [Hos, Assumption 2.9]. For $A \subseteq \Omega_m$ let $1_A \in \mathbb{Z}^{\Omega_m}$ have entries 1 on A and 0 elsewhere. Throughout, the character is

$$\rho = \det = 1_{\Omega_m} :\ T_m \to \mathbb{C}^\times, \qquad t \mapsto \prod_d t_d. \tag{1.14}$$

In King's notation for the quiver Γ_m, this is the stability parameter which is 1 at every slot [Hos, §5.1].

Definition 1.6 ([Hos, Definition 2.1])**.** A point $B \in E_m$ is *ρ-semistable* if there are $n > 0$ and $f \in \mathbb{C}[E_m]$ with $f(t\cdot B'') = \rho(t)^n f(B'')$ for all $t \in T_m$, $B'' \in E_m$, and $f(B) \neq 0$. It is *ρ-stable* if moreover its stabilizer is finite and T_m acts on $\{f \neq 0\}$ with closed orbits, for some such f. Otherwise B is *ρ-unstable*. We write $E_m^{\rho\text{-}s} \subseteq E_m^{\rho\text{-}ss}$ for the open subsets of ρ-stable and ρ-semistable points, and $E_m /\!/_\rho T_m$ for the GIT quotient.

For $B \in E_m$ let $\mathrm{wt}(B)$ be the set of weights χ in (1.13) such that the component of B in the χ-weight space is nonzero. Following [Hos, (4)], the *cone of allowable 1-PSs* of B is

$$C_B = \bigcap_{\chi\in \mathrm{wt}(B)} H_\chi \subseteq \mathbb{R}^{\Omega_m}, \qquad H_\chi = \{\lambda : (\chi,\lambda) \ge 0\}. \tag{1.15}$$

A 1-PS λ lies in C_B if and only if $\lim_{\tau\to 0}\lambda(\tau)\cdot B$ exists. Put $H_\rho = \{\lambda : (\rho,\lambda)\geq 0\}$ and $H_\rho^\circ = \{\lambda : (\rho,\lambda) > 0\}$.

Proposition 1.7 (Hilbert–Mumford criterion [Hos, §2.1])**.** *Let $B \in E_m$ and recall C_B from (1.15). Then B is ρ-semistable if and only if $C_B \subseteq H_\rho$, and ρ-stable if and only if $C_B\setminus\{0\} \subseteq H_\rho^\circ$.*

1.8. **Reachability and U_m.**

Definition 1.8. Let $B = (B', a) \in E_m$ (Definition 1.4). A slot d is *reachable* in B if there are slots $d_1, \dots, d_r = d$ ($r \geq 1$) and labels such that $a_{d_1}^{(p_0)} \neq 0$ and $B_{d_j\to d_{j+1}}^{(p_j)} \neq 0$ for $1 \leq j < r$. Let $R(B)$ be the set of reachable slots and $S(B) = \Omega_m \setminus R(B)$ the set of unreachable ones. Put

$$U_m = \{B \in E_m : R(B) = \Omega_m\}.$$

For $S \subseteq \Omega_m$ let $\lambda_S = -1_S$ be the 1-PS with $\lambda_S(\tau)_d = \tau^{-1}$ for $d \in S$ and 1 otherwise; $\lambda_\emptyset$ is the trivial 1-PS.

"d is reachable" is a finite union of non-vanishing conditions, so U_m is open; it is stable under $T_m \rtimes W_m$ (§1.6).

Lemma 1.9. *Let $B \in E_m$. Recall the sets $R(B)$, $S(B)$ and the 1-PS λ_S of Definition 1.8, and the cone C_B of (1.15).*

(1) Every $\lambda \in C_B$ satisfies $\lambda_d \geq 0$ for all $d \in R(B)$.
(2) $\lambda_{S(B)} \in C_B$, and $(\rho, \lambda_{S(B)}) = -|S(B)|$.

Proof. (1) Take a path $d_1, \dots, d_r = d$ as in Definition 1.8. By (1.13), $x_{d_1} \in \mathrm{wt}(B)$ and $x_{d_{j+1}} - x_{d_j} \in \mathrm{wt}(B)$, so $\lambda_{d_1} \geq 0$ and $\lambda_{d_{j+1}} \geq \lambda_{d_j}$; hence $\lambda_d \geq 0$.

(2) Put $S = S(B)$ and $R = R(B)$. A nonzero framing coordinate a_d with $d \in S$, or a nonzero coordinate $B_{e\to d}$ with $e \in R$, $d \in S$, would make d reachable; so these vanish. The remaining weights in $\mathrm{wt}(B)$ are x_d with $d \in R$, $x_e - x_d$ with d, e both in R or both in S, and $x_e - x_d$ with $d \in S$, $e \in R$. Their pairings with $\lambda_S = -1_S$ are 0, 0 and 1, all ≥ 0. Finally $(\rho, \lambda_S) = -|S|$ by (1.14). □

Lemma 1.10. *Let $B \in E_m$, and recall ρ-(semi)stability from Definition 1.6.*

(1) The smallest subrepresentation of B containing the space at ∞ is supported on $\{\infty\}\cup R(B)$. Hence $B \in U_m$ if and only if B is cyclic, i.e. generated by the space at ∞.
(2) $U_m = E_m^{\rho\text{-}ss} = E_m^{\rho\text{-}s}$.
(3) T_m acts freely on U_m. The quotient $U_m/T_m = E_m/\!/_\rho T_m$ is a smooth quasi-projective toric variety of dimension $\dim E_m - |l(m)|$ (if nonempty), and

$$H^*_{T_m}(U_m) = H^*(U_m/T_m), \qquad H^*_{T_m}(U_m)^{W_m} = H^*\big((U_m/T_m)/W_m\big).$$

Proof. (1) All vertex spaces are one-dimensional, so a subrepresentation is a subset of vertices closed under nonzero arrows. The smallest one containing ∞ is $\{\infty\} \cup R(B)$.

(2) If $B \in U_m$, then every $0 \neq \lambda \in C_B$ has $\lambda_d \geq 0$ for all d by Lemma 1.9(1), hence $(\rho, \lambda) = \sum_d \lambda_d > 0$. By Proposition 1.7, B is ρ-stable. If $S(B) \neq \emptyset$, then Lemma 1.9(2) gives $\lambda_{S(B)} \in C_B$ with $(\rho, \lambda_{S(B)}) < 0$, so $C_B \not\subseteq H_\rho$ and B is ρ-unstable.

(3) If $t\cdot B = B$, then $t_d = 1$ whenever $a_d \neq 0$, and $t_e = t_d$ whenever $B_{d\to e} \neq 0$; following paths from ∞ gives $t = 1$. By (2) and [Hos, Theorem 2.2], $U_m \to E_m/\!/_\rho T_m$ is a geometric quotient, smooth because the action is free. The torus $(\mathbb{C}^\times)^{\dim E_m}$ scaling all coordinates commutes with T_m and preserves U_m, since reachability only depends on which coordinates are nonzero. If $U_m \neq \emptyset$, it contains the open orbit where all coordinates are nonzero, so U_m/T_m is toric. The last assertions hold because the action is free and W_m is finite. □

1.9. **The toric hyperKähler variety Y_m.** Choose an *orientation* o: for every unordered pair $\{d,e\}$ of distinct slots and every $1 \le p \le b_{de}$, declare one of $(d \to e, p)$, $(e \to d, p)$ to be *original*. Let V_m^o be the framed quiver with vertex set $\Omega_m \sqcup \{\infty\}$, the original arrows and the framing arrows, and let

$$L_m^o = \bigoplus_{\text{original arrows}} \mathbb{C} \;\oplus\; F_m$$

be its representation space with dimension vector 1. The coordinate of the reverse $\bar\gamma$ of an original arrow γ has the opposite weight, so it is naturally dual to the coordinate B_γ. Hence

$$T^*L_m^o = L_m^o \oplus (L_m^o)^\vee = E_m \oplus F_m^\vee, \qquad F_m^\vee = \bigoplus_d \left(\mathbb{C}^{w_{i(d)}}\right)^\vee,$$

where the coordinates $b_d = (b_d^{(p)})_p$ of $F_m^\vee$ are dual framing arrows $d \to \infty$, of weight $-x_d$. Following the recent literature we write b for the dual framing; it should not be confused with the integers b_{de} of (1.8) or the bilinear form $b(k,m)$ of (1.5). The moment map $\mu : E_m \oplus F_m^\vee \to \mathrm{Lie}(T_m)^\vee = \mathbb{C}^{\Omega_m}$, defined by $\langle \mu(s,p), \xi\rangle = \langle p, \xi \cdot s\rangle$ as in [Hik], is

$$\mu_d(B,b) = \sum_{\substack{\gamma \text{ original}\\ h(\gamma)=d}} B_\gamma B_{\bar\gamma} - \sum_{\substack{\gamma \text{ original}\\ t(\gamma)=d}} B_\gamma B_{\bar\gamma} + \langle b_d, a_d\rangle, \qquad \langle b_d, a_d\rangle = \sum_p b_d^{(p)} a_d^{(p)}.$$

Definition 1.11. $Y_m = \{(B,b) \in U_m \times F_m^\vee : \ \mu(B,b) = 0\}/T_m$, where $B = (B', a) \in U_m$ and $b \in F_m^\vee$.

Remark 1.12. (i) Y_m is the quiver variety [Nak94] of V_m^o with dimension vector $(1,\dots,1)$ and framing w, in its holomorphic description at determinant stability (cf. [Nak94, Theorem 3.2, Proposition 3.5]). The multipartition quiver has no edge loops, so this is within the setting of [Nak94]. Indeed, by the argument of Lemma 1.10(1), cyclicity in the doubled quiver means reachability using the framing arrows $\infty \to d$ and the arrows between slots in both directions; that is exactly the condition $B \in U_m$, and the dual framing b plays no role.

(ii) T_m acts freely on $U_m \times F_m^\vee$ (Lemma 1.10(3)), so μ is a submersion there and Y_m is smooth of dimension

$$\dim Y_m = 2g_m, \qquad g_m = \sum_{\{d,e\}} b_{de} + w(m) - |l(m)|.$$

Here g_m is the number of edges minus the number of slots of the underlying graph of Γ_m, i.e. its first Betti number when that graph is connected.

(iii) Changing the orientation changes the sign of some terms of μ. Changing the sign of the corresponding reverse coordinates is a T_m-equivariant isomorphism, so Y_m does not depend on o up to isomorphism. Composing with such sign changes, W_m acts on Y_m.

(iv) *Relation with Konno* [Kon00, Kon01] *and Hausel–Sturmfels* [HS]. Let A be the integer matrix with $|l(m)|$ rows whose columns are the T_m-weights (1.13) of the coordinates of L_m^o, one column for each original arrow and each framing arrow. This is the boundary map [HS, (40)] of the quiver V_m^o, with $v_0 = \infty$; it is unimodular [HS, Lemma 8.1], and T_m plays the role of the torus $\mathbb{T}_{\mathbb{C}}^d$ of [HS]. The character $\rho = \det$ of (1.14) is the vector $\theta = (1,\dots,1)$ of [HS]. A hyperplane spanned by columns of A is spanned by the arrows of a forest with two components; if $S \subseteq \Omega_m$ is the component not containing ∞, the hyperplane is $\{y : \sum_{d\in S} y_d = 0\}$. Since $\sum_{d\in S}\theta_d = |S| \neq 0$, θ is generic in the sense of [HS, §4]. In this notation:

- The coordinates z_i, w_i of [HS, §4] are the coordinates of L_m^o and their duals, so the Lawrence toric variety of [HS, Definition 4.1, Theorem 1.1(2)] is the GIT quotient of

$T^*L^o_m = E_m \oplus F_m^\vee$ without the moment map equation:

$$X_m^{\mathrm{Law}} := X(A^\pm, \theta) = (E_m \oplus F_m^\vee) /\!/_\rho T_m.$$

- Y_m is the toric quiver variety $Y(V^o_m, \theta) = Y(A, \theta)$ of [HS, §8]: the subvariety of X_m^{Law} cut out by the components of the complex moment map [HS, (29), Definition 6.1], biholomorphic to the hyperKähler quotient [HS, Theorem 6.3]. (In [HS, §6] the Gale dual matrix is assumed to have no zero row, i.e. the underlying graph has no bridge, so that the ideal (29) is prime; in general we take Definition 1.11 as the definition.)
- X_m^{Law} is a vector bundle over the toric manifold U_m/T_m, whose fiber consists of the linear maps *to* the framing vector spaces. Indeed, the coordinates b_d of $F_m^\vee$ are arrows $d \to \infty$, which do not help to reach slots from ∞; so Lemma 1.9(1) still holds for the cone of allowable 1-PSs of (B, b). Their weights $-x_d$ pair with $\lambda_{S(B)}$ to 0 or 1, so Lemma 1.9(2) also holds. The proof of Lemma 1.10(2) then shows that the semistable locus is $U_m \times F_m^\vee$, on which T_m acts freely. Therefore

$$X_m^{\mathrm{Law}} = U_m \times^{T_m} F_m^\vee = \bigoplus_{d \in \Omega_m} \mathrm{Hom}\left(\mathcal{L}_d,\, \mathbb{C}^{w_{i(d)}}\right),$$

  where $\mathcal{L}_d$ is the line bundle on U_m/T_m associated with the character $t \mapsto t_d$ (the tautological line at the slot d), and $\mathbb{C}^{w_{i(d)}}$ is a trivial bundle. The zero section is U_m/T_m, and $Y_m \subset X_m^{\mathrm{Law}}$ is the closed subvariety $\mu = 0$. For a single slot with w framing arrows this is the example in [HS, §1]: a rank w bundle over $\mathbb{P}^{w-1}$ containing $Y_m = T^*\mathbb{P}^{w-1}$.
- By [HS, Theorem 1.1] and the remark after its proof, Y_m, X_m^{Law} and their common core are deformation retracts of one another, and their cohomology (with $\mathbb{Z}$ coefficients, as A is unimodular) is $\mathbb{Z}[x_1, \dots, x_N]/(\mathrm{Circ}(A) + M^*(A))$, where N is the number of columns of A and there is one variable for each column. By [HS, Theorem 8.3], $M^*(A)$ is generated by the monomials of the cuts of the underlying graph. In the presentation [HS, (37)], which eliminates $\mathrm{Circ}(A)$, each variable is replaced by the weight of its arrow, a linear form in the x_d; the monomial of the cut separating S from $(\Omega_m \setminus S) \cup \{\infty\}$ then becomes $\pm q_S$ (Definition 1.5). Hence

$$H^*(Y_m) \cong H^*(X_m^{\mathrm{Law}}) \cong P_m/J_m,$$

  and J_m is the image of the cocircuit ideal; compare [Hik, Section 5.1].
  The presentation of $H^*(Y_m)$ is due to Konno [Kon00, Theorem 3.1], for smooth toric hyperKähler manifolds and with $\mathbb{Z}$ coefficients, earlier than [HS]; see [HS, §1]. In his notation k is the Lie algebra of the acting torus, i.e. the row space of A. The monomials $\prod_{b_i \neq 0} u_i$, for nonzero $\sum_i b_i X_i \in k$, are the products over unions of cocircuits, and his linear forms, the elements of $\ker \iota^* \cap (\mathfrak{t}^N_{\mathbb{Z}})^*$, form $\mathrm{Circ}(A)$. So [Kon00, Theorem 3.1] is the presentation $\mathbb{Z}[x]/(\mathrm{Circ}(A) + M^*(A))$ above, and [Kon00, Theorem 3.2] gives the equivalent description in terms of the associated hyperplane arrangement. The equivariant cohomology for the residual torus, which is the same ring without the linear forms, is [Kon01, Theorem 2.4]. Konno's proofs [Kon00, §7], [Kon01, §3] proceed by induction on the number of coordinates, using Thom isomorphisms and split short exact sequences in equivariant cohomology; this is close in spirit to the proof of Proposition 2.1(a) in Section 2, which is organized instead by Hesselink's stratification.
  Combined with the vector bundle structure, the isomorphism $H^*(X_m^{\mathrm{Law}}) \cong P_m/J_m$ gives a second proof of Proposition 2.1(a), where the map $P_m \to H^*(U_m/T_m)$ sends x_d to $c_1(\mathcal{L}_d)$. By [HS, Corollary 8.4], the Euler characteristic of Y_m is the number of spanning trees of the underlying graph; for m_b in Example 1.13 it is 4. (If some component of the graph misses ∞, then $U_m = \emptyset$ and $J_m = P_m$, so all these spaces are empty.)


Apart from the alternative argument for the finite-dimensionality of P_m/J_m in §1.5, none of this is used in what follows.

(v) Applying Crawley-Boevey–Van den Bergh [CBVdB] to V_m^o with dimension vector $(1,\dots,1)$, and using (iv) and purity, the number $a_m(q)$ of absolutely indecomposable representations of V_m^o of this dimension over $\mathbb{F}_q$ is $a_m(q) = q^{-g_m}\#Y_m(\mathbb{F}_q) = q^{g_m}\operatorname{Hilb}(P_m/J_m; q^{-1})$. By Kac, a_m does not depend on o. This is also not used below.

Example 1.13. Let Q be the quiver $1 \to 2$ with a loop added at each vertex, so Q' is $1 \to 2$, $c_{12} = c_{21} = 1$ and $c_{11} = c_{22} = 0$ in (1.1). Take $v = (2,1)$ and $w = (1,0)$. Then $C(v) = 2$, and $\mathcal{M}(v)$ consists of

$$m_a:\ \lambda^1 = (2),\ \lambda^2 = (1), \qquad m_b:\ \lambda^1 = (1,1),\ \lambda^2 = (1).$$

For m_b there are slots d_1, d_2 of type $(1,1)$ and e of type $(2,1)$, with $b_{d_1e} = b_{d_2e} = 1$ and $b_{d_1d_2} = c_{11} = 0$; $W_{m_b} = \mathfrak{S}_2$ swaps d_1, d_2. The multipartition quiver is



so $\dim E_{m_b} = 6$ and U_{m_b}/T_{m_b} is a smooth toric threefold. Its underlying graph is a 4-cycle, $g_{m_b} = 1$, and Y_{m_b} is the A_3-surface. Writing $a = x_{d_1}$, $b = x_{d_2}$, $c = x_e$, the cut polynomials of Definition 1.5 are

$$a(a-c),\ b(b-c),\ (c-a)(c-b),\ a(c-b),\ b(c-a),\ ab,\ ab(a-c)(b-c),$$

for $S = \{d_1\},\{d_2\},\{e\},\{d_1,e\},\{d_2,e\},\{d_1,d_2,e\},\{d_1,d_2\}$ respectively. They generate $J_{m_b} = (a,b,c)^2$. Hence P_{m_b}/J_{m_b} has basis $1, a, b, c$, with Hilbert series $1+3q$, and $h_{m_b} = 1+2q$ (invariants 1, $a+b$, c).

For m_a there are slots d of type $(1,2)$ and e of type $(2,1)$, with $b_{de} = \min(2,1) = 1$; the multipartition quiver is $\infty \to d \rightleftarrows e$. Then $J_{m_a} = (x_d, x_e)$ and $h_{m_a} = 1$.

2. Proposition 6.2 via Kirwan's equivariant perfection

Recall the multipartition space E_m (Definition 1.4) and its open subset U_m (Definition 1.8), the torus T_m and $P_m = H^*_{T_m}(\mathrm{pt})$ (§1.5), the cut ideal J_m and the polynomial h_m (Definition 1.5), and the notation $(q)_m$, $w(k)$, $b(k,m)$ of Definition 1.2.

Proposition 2.1 ([Hik, Proposition 6.2 and (6.2)])**.** *For every multipartition m:*

(a) $H^*_{T_m}(U_m)$ is concentrated in even degrees, and restriction $P_m = H^*_{T_m}(E_m) \to H^*_{T_m}(U_m)$ is surjective with kernel J_m. In particular $H^*(U_m/T_m) \cong P_m/J_m$, and h_m is the Poincaré polynomial of $(U_m/T_m)/W_m$.

(b) In $\mathbb{Z}[[q]]$,

$$\frac{1}{(q)_m} = \sum_{k \le m} q^{\,w(k)+b(k,m-k)}\ \frac{h_{m-k}(q)}{(q)_k}. \tag{6.1}$$

The term $k = \emptyset$ in (6.1) is h_m, so (6.1) determines h_m recursively. The proof below is Kirwan's argument [Kir], applied to Hesselink's stratification of E_m for the torus T_m and the character $\rho = \det$ of (1.14), and made W_m-equivariant. For the toric hyperKähler manifold Y_m, the corresponding presentation of the cohomology was proved earlier by Konno [Kon00, Theorem 3.1], also by splitting exact sequences inductively; see Remark 1.12(iv).

2.1. **Recollection: Hesselink's stratification.** We recall [Hos, §§2.2–2.3] for a torus T acting linearly on a complex vector space V, with a character ρ and a norm $\|\cdot\|$ on 1-PSs as in §1.7. For $v \in V$ let C_v be the cone of allowable 1-PSs, defined as in (1.15), so that $\lambda \in C_v$ if and only if $\lim_{\tau\to 0} \lambda(\tau)\cdot v$ exists. For a ρ-unstable v, [Hos, Definition 2.12] puts

$$M^\rho(v) = \inf\Big\{\frac{(\rho,\lambda)}{\|\lambda\|} :\ 0 \neq \lambda \in C_v\Big\} < 0.$$

A 1-PS λ is *ρ-adapted* to v if it attains this infimum, and $\wedge^\rho(v)$ denotes the set of indivisible ρ-adapted 1-PSs; for a torus it has exactly one element [Hos, Lemma 2.13]. For a 1-PS λ the *Hesselink stratum* (Hoskins' blade, which equals the stratum $S_{[\lambda]}$ since T is abelian) and the *limit set* are [Hos, (5), (6)]

$$S_\lambda = \{v \ \rho\text{-unstable} :\ \lambda \in \wedge^\rho(v)\}, \qquad Z_\lambda = \{v \in S_\lambda :\ \lambda(\tau)\cdot v = v \text{ for all } \tau\},$$

and $S_0 := V^{\rho\text{-}ss}$ for the trivial 1-PS. Let V^λ be the fixed locus of λ, V^λ_+ the set of points v for which $\lim_{\tau\to 0}\lambda(\tau)\cdot v$ exists, and $p_\lambda : V^\lambda_+ \to V^\lambda$ the limit map. These are linear subspaces and a linear projection. By [Hos, Proposition 2.18, Lemma 2.19],

$$S_\lambda = p_\lambda^{-1}(Z_\lambda),$$

and Z_λ is the semistable locus for T acting on V^λ with respect to the character $\rho_\lambda = \|\lambda\|^2\rho - (\rho,\lambda)\lambda^*$, where λ^* is the character dual to λ under the inner product. In particular Z_λ is open in V^λ and S_λ is open in V^λ_+; the normal space of S_λ is $V^\lambda_- = \bigoplus_{(\chi,\lambda)<0} V_\chi$. By [Hos, Theorem 2.16, Remark 2.17], $V = \bigsqcup_\lambda S_\lambda$ is a decomposition into finitely many T-invariant locally closed subvarieties, *Hesselink's stratification*, and $\partial S_\lambda \cap S_{\lambda'} \neq \emptyset$ only if $\lambda < \lambda'$, where $\lambda < \lambda'$ means $(\rho,\lambda)/\|\lambda\| > (\rho,\lambda')/\|\lambda'\|$. By [Hos, Theorems 4.12 and 4.18] it coincides with Kirwan's stratification and with the Morse stratification of the norm square of the shifted moment map.

Kirwan's equivariant perfection [Kir]: if the Euler class of each V^λ_- is a non-zero-divisor in $H^*_T(S_\lambda)$ (the Atiyah–Bott criterion [AB]), then the Thom–Gysin sequences split, the restriction $H^*_T(V) \to H^*_T(V^{\rho\text{-}ss})$ is surjective, and

$$\operatorname{Hilb} H^*_T(V) = \operatorname{Hilb} H^*_T(V^{\rho\text{-}ss}) + \sum_{\lambda\neq 0} q^{\dim V^\lambda_-} \operatorname{Hilb} H^*_T(S_\lambda).$$

Below we verify everything we need directly for $V = E_m$.

2.2. **The adapted 1-PS of a point of E_m.**

Lemma 2.2. *Let $B \in E_m$. Recall the sets $R(B)$, $S(B)$ of reachable and unreachable slots and the 1-PS $\lambda_S = -1_S$ from Definition 1.8, and the cone C_B of allowable 1-PSs from (1.15). If $S(B) \neq \emptyset$, then B is ρ-unstable,*

$$M^\rho(B) = -\sqrt{|S(B)|}, \qquad \wedge^\rho(B) = \{\lambda_{S(B)}\}.$$

Proof. Put $S = S(B)$ and $R = R(B)$; B is ρ-unstable by Lemma 1.10(2). Let $0 \neq \lambda \in C_B$. By Lemma 1.9(1), $\lambda_d \geq 0$ for $d \in R$, so by Cauchy–Schwarz

$$(\rho,\lambda) = \sum_{d\in R}\lambda_d + \sum_{d\in S}\lambda_d \ \geq\ \sum_{d\in S}\lambda_d \ \geq\ -\sqrt{|S|}\Big(\sum_{d\in S}\lambda_d^2\Big)^{1/2} \ \geq\ -\sqrt{|S|}\,\|\lambda\|.$$

By Lemma 1.9(2), $\lambda_S \in C_B$ and $(\rho,\lambda_S)/\|\lambda_S\| = -|S|/\sqrt{|S|} = -\sqrt{|S|}$. Hence $M^\rho(B) = -\sqrt{|S|}$, attained by λ_S. Equality throughout forces $\lambda_d = 0$ for $d \in R$ and $\lambda|_S$ to be a negative multiple of 1_S, i.e. $\lambda \in \mathbb{R}_{>0}\lambda_S$. Since λ_S is indivisible, $\wedge^\rho(B) = \{\lambda_S\}$. □

Corollary 2.3. *Hesselink's stratification of E_m (§2.1) is*

$$E_m = \bigsqcup_{S\subseteq\Omega_m} S_{\lambda_S}, \qquad S_{\lambda_S} = \{B \in E_m : S(B) = S\},$$

with $S_{\lambda_\emptyset} = S_0 = U_m$; all other 1-PSs index empty strata. The ordering is $\lambda_S < \lambda_{S'}$ if and only if $|S| < |S'|$. For $0 \le r \le |l(m)|$ put

$$U_{m,r} = \{B \in E_m : \ |S(B)| \le r\} = \bigsqcup_{|S|\le r} S_{\lambda_S}.$$

Then $U_{m,0} = U_m$, $U_{m,|l(m)|} = E_m$, $U_{m,r} \setminus U_{m,r-1} = \bigsqcup_{|S|=r} S_{\lambda_S}$, and each $U_{m,r}$ is open in E_m.

Proof. The description of the strata follows from Lemma 2.2 and Lemma 1.10(2), and the ordering from $M^\rho = -\sqrt{|S|}$ on S_{λ_S}. The set $U_{m,r}$ is the union over $A \subseteq \Omega_m$ with $|A| = |l(m)|-r$ of the open sets $\{B : A \subseteq R(B)\}$. □

2.3. **The Hesselink strata of E_m.** Fix a nonempty $S \subseteq \Omega_m$, and put $R = \Omega_m \setminus S$, $k = \kappa(S)$ (the type of S, §1.3) and $m' = m - k$. Decompose

$$E_m = E_{RR} \oplus E_{SS} \oplus E_{SR} \oplus E_{RS} \oplus F_R \oplus F_S,$$

where E_{XY} collects the coordinates $B^{(p)}_{d\to e}$ with $d \in X$, $e \in Y$, and F_X the framing coordinates $a^{(p)}_d$ with $d \in X$. Via the identification (1.10), $R \cong \Omega_{m'}$, we identify $E_{RR} \oplus F_R = E_{m'}$ and $T_m = T_{m'} \times T_S$, where $T_S = (\mathbb{C}^\times)^S$. By (1.13), the λ_S-weights of the coordinates are -1 on $E_{RS} \oplus F_S$, $+1$ on E_{SR} and 0 otherwise. Hence, in the notation of §2.1,

$$E_m^{\lambda_S} = E_{RR} \oplus E_{SS} \oplus F_R, \qquad (E_m)^{\lambda_S}_+ = E_m^{\lambda_S} \oplus E_{SR}, \qquad N_S := (E_m)^{\lambda_S}_- = E_{RS} \oplus F_S, \tag{2.1}$$

and p_{λ_S} kills E_{SR}.

Lemma 2.4. *In this notation, recall $U_{m,r}$ from Corollary 2.3 and q_S from Definition 1.5.*

(a) The limit set is $Z_{\lambda_S} = U_{m'} \times E_{SS} \subseteq E_m^{\lambda_S}$, and the Hesselink stratum is $S_{\lambda_S} = p_{\lambda_S}^{-1}(Z_{\lambda_S}) = U_{m'} \times E_{SS} \times E_{SR}$.

(b) $S_{\lambda_S} = (E_m)^{\lambda_S}_+ \cap U_{m,|S|}$. In particular S_{λ_S} is smooth and closed in $U_{m,|S|}$.

(c) The normal bundle of S_{λ_S} in $U_{m,|S|}$ is the trivial bundle with fiber N_S. Its rank is $c_S = w(k) + b(k, m')$, and its equivariant Euler class is $e_S = q_S$.

(d) $H^*_{T_m}(S_{\lambda_S}) \cong H^*_{T_{m'}}(U_{m'}) \otimes \mathbb{C}[x_d : d \in S]$, compatibly with restriction from $P_m = P_{m'} \otimes \mathbb{C}[x_d : d \in S]$.

Proof. (a) By Corollary 2.3, $S_{\lambda_S} = \{B : S(B) = S\}$. If $S(B) = S$, then $\lambda_S \in C_B$ by Lemma 1.9(2), i.e. $B \in (E_m)^{\lambda_S}_+$. For $B \in (E_m)^{\lambda_S}_+$ the coordinates in $E_{RS} \oplus F_S$ vanish, so a path from ∞ never enters S; hence $S \subseteq S(B)$, and reachability of the slots of R only involves the coordinates in $E_{RR} \oplus F_R = E_{m'}$. Thus $S(B) = S$ if and only if $B|_{E_{m'}} \in U_{m'}$, with no condition on E_{SS} and E_{SR}. This gives S_{λ_S}, and $Z_{\lambda_S} = S_{\lambda_S} \cap E_m^{\lambda_S} = U_{m'} \times E_{SS}$.

(b) If $B \in (E_m)^{\lambda_S}_+ \cap U_{m,|S|}$, then $S \subseteq S(B)$ by the proof of (a) and $|S(B)| \le |S|$, so $S(B) = S$. The converse follows from (a). Since $(E_m)^{\lambda_S}_+$ is closed, S_{λ_S} is closed in $U_{m,|S|}$; by (a) it is open in the linear space $(E_m)^{\lambda_S}_+$, hence smooth.

(c) Since S_{λ_S} is open in $(E_m)^{\lambda_S}_+$, its normal bundle is $S_{\lambda_S} \times (E_m/(E_m)^{\lambda_S}_+) = S_{\lambda_S} \times N_S$. By (1.13), the weights of N_S are x_d, with multiplicity $w_{i(d)}$, for the framing coordinates at $d \in S$, and $x_d - x_e$, with multiplicity $b_{ed} = b_{de}$, for the coordinates $B_{e\to d}$ with $e \in R$, $d \in S$. So the rank is $w(k)+b(k,m')$ by (1.3) and (1.11), and the Euler class is $\prod_{d\in S} x_d^{w_{i(d)}} \prod_{d\in S, e\in R}(x_d - x_e)^{b_{de}} = q_S$.

(d) The linear contraction of $E_{SS} \times E_{SR}$ retracts S_{λ_S} equivariantly onto $U_{m'}$, on which T_S acts trivially; so $H^*_{T_{m'}\times T_S}(U_{m'}) = H^*_{T_{m'}}(U_{m'}) \otimes H^*(BT_S)$. □

Remark 2.5. This agrees with [Hos, Proposition 2.18]: $\lambda_S^* = -1_S$, so $\rho_{\lambda_S} = |S|\,1_{\Omega_m} - |S|\,1_S = |S|\,1_R$. A point of $E_m^{\lambda_S} = E_{m'} \oplus E_{SS}$ is semistable for this character exactly when its $E_{m'}$-component lies in $U_{m'}$, by Lemma 1.10(2) for m', since 1_R is trivial on T_S.

2.4. **The Atiyah–Bott criterion.**

Lemma 2.6. *Let $S \neq \emptyset$. Multiplication by the Euler class $e_S = q_S$ of Lemma 2.4(c) on $H^*_{T_m}(S_{\lambda_S})$ is injective.*

Proof. By Lemma 2.4(d), $H^*_{T_m}(S_{\lambda_S}) = A[x_d : d \in S]$ with $A = H^*_{T_{m'}}(U_{m'})$, an arbitrary commutative ring (possibly with nilpotents) containing the x_e, $e \in R$. Regard q_S (Definition 1.5) as a polynomial in the variables x_S over A. Expanding $(x_d - x_e)^{b_{de}}$ shows that its part of top total degree in x_S is the single monomial $\prod_{d\in S} x_d^{w_{i(d)}+\sum_{e\in R} b_{de}}$, with coefficient 1. If $f \neq 0$, the top-degree part of $q_S f$ is this monomial times the top-degree part of f, which is nonzero. □

Remark 2.7. This is the Atiyah–Bott criterion [AB]. The 1-PS λ_S^{-1} acts trivially on the retract $U_{m'} \times \{0\}$ of S_{λ_S} and with weight $+1$ on every coordinate of N_S. Restricting to it sends every factor of q_S to the generator $u \in H^2(B\mathbb{C}^\times)$, so $q_S \mapsto u^{c_S}$; this is exactly the leading monomial used above.

2.5. **Equivariant perfection.**

Proposition 2.8. *Recall $U_{m,r}$ from Corollary 2.3 and c_S from Lemma 2.4(c). For $1 \leq r \leq |l(m)|$ there is a short exact sequence*

$$0 \to \bigoplus_{|S|=r} H^{*-2c_S}_{T_m}(S_{\lambda_S}) \xrightarrow{\sum_S i_{S*}} H^*_{T_m}(U_{m,r}) \to H^*_{T_m}(U_{m,r-1}) \to 0,$$

where $i_S : S_{\lambda_S} \hookrightarrow U_{m,r}$ is the inclusion.

Proof. By Corollary 2.3 and Lemma 2.4(b), the union of the strata S_{λ_S} with $|S| = r$ is a closed smooth subvariety of $U_{m,r}$, with complement $U_{m,r-1}$. So there is the equivariant Thom–Gysin long exact sequence

$$\cdots \to \bigoplus_{|S|=r} H^{k-2c_S}_{T_m}(S_{\lambda_S}) \to H^k_{T_m}(U_{m,r}) \to H^k_{T_m}(U_{m,r-1}) \xrightarrow{\delta} \bigoplus_{|S|=r} H^{k+1-2c_S}_{T_m}(S_{\lambda_S}) \to H^{k+1}_{T_m}(U_{m,r}) \to \cdots.$$

By the self-intersection formula $i_S^* i_{S*} = e_S \cup (-) = q_S \cup (-)$, which is injective by Lemma 2.6. For $S' \neq S$ with $|S'| = |S|$ the strata are disjoint, so $i_{S'}^* i_{S*} = 0$. Hence $\sum_S i_{S*}$ is injective in every degree, the connecting maps δ vanish, and the long exact sequence splits into short exact sequences. □

In particular, restriction $H^*_{T_m}(E_m) \to H^*_{T_m}(U_m)$ is surjective: this is Kirwan surjectivity for the toric quotient U_m/T_m. Odd cohomology vanishes at every stage, because $\delta = 0$ and $H^{\mathrm{odd}}_{T_m}(E_m) = 0$.

2.6. **Proof of Proposition 2.1(a).** Recall the cut polynomials q_S and the cut ideal J_m from Definition 1.5. We use induction on the number of slots $|l(m)|$; for $m = \emptyset$ there is nothing to prove. Assume Proposition 2.1(a) for all multipartitions with fewer slots than m. For $0 \leq r \leq |l(m)|$ let $J_{m,r} \subseteq P_m$ be the ideal generated by the q_S with $|S| > r$, so that $J_{m,0} = J_m$. We show by descending induction on r that restriction $P_m \to H^*_{T_m}(U_{m,r})$ is surjective with kernel $J_{m,r}$. For $r = |l(m)|$ we have $U_{m,r} = E_m$ and $J_{m,r} = 0$.

Assume the claim for r. By Proposition 2.8, $H^*_{T_m}(U_{m,r-1})$ is the quotient of $H^*_{T_m}(U_{m,r}) = P_m/J_{m,r}$ by the images of the Gysin maps i_{S*}, $|S| = r$. Since $S \neq \emptyset$, $m' = m - \kappa(S)$ has fewer slots, so by induction and Lemma 2.4(d), $P_m = P_{m'} \otimes \mathbb{C}[x_S] \to H^*_{T_m}(S_{\lambda_S})$ is surjective. Thus every class on S_{λ_S} is $i_S^* f$ for some $f \in P_m$, and by the projection formula $i_{S*}(i_S^* f) = f \cup i_{S*}(1)$. Moreover $i_{S*}(1) = [S_{\lambda_S}]$ is the restriction to $U_{m,r}$ of the equivariant class of the linear subspace $(E_m)_+^{\lambda_S} \subseteq E_m$, which is the Euler class $e(N_S) = q_S$; here we use $S_{\lambda_S} = (E_m)_+^{\lambda_S} \cap U_{m,r}$ (Lemma 2.4(b)). Hence the image of the Gysin maps is the ideal generated by the q_S with $|S| = r$, and $H^*_{T_m}(U_{m,r-1}) = P_m/(J_{m,r} + (q_S : |S| = r)) = P_m/J_{m,r-1}$. For $r = 0$ this gives $H^*_{T_m}(U_m) = P_m/J_m$. Evenness was shown after Proposition 2.8, and the statements about U_m/T_m follow from Lemma 1.10(3). □

2.7. **Proof of Proposition 2.1(b): taking W_m-invariants.** Recall the action of W_m on E_m and on equivariant cohomology from §1.6, and the types of subsets and their stabilizers from §1.3. Since $S(\sigma B) = \sigma S(B)$, W_m preserves each $U_{m,r}$ and permutes the strata: $\sigma(S_{\lambda_S}) = S_{\lambda_{\sigma S}}$. Restriction maps commute with W_m. The Gysin maps do as well, because σ induces complex-linear isomorphisms $N_S \to N_{\sigma S}$ (permutations of coordinates), so it preserves the Thom classes defined by the complex orientations, without signs. Hence the sequences of Proposition 2.8 are sequences of W_m-modules, and they stay exact after taking invariants (§1.6(a)).

Fix $r \geq 1$. For each $k \leq m$ with $|l(k)| = r$ choose $S_k \subseteq \Omega_m$ of type k. The subsets of type k form the orbit $W_m \cdot S_k$, with stabilizer $W_k \times W_{m-k}$, and $c_S = w(k) + b(k, m-k)$ is constant on it. For a W_m-module $\bigoplus_{S \in W_m S_k} M_S$ permuted transitively, projection to the S_k-summand identifies the invariants with $M_{S_k}^{W_k \times W_{m-k}}$; in particular no binomial coefficients appear. By Lemma 2.4(d) and part (a) for $m' = m - k$, compatibly with the actions of $W_{m'}$ and W_k,

$$H^*_{T_m}(S_{\lambda_{S_k}})^{W_k \times W_{m'}} \cong (P_{m'}/J_{m'})^{W_{m'}} \otimes \mathbb{C}[x_d : d \in S_k]^{W_k},$$

whose Hilbert series is $h_{m'}(q)/(q)_k$ by Definition 1.5 and §1.6(c). Therefore

$$\mathrm{Hilb}\left(H^*_{T_m}(U_{m,r})^{W_m}\right) - \mathrm{Hilb}\left(H^*_{T_m}(U_{m,r-1})^{W_m}\right) = \sum_{\substack{k \leq m \\ |l(k)| = r}} q^{\,w(k) + b(k,m-k)} \frac{h_{m-k}(q)}{(q)_k}.$$

Summing over $r = 1, \ldots, |l(m)|$ telescopes. At $r = |l(m)|$ we have $U_{m,r} = E_m$ and $\mathrm{Hilb}(P_m^{W_m}) = 1/(q)_m$ by §1.6(c); at $r = 0$ we have $\mathrm{Hilb}(H^*_{T_m}(U_m)^{W_m}) = h_m$ by part (a). This gives (6.1). □

2.8. **Dictionary and an example.** The following table summarizes the correspondence with [Hos, §2] and [Kir].

Hoskins [Hos, §2]	multipartition space E_m, character $\rho = \det$
ρ-semistable locus $V^{\rho\text{-}ss} = S_0$	U_m (Lemma 1.10(2))
cone of allowable 1-PSs C_v	C_B; $\lambda_d \geq 0$ on $R(B)$ (Lemma 1.9)
$M^\rho(v)$, $\wedge^\rho(v)$	$-\sqrt{\lvert S(B)\rvert}$, $\{\lambda_{S(B)}\}$ (Lemma 2.2)
ordering $\lambda < \lambda'$	$\lvert S\rvert < \lvert S'\rvert$ (Corollary 2.3)
V^λ, V^λ_+	$E_{RR} \oplus E_{SS} \oplus F_R$, $E_m^{\lambda_S} \oplus E_{SR}$ (2.1)
limit set Z_λ	$U_{m'} \times E_{SS}$ (Lemma 2.4(a))
Hesselink stratum $S_\lambda = p_\lambda^{-1}(Z_\lambda)$	$\{S(B) = S\} = U_{m'} \times E_{SS} \times E_{SR}$
normal space V^λ_-, codimension	$N_S = E_{RS} \oplus F_S$, $w(k) + b(k, m')$
Euler class of V^λ_-	cut polynomial q_S (Lemma 2.4(c))
Atiyah–Bott criterion	leading monomial of q_S (Lemma 2.6)
Kirwan's formula	(6.1), in W_m-invariants

Example 2.9. For m_b in Example 1.13, $(q)_{m_b} = (1-q)^2(1-q^2)$, and the six $k \leq m_b$, written as $(k_{1,1}, k_{2,1})$, contribute to (6.1):

k	$w(k) + b(k, m_b - k)$	$h_{m_b - k}$	term
$(0,0)$	0	$1+2q$	$1+2q$
$(1,0)$	$1+1$	1	$q^2/(1-q)$
$(2,0)$	$2+2$	0	0 (slot e alone has no framing, $U = \emptyset$)
$(0,1)$	$0+2$	1	$q^2/(1-q)$
$(1,1)$	$1+1$	1	$q^2/(1-q)^2$
$(2,1)$	$2+0$	1	$q^2/\big((1-q)^2(1-q^2)\big)$

Multiplying by $(1-q)^2(1-q^2)$, the right side of (6.1) becomes $(1+2q)(1-2q+2q^3-q^4) + 2q^2(1-q)(1-q^2) + q^2(1-q^2) + q^2 = 1$, as it should.

3. Hausel's formula and Corollary 6.3

3.1. **Hausel's formula.** In this subsection Q is an arbitrary finite quiver; loops and multiple arrows are allowed. For dimension vectors v, w let $G = \prod_i GL(V_i)$, $N = \bigoplus_{a \in Q_1} \mathrm{Hom}(V_{t(a)}, V_{h(a)}) \oplus \bigoplus_i \mathrm{Hom}(V_i, W_i)$, let $\mu : T^*N \to \mathfrak{g}^\vee \cong \mathfrak{g}$ be the moment map, and put

$$\mathfrak{M}_\eta(v,w) = \mu^{-1}\big((\mathrm{id}_{V_i})_i\big)/\!/G.$$

When nonempty, this is smooth of dimension $2\delta(v,w)$, where

$$\delta(v,w) = \sum_{a \in Q_1} v_{t(a)} v_{h(a)} - \sum_i v_i^2 + w \cdot v.$$

Its odd cohomology vanishes; write $P_{\mathfrak{M}_\eta(v,w)}(q)$ for its Poincaré polynomial in the half-degree variable q. For partitions μ, ν let $\langle \mu, \nu \rangle = \sum_{a,b} \min(\mu_a, \nu_b)$, let (1^{w_i}) be the partition with w_i parts equal to 1, and for a Q_0-tuple of partitions λ put $|\lambda| = (|\lambda^i|)_i$ and

$$E_Q(\lambda) = \sum_{a \in Q_1} \langle \lambda^{t(a)}, \lambda^{h(a)} \rangle - \sum_i \langle \lambda^i, \lambda^i \rangle.$$

Theorem 3.1 (Hausel [Hau]; [Hik, (6.9)]). *In* $\mathbb{C}(q)[[z_i : i \in Q_0]]$,

$$\sum_v q^{-\delta(v,w)} P_{\mathfrak{M}_\eta(v,w)}(q)\, z^v = \frac{\displaystyle\sum_\lambda z^{|\lambda|} \frac{q^{-E_Q(\lambda)-\sum_i \langle \lambda^i, (1^{w_i})\rangle}}{(q)_{m(\lambda)}}}{\displaystyle\sum_\lambda z^{|\lambda|} \frac{q^{-E_Q(\lambda)}}{(q)_{m(\lambda)}}}, \tag{3.1}$$

where λ *runs over all* Q_0*-tuples of partitions, including the empty one, and* $(q)_{m(\lambda)}$ *is as in* (1.4).

We do not discuss Hausel's proof, which uses an arithmetic Fourier transform. Remark 3.2 explains the alternative point-count proof of [Nak04], and Remark 3.3 relates $\mathfrak{M}_\eta(v,w)$ to the quiver variety at determinant stability.

Remark 3.2 (The point-count proof, after [Nak04]). We explain the right side of (3.1) following the unpublished note [Nak04, §1], which obtains (3.1) from the Kac–Stanley formula [Kac, Hua], Crawley-Boevey's trick [CB] and a theorem of Crawley-Boevey and Van den Bergh [CBVdB]. Work over a finite field $\mathbb{F}_u$ and put

$$R_Q(V) = \bigoplus_{a\in Q_1} \mathrm{Hom}(V_{t(a)}, V_{h(a)}), \qquad R_Q(V,W) = R_Q(V) \oplus \bigoplus_i \mathrm{Hom}(W_i, V_i),$$

with $G_V = \prod_i GL(V_i)$ acting trivially on W.

(a) The series P_Q *and* P'_Q. For a partition λ let $b_\lambda(t) = \prod_{s\ge 1} \prod_{j=1}^{m_s(\lambda)} (1-t^j)$ [Mac, III, (2.12)], where $m_s(\lambda)$ is the number of parts equal to s. Note $\langle \lambda, (1)\rangle = l(\lambda)$, the number of parts. With $X^{|\lambda|} = \prod_i X_i^{|\lambda^i|}$, define

$$P_Q(X,u) = \sum_\lambda \frac{u^{\sum_{a\in Q_1} \langle \lambda^{t(a)}, \lambda^{h(a)}\rangle}}{\prod_i u^{\langle \lambda^i, \lambda^i\rangle}\, b_{\lambda^i}(u^{-1})} X^{|\lambda|},$$

$$P'_Q(X,u) = \sum_\lambda \frac{u^{\sum_{a\in Q_1} \langle \lambda^{t(a)}, \lambda^{h(a)}\rangle + \sum_i w_i \langle \lambda^i, (1)\rangle}}{\prod_i u^{\langle \lambda^i, \lambda^i\rangle}\, b_{\lambda^i}(u^{-1})} X^{|\lambda|}.$$

Since $\prod_i b_{\lambda^i}(u^{-1}) = (u^{-1})_{m(\lambda)}$ and $w_i\langle \lambda^i, (1)\rangle = \langle \lambda^i, (1^{w_i})\rangle$, the substitution $u = q^{-1}$, $X = z$ turns P_Q and P'_Q into the denominator and the numerator of the right side of (3.1).

(b) Burnside's lemma and the Kac–Stanley formula. For a finite G_V-set R, Burnside's lemma gives $|R/G_V| = |G_V|^{-1} \sum_{g\in G_V} |R^g|$. For $g \in G_V$, decompose each V_i into the generalized eigenspaces $V_i^{(f)}$ of g_i, indexed by monic irreducible polynomials $f \ne t$ over $\mathbb{F}_u$. A linear map $\phi : V_i \to V_j$ with $g_j\phi = \phi g_i$ maps $V_i^{(f)}$ into $V_j^{(f)}$. Hence

$$R_Q(V)^g = \bigoplus_f R_Q(V^{(f)})^{g_f}, \qquad Z_{G_V}(g) = \prod_f Z(g_f),$$

so the Burnside sum factorizes over f. Consider the factor $f = t-1$, where the component of g is unipotent. If $g_i - 1$ is nilpotent of Jordan type λ^i, then $R_Q(V)^g$ consists of the $\mathbb{F}_u[t]$-linear maps between the modules V_i, with t acting by $g_i - 1$. Therefore

$$|R_Q(V)^g| = u^{\sum_a \langle \lambda^{t(a)}, \lambda^{h(a)}\rangle}, \qquad |Z_{G_V}(g)| = \prod_i u^{\langle \lambda^i, \lambda^i\rangle} b_{\lambda^i}(u^{-1}),$$

and the unipotent elements of type λ contribute $|R^g|/|Z_{G_V}(g)|$, the λ-term of P_Q. The same computation over $\mathbb{F}_{u^d}$ gives the local factor $P_Q(X^d, u^d)$ for f of degree d, and one obtains the

Kac–Stanley formula [Hua, Theorem 4.3]:

$$\sum_V |R_Q(V)/G_V|\, X^{\dim V} = \prod_{d\geq 1} P_Q(X^d, u^d)^{\varphi_d(u)},$$

where $\varphi_1(u) = u-1$ and, for $d \geq 2$, $\varphi_d(u)$ is the number of monic irreducible polynomials of degree d over $\mathbb{F}_u$.

(c) Why only eigenvalue 1 *survives.* Since G_V acts trivially on W, an element g fixes a framing map $a_i \in \mathrm{Hom}(W_i, V_i)$ if and only if $g_i a_i = a_i$, i.e. $\mathrm{im}\, a_i \subseteq \ker(g_i - 1)$. Thus

$$|\mathrm{Hom}(W_i, V_i)^g| = u^{\,w_i \dim \ker(g_i - 1)},$$

which depends only on the component of g at the eigenvalue 1. For g_i unipotent of type λ^i, $\dim\ker(g_i-1) = l(\lambda^i) = \langle \lambda^i, (1)\rangle$. Adding the framing therefore changes only the local factor at $f = t-1$, from P_Q to P'_Q:

$$\sum_V |R_Q(V,W)/G_V|\, X^{\dim V} = P'_Q(X,u)\, P_Q(X,u)^{u-2} \prod_{d\geq 2} P_Q(X^d, u^d)^{\varphi_d(u)}.$$

Dividing by the Kac–Stanley formula, all local factors at eigenvalues different from 1 cancel:

$$\frac{\sum_V |R_Q(V,W)/G_V|\, X^{\dim V}}{\sum_V |R_Q(V)/G_V|\, X^{\dim V}} = \frac{P'_Q(X,u)}{P_Q(X,u)}. \tag{3.2}$$

In [Nak04] this is derived from the Kac–Stanley formula for the quiver $Q^{\#}$ of Crawley-Boevey's trick, by taking the coefficients of Y^0 and Y^1, where Y records the dimension at the extra vertex ∞. There an element $(g,c) \in G_V \times \mathbb{F}_u^\times$ fixes the framing arrows $\infty \to i$ exactly when they land in the c-eigenspace of g_i. Since the diagonal scalars act trivially, (g,c) and $(c^{-1}g, 1)$ have the same fixed points, so the $u-1$ values of c contribute equally; this factor $u-1$ cancels against $|GL_1(\mathbb{F}_u)| = u^{\langle (1),(1)\rangle} b_{(1)}(u^{-1})$. Normalizing $c=1$ gives the description above.

(d) Krull–Schmidt and Crawley-Boevey–Van den Bergh. A point $(B,a) \in R_Q(V,W)$ is a representation of $Q^{\#}$ with dimension 1 at ∞ [CB]. It decomposes uniquely as $(B_1,a_1) \oplus B_2$, where (B_1,a_1) is indecomposable, hence absolutely indecomposable since the dimension at ∞ is 1, and B_2 is a representation of Q. Hence

$$\sum_V |R_Q(V,W)/G_V|\, X^{\dim V} = \sum_V |R_Q(V)/G_V|\, X^{\dim V} \times \sum_V |R_Q(V,W)^{\mathrm{a.i.}}/G_V|\, X^{\dim V},$$

that is, framed = (absolutely indecomposable framed) × unframed. By [CBVdB, Proposition 2.2.1], in the form [Nak04, Theorem 1.1], for $W \neq 0$ and a generic parameter $\zeta \in \mathbb{F}_u^{Q_0}$ (for instance $\zeta = \eta$ when the characteristic exceeds $\sum_i v_i$),

$$|R_Q(V,W)^{\mathrm{a.i.}}/G_V| = u^{-\delta(v,w)}\, \#\mathfrak{M}_\zeta(v,w)(\mathbb{F}_u).$$

Comparing with (3.2) gives $\sum_v u^{-\delta(v,w)} \#\mathfrak{M}_\zeta(v,w)(\mathbb{F}_u)\, X^v = P'_Q(X,u)/P_Q(X,u)$ [Nak04, Theorem 1.2], which is Hausel's formula. Finally $\#\mathfrak{M}_\zeta(v,w)(\mathbb{F}_u) = u^{2\delta(v,w)} P_{\mathfrak{M}_\eta(v,w)}(u^{-1})$, since the count is the compactly supported Poincaré polynomial of the complex quiver variety (see [Nak04, §2]), and the substitution $u = q^{-1}$ gives (3.1).

(e) Loops. If Q has a loop at i, its fixed points $\mathrm{End}(V_i)^g$ form the centralizer algebra of g_i. For g_i unipotent of type λ^i it has $u^{\langle \lambda^i, \lambda^i\rangle}$ elements, which cancels the factor $u^{\langle \lambda^i, \lambda^i\rangle}$ in $|Z_{G_V}(g)|$. This is the cancellation of Lemma 3.4 below, seen from the Burnside side.

Remark 3.3 (Determinant stability). Let $X(v,w)$ be the quiver variety at determinant stability, i.e. the GIT quotient of $\mu^{-1}(0)$ by G for the character det. For $\zeta = (\zeta_{\mathbb{R}}, \zeta_{\mathbb{C}}) \in \mathbb{R}^{Q_0} \oplus \mathbb{C}^{Q_0}$ let $\mathfrak{M}_\zeta(v,w)$ be the hyperKähler quotient of [Nak94, §2], with the convention $\mu = -\zeta$ used there. Then:

- by [Nak94, Theorem 3.2 and Proposition 3.5], $X(v,w)$ is homeomorphic to $\mathfrak{M}_{(\zeta_{\mathbb{R}},0)}(v,w)$ for a suitable nonzero real multiple $\zeta_{\mathbb{R}}$ of $(1,\dots,1)$. ([Nak94, Proposition 3.5] treats $\zeta_{\mathbb{R}}$ with positive entries, where stability is a condition on the maps $V\to W$; changing the sign of $\zeta_{\mathbb{R}}$ gives the dual condition, cyclicity with respect to the maps $W\to V$, which is the determinant stability used here.)
- by [Nak94, Theorem 3.1], $\mathfrak{M}_{\eta}(v,w)$ is homeomorphic to $\mathfrak{M}_{(0,-\eta)}(v,w)$, where $\eta=(1,\dots,1)$.

Both parameters lie off every wall $\mathbb{R}^3\otimes D_\theta$ with $0\neq\theta\in\mathbb{Z}^{Q_0}_{\geq 0}$, because $\sum_k\theta_k\neq 0$; so they are generic. By [Nak94, Corollary 4.2], $\mathfrak{M}_{(\zeta_{\mathbb{R}},0)}(v,w)$ and $\mathfrak{M}_{(0,-\eta)}(v,w)$ are diffeomorphic; in particular one is empty if and only if the other is. Hence $P_{X(v,w)}=P_{\mathfrak{M}_\eta(v,w)}$. This is the argument of [Hik, proof of Proposition 6.4].

The paper [Nak94] assumes that no edge joins a vertex with itself, whereas here Q may have loops, and from §3.2 on it has a loop at every vertex. The proof of [Nak94, Corollary 4.2] does not use this assumption: it relies only on [Nak94, Theorem 4.1(1),(2)] and on rotating the complex structures I,J,K.

3.2. **Quivers with a loop at every vertex.** From now on Q again has a loop at every vertex. Recall Q' and c_{ij} from (1.1), $C(v)$ from (1.2), and $l(m)$, $w(m)$, $(q)_m$, D_m from Definition 1.2.

Lemma 3.4 (Loop cancellation)**.** *For every Q_0-tuple of partitions λ, with $m=m(\lambda)$,*

$$E_Q(\lambda)=D_m,\qquad \sum_i\langle\lambda^i,(1^{w_i})\rangle=w(m),\qquad \delta(v,w)=C(v)+w\cdot v.$$

Proof. The chosen loops contribute $\sum_i\langle\lambda^i,\lambda^i\rangle$ to $\sum_{a\in Q_1}\langle\lambda^{t(a)},\lambda^{h(a)}\rangle$, which cancels the vertex term; the remaining arrows give D_m by (1.6). Next, $\langle\lambda^i,(1^{w_i})\rangle=\sum_a\sum_{b=1}^{w_i}\min(\lambda^i_a,1)=w_i\,l_i(m)$. The same cancellation applied to $\sum_a v_{t(a)}v_{h(a)}-\sum_i v_i^2$ gives $C(v)$. □

Using $\lambda\leftrightarrow m(\lambda)$, and with h_m as in Definition 1.5, define the following series in $\mathbb{C}(q)[[z]]$:

$$\mathcal{A}^{\mathrm{fr}}(z)=\sum_{m\in\mathcal{M}}z^{\|m\|}\frac{q^{-D_m-w(m)}}{(q)_m},\qquad \mathcal{A}^0(z)=\sum_{m\in\mathcal{M}}z^{\|m\|}\frac{q^{-D_m}}{(q)_m},\tag{3.3}$$
$$\mathcal{H}(z)=\sum_{m\in\mathcal{M}}z^{\|m\|}q^{-D_m-w(m)}h_m(q).$$

Each z^v-coefficient is a finite sum over $\mathcal{M}(v)$, and $\mathcal{A}^0$ has constant term 1, so $\mathcal{A}^0$ is invertible. By Lemma 3.4 and Remark 3.3, Theorem 3.1 becomes

$$\sum_v q^{-C(v)-w\cdot v}\,P_{X(v,w)}(q)\,z^v=\frac{\mathcal{A}^{\mathrm{fr}}(z)}{\mathcal{A}^0(z)}.\tag{3.4}$$

3.3. **Computing the ratio by Proposition 6.2.**

Corollary 3.5 ([Hik, Corollary 6.3])**.** *$\mathcal{A}^{\mathrm{fr}}(z)=\mathcal{A}^0(z)\cdot\mathcal{H}(z)$; equivalently $\mathcal{H}=\mathcal{A}^{\mathrm{fr}}/\mathcal{A}^0$.*

Proof. Recall the identity (6.1) of Proposition 2.1(b) and the series (3.3). Multiply (6.1) by $z^{\|m\|}q^{-D_m-w(m)}$. The left side becomes the m-th term of $\mathcal{A}^{\mathrm{fr}}$. For $k\leq m$ with $m'=m-k$, (1.7) gives

$$-D_m-w(m)+w(k)+b(k,m')=-D_k-\big(D_{m'}+w(m')\big),$$

and $z^{\|m\|}=z^{\|k\|}z^{\|m'\|}$. So the k-th term on the right factors as

$$\Big(z^{\|k\|}\frac{q^{-D_k}}{(q)_k}\Big)\cdot\Big(z^{\|m'\|}q^{-D_{m'}-w(m')}h_{m'}(q)\Big),$$

a term of $\mathcal{A}^0$ times a term of $\mathcal{H}$. Since $(m,k)\mapsto(k,m-k)$ is a bijection from $\{(m,k):k\le m\}$ onto $\mathcal{M}\times\mathcal{M}$, summing over all m gives $\mathcal{A}^{\mathrm{fr}}=\mathcal{A}^0\cdot\mathcal{H}$. □

Comparing the z^v-coefficients of (3.4) and Corollary 3.5, and multiplying by $q^{C(v)+w\cdot v}$, gives the formula of [Hik, (6.8)]:

Corollary 3.6 ([Hik, Proposition 6.4])**.** *Let $X(v,w)$ be the quiver variety at determinant stability (Remark 3.3). Then*

$$P_{X(v,w)}(q)=\sum_{m\in\mathcal{M}(v)} q^{\,C(v)-D_m+w\cdot(v-l(m))}\,h_m(q).$$

3.4. **Examples.**

Example 3.7 (Jordan quiver)**.** Let Q be the Jordan quiver, so Q' has no arrows and $c=0$. Multipartitions are partitions λ, with m_s parts of size s. The multipartition quiver Γ_m has no arrows between slots, and every slot receives w framing arrows. Hence $U_m=\prod_d(\mathbb{C}^w\setminus 0)$, $U_m/T_m=(\mathbb{P}^{w-1})^{l(\lambda)}$, $q_S=\prod_{d\in S}x_d^w$, $J_m=(x_d^w:d\in\Omega_m)$, and

$$h_m(q)=\prod_s\begin{bmatrix}m_s+w-1\\ m_s\end{bmatrix}_q,$$

the Poincaré polynomial of $\prod_s\mathrm{Sym}^{m_s}\mathbb{P}^{w-1}$. The Hesselink strata S_{λ_S} (Corollary 2.3) are $\{a_d=0\ (d\in S),\ a_d\neq 0\ (d\notin S)\}$, of codimension $w|S|$. Here $D_m=0$, and every sum factors over part sizes. Euler's identity $\sum_{k\ge0}t^k/(q)_k=\prod_{j\ge0}(1-tq^j)^{-1}$ gives

$$\mathcal{A}^{\mathrm{fr}}=\prod_{s\ge1}\prod_{j\ge0}\frac{1}{1-z^sq^{j-w}},\qquad \mathcal{A}^0=\prod_{s\ge1}\prod_{j\ge0}\frac{1}{1-z^sq^j},\qquad \mathcal{H}=\frac{\mathcal{A}^{\mathrm{fr}}}{\mathcal{A}^0}=\prod_{s\ge1}\prod_{k=1}^{w}\frac{1}{1-z^sq^{-k}}.$$

Corollary 3.6 gives the Betti numbers of the Gieseker space $X(n,w)$:

$$P_{X(n,w)}(q)=\sum_{\lambda\vdash n}q^{\,w(n-l(\lambda))}\prod_s\begin{bmatrix}m_s+w-1\\ m_s\end{bmatrix}_q.$$

For $w=1$ this is $\sum_{\lambda\vdash n}q^{\,n-l(\lambda)}$ for $\mathrm{Hilb}^n(\mathbb{C}^2)$; for $w=n=2$ it is $(1+q+q^2)+q^2(1+q)=1+q+2q^2+q^3$.

Example 3.8 (The A_2 example continued)**.** In Example 1.13, $C(v)=2$. For m_b: $D_{m_b}=2$, $l(m_b)=(2,1)$, $w\cdot(v-l)=0$, so the term is $q^0(1+2q)$. For m_a: $D_{m_a}=1$, $l(m_a)=(1,1)$, $w\cdot(v-l)=1$, so the term is $q^2\cdot 1$. Corollary 3.6 gives

$$P_{X((2,1),(1,0))}(q)=(1+2q)+q^2=1+2q+q^2,$$

for an 8-dimensional variety. Note that the unlooped A_2 variety with these data is empty; the formula concerns the quiver with loops.

References

[AB] M. F. Atiyah and R. Bott, *The Yang–Mills equations over Riemann surfaces*, Philos. Trans. Roy. Soc. London Ser. A **308** (1983), 523–615.

[CB] W. Crawley-Boevey, *Geometry of the moment map for representations of quivers*, Compos. Math. **126** (2001), 257–293.

[CBVdB] W. Crawley-Boevey and M. Van den Bergh, *Absolutely indecomposable representations and Kac–Moody Lie algebras*, Invent. Math. **155** (2004), 537–559.

[Hau] T. Hausel, *Betti numbers of holomorphic symplectic quotients via arithmetic Fourier transform*, Proc. Natl. Acad. Sci. USA **103** (2006), 6120–6124.

[Hes] W. H. Hesselink, *Desingularizations of varieties of nullforms*, Invent. Math. **55** (1979), 141–163.

[Hik] OpenAI, *The equivariant cohomological Hikita conjecture for arbitrary quivers*, preprint dated September 24, 2026.

[Hos] V. Hoskins, *Stratifications associated to reductive group actions on affine spaces*, Q. J. Math. **65** (2014), 1011–1047; doi:10.1093/qmath/hat046.

[HS] T. Hausel and B. Sturmfels, *Toric hyperKähler varieties*, Doc. Math. **7** (2002), 495–534; arXiv:math/0203096.

[Hua] J. Hua, *Counting representations of quivers over finite fields*, J. Algebra **226** (2000), 1011–1033.

[Kac] V. G. Kac, *Root systems, representations of quivers and invariant theory*, in: Invariant Theory, Lecture Notes in Math. **996**, Springer, 1983, 74–108.

[Kem] G. R. Kempf, *Instability in invariant theory*, Ann. of Math. (2) **108** (1978), 299–316.

[Kin] A. D. King, *Moduli of representations of finite dimensional algebras*, Quart. J. Math. Oxford Ser. (2) **45** (1994), 515–530.

[Kir] F. C. Kirwan, *Cohomology of Quotients in Symplectic and Algebraic Geometry*, Mathematical Notes **31**, Princeton University Press, 1984.

[Kon00] H. Konno, *Cohomology rings of toric hyperKähler manifolds*, Internat. J. Math. **11** (2000), no. 8, 1001–1026.

[Kon01] H. Konno, *Equivariant cohomology rings of toric hyperKähler manifolds*, in: Proceedings of the Second Meeting on Quaternionic Structures in Mathematics and Physics (Rome, 6–10 September 1999), pp. 231–240, 2001.

[Mac] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, 2nd ed., Oxford University Press, 1995.

[Nak94] H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac–Moody algebras*, Duke Math. J. **76** (1994), 365–416.

[Nak04] H. Nakajima, unpublished note, 2004.

[Nes] L. Ness, *A stratification of the null cone via the moment map*, Amer. J. Math. **106** (1984), 1281–1329.