

Biotransformations from and to methylated flavonoids

Subtitle

Benjamin Weigel
Leibniz-Institute of Plant Biochemistry
Department of Bioorganic Chemistry
Weinberg 3
06120 Halle(Saale)
May 6, 2015

Advisor: Prof. Dr. Ludger A. Wessjohann
wessjohann@ipb-halle.de
+49 (345) 5582-1301

noch nicht bekannt

1 **Contents**

2	Notes of Revisors	viii
3	I Preface	1
4	<hr/>	
5	1 Abstracts	2
6	1.1 English Abstract	2
7	1.2 Deutsche Zusammenfassung	2
8	II Thesis	3
9	<hr/>	
10	2 Introduction	4
11	2.1 Natural products and secondary metabolites	4
12	2.1.1 General.	4
13	2.1.2 Classes of natural products	4
14	2.2 Alkylating reactions in nature	4
15	2.2.1 Methylation	4
16	2.2.2 Prenylation	4
17	2.2.3 Glycosylation	5
18	2.3 Usage and expansion of natures reaction toolbox	5
19	2.3.1 Terpene synthases and elongases	5
20	2.3.2 Methyl transferases.	5
21	2.3.3 Glycosyl transferases.	5
22	2.3.4 Other important enzymes in biotech research.	5
23	2.4 Conclusion	5

1	3 Material And Methods	6
2	3.1 Materials	6
3	3.1.1 Chemicals	6
4	3.1.2 Commonly used solutions and buffers	6
5	3.1.3 Culture media used to grow bacteria	7
6	3.1.4 Bacterial strains	7
7	3.1.5 Plasmids	8
8	3.1.6 Oligonucleotides and synthetic genes	9
9	3.1.7 Instruments	9
10	3.1.8 Software	10
11	3.2 Microbiology	10
12	3.3 Molecular Biology	10
13	3.3.1 Golden Gate Cloning	11
14	3.3.2 Subcloning of genes	11
15	3.4 Protein biochemistry	11
16	3.4.1 Protein production test (expression test)	11
17	3.4.2 Protein subfractionation	12
18	3.4.3 Protein sample concentration by TCA precipitation	12
19	3.4.4 Preparation of periplasmic protein	13
20	3.4.5 Discontinuous SDS-polyacrylamide gel electrophoresis (SDS-	
21	PAGE)	13
22	3.4.6 Buffer change of protein samples	14
23	3.4.7 Production of recombinant protein	14
24	3.4.8 Preparation of IB	15
25	3.4.9 Purification of His-tagged proteins using immobilized metal	
26	affinity chromatography (IMAC)	15
27	3.5 Analytics	16
28	3.5.1 In vitro determination of glucose	16
29	3.5.2 HPLC measurements	18
30	4 Evaluation of PFOMT towards the acceptance of long-chain SAM	
31	analogues	19
32	4.1 Introduction	19
33	4.2 Substrate binding studies using ITC.	19
34	4.3 Determination of the structure of <i>apo</i> -PFOMT.	19
35	4.3.1 PFOMT activity in deep eutectic solvent (DES) / Solubility-	
36	enhancing effects of DES.	19
37	4.4 Study of variants for long-chain alkylations	19
38	4.4.1 PFOMT-Paper (DIM)	19
39	4.4.2 Dockings???.	19

1	4.5 Colclusion/Discussion	19
2	5 Enzymatic methylation of Non-catechols	20
3	5.1 Introduction	20
4	5.2 SOMT-2.	20
5	5.2.1 In vivo methylation studies using <i>N. benthamiana</i>	20
6	5.2.2 In vivo studies in <i>E. coli</i>	20
7	5.2.3 In vitro studies using recombinantly produced SOMT-2	20
8	5.3 PFOMT	20
9	5.3.1 Acidity and Nucleophilicity of phenolic hydroxyl-groups	20
10	5.3.2 pH-Profiles of PFOMT-catalysis	20
11	5.3.3 Influence of Mg^{2+} on PFOMT activity	20
12	5.4 Consensus or Bioinformatic points-of-view (COMT)???	20
13	5.5 Conclusion/Discussion	20
14	6 Development of an whole cell methyl transferase screening sys-	
15	tem	21
16	6.1 Introduction	21
17	6.2 Theoretical considerations / design of system	21
18	6.3 Detectability of S-adenosyl-textscL-homocysteine (SAH).	21
19	6.4 Usage of the lsr-promoter for true autoinduction	21
20	6.5 Conclusion/Discussion	21
21	7 DES in protein crystallography	22
22	7.1 Introduction	22
23	7.2 Solubility enhancement of hydrophobic substances by addition of	
24	DES	22
25	7.3 Enzymatic O-methylation in DES	22
26	7.4 DES as precipitants in protein crystallization	22
27	7.5 Conclusion/Discussion	22
28	8 Acknowledgements	23
29	III Appendix	24
30		
31	A Figures	25
32	B Tables	26
33	C Affidavit	28

¹	Acronyms	31
²	Glossary	33



¹**List of Figures**

²	3.1 GOD assay	17
³	A.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean com-	
⁴	modo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus	
⁵	et magnis dis parturient montes, nascetur ridiculus mus. Donec quam	
⁶	felis, ultricies nec, pellentesque eu, pretium quis, sem.	25

List of Tables

3.1	Plasmids used in this work.	8
3.2	Primers used in this work. Recognition sites for endonucleases are underlined. Positions used for site directed mutagenesis are in lower case font.	9
B.1	Overview over the constructs produced for the present thesis. Each step during the production of the construct is given in the workflow steps column. Primers (italic font) or restriction sites used during each step are displayed in parenthesis.	27

¹ **Notes of Revisors**

²	 BWEIGEL: Überprüfen!	15
³	 BWEIGEL: complete	15

1

2

I

Preface


1 Abstracts

1.1 English Abstract

Test Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

1.2 Deutsche Zusammenfassung

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.



Thesis

1

2

¹ 2 Introduction

² Some introductory text

³ 2.1 Natural products and secondary metabolites

⁴ 2.1.1 General

⁵ 2.1.2 Classes of natural products

⁶ Terpenoids and Steroids

⁷ ... here is some text

⁸ Polyketides and non-ribosomal peptides

⁹ ... here is some text

¹⁰ Alkaloids

¹¹ ... here is some text

¹² Phenylpropanoids

¹³ ... here is some text

¹⁴ 2.2 Alkylating reactions in nature

¹⁵ 2.2.1 Methylation

¹⁶ 2.2.2 Prenylation

¹ 2.2.3 Glycosylation

² 2.3 Usage and expansion of nature's reaction toolbox
³

⁴ 2.3.1 Terpene synthases and elongases

⁵ 2.3.2 Methyl transferases

⁶ 2.3.3 Glycosyl transferases

⁷ 2.3.4 Other important enzymes in biotech research

⁸ BMVOs

⁹ Esterases/Lipases

¹⁰ Oxidases

¹¹ Lyases

¹² Transaminases

¹³ 2.4 Conclusion

3 Material And Methods

Within this section percentages refer to volume per volume (v/v) percentages unless otherwise specified.

3.1 Materials

3.1.1 Chemicals

Enzymes and buffers used for molecular cloning were obtained from Life Technologies (Darmstadt, Germany), unless otherwise noted. Flavonoid HPLC standards were purchased from Extrasynthese (Genay, France). Deuterated solvents were aquired from Deutero GmbH (Kastellaun, Germany). Solvents, purchased from VWR (Poole, England), were distilled in-house before use. All other chemicals were obtained from either Sigma-Aldrich (Steinheim, Germany), Applichem (Darmstadt, Germany), Carl Roth (Karlsruhe, Germany) or Merck (Darmstadt, Germany).

3.1.2 Commonly used solutions and buffers

50× 5052	25 % glycerol, 2.5 % (w/v) glucose, 10 % (w/v) α-lactose
binding buffer	50 mM Tris/HCl, 500 mM NaCl, 10 % glycerol, 2.5 mM imidazole pH 7
elution buffer	50 mM Tris/HCl, 500 mM NaCl, 10 % glycerol, 250 mM imidazole pH 7
lysis buffer	50 mM Tris/HCl, 500 mM NaCl, 10 % glycerol, 2.5 mM imidazole, 0.2 % Tween-20 pH 7
1 M MMT pH 4 (10×)	26.8 g/l L-malic acid, 78.1 g/l MES, 26.8 g/l Tris, 2.1 % 10 M HCl
1 M MMT pH 9 (10×)	26.8 g/l L-malic acid, 78.1 g/l MES, 26.8 g/l Tris, 6.7 % 10 M NaOH
20× NPS	1 M Na ₂ HPO ₄ , 1 M KH ₂ PO ₄ , 0.5 M (NH ₄) ₂ SO ₄

5× SDS sample buffer	10 % (w/v) SDS, 10 mM β-mercapto ethanol, 20 % glycerol, 0.2 M Tris/HCl pH 6.8, 0.05 % (w/v) bromophenolblue
1000× trace elements	50 mM FeCl ₃ , 20 mM CaCl ₂ , 10 mM MnCl ₂ , 10 mM ZnSO ₄ , 2 mM CoCl ₂ , 2 mM CuCl ₂ , 2 mM NiCl ₂ , 2 mM Na ₂ MoO ₄ , 2 mM Na ₂ SeO ₃ , 2 mM H ₃ BO ₃

1

3.1.3 Culture media used to grow bacteria

LB-medium	10 g/l NaCl, 10 g/l tryptone, 5 g/l yeast extract, pH 7.5
LB-agar	LB + 1.5 % (w/v) agar-agar
TB-medium	12 g/l tryptone, 24 g/l yeaxst extract, 0.4 % glycerol, 72 mM K ₂ HPO ₄ , 17 mM KH ₂ PO ₄
ZY	10 g/l tryptone, 5 g/l yeast extract
ZYP-5052	in proportions: 0.928 ZY, 0.05 20× NPS, 0.02 50× 5052, 0.002 1 M MgSO ₄ , 0.0002 1000× trace elements

2

3.1.4 Bacterial strains

3

E.coli

BL21(DE3)	F ⁻ <i>ompT hsdSB</i> (r _B ⁻ ,m _B ⁻) <i>gal dcm</i> λ(DE3) Invitrogen, Karlsruhe (Germany)
C41(DE3)	F ⁻ <i>ompT hsdSB</i> (r _B ⁻ ,m _B ⁻) <i>gal dcm</i> λ(DE3) Lucigen, Wisconsin (USA)
C43(DE3)	F ⁻ <i>ompT hsdSB</i> (r _B ⁻ ,m _B ⁻) <i>gal dcm</i> λ(DE3) Lucigen, Wisconsin (USA)
DH5α	F ⁻ Φ80 <i>lacZ</i> ΔM15 Δ(<i>lacZYA-argF</i>) U169 <i>recA1 endA1 hsdR17</i> (r _K ⁻ m _K ⁺) <i>phoA supE44</i> λ ⁻ <i>thi-1 gyrA96 relA1</i> Invitrogen, Karlsruhe (Germany)
JM110	<i>rpsL thr leu thi lacY galK galT ara tonA tsx dam dcm glnV44</i> Δ(<i>lac-proAB</i>) e14- [F' <i>traD36 proAB</i> ⁺ <i>lacI</i> ^q <i>lacZ</i> ΔM15] <i>hsdR17</i> (r _K ⁻ m _K ⁺) Martin-Luther-University Halle-Wittenberg
JW1593 (BW25113 derivative)	<i>rrnB</i> Δ <i>lacZ</i> 4787 <i>HsdR</i> 514 Δ(<i>araBAD</i>)568 <i>rph-1</i> Δ <i>ydgG</i> (Kan ^R) Keio Collection, National Institute of Genetics (Japan)
MG1655	F ⁻ λ ⁻ <i>ilvG</i> ⁻ <i>rfb-50 rph-1</i> DSMZ, Hamburg (Germany)

One Shot TOP10	F [−] Φ80 <i>lacZ</i> ΔM15 Δ(<i>mrr</i> - <i>hsdRMS</i> - <i>mcrBC</i>) <i>recA1 endA1 mcrA</i> Δ <i>lacX74 araD139</i> Δ(<i>ara-leu</i>)7697 <i>galU galK rpsL</i> (Str ^R) λ [−] <i>nupG</i> Invitrogen, Karlsruhe (Germany)
Origami(DE3)	Δ(<i>ara-leu</i>)7697 Δ <i>lacX74</i> Δ <i>phoA</i> <i>Pvull phoR araD139 ahpC galE galK rpsL</i> F' [<i>lac</i> + <i>lacI q pro</i>] (DE3) <i>gor522::Tn10 trxB</i> (Kan ^R , Str ^R , Tet ^R) Novagen, Wisconsin (USA)
Rosetta(DE3)	F [−] <i>ompT hsdSB</i> (r _B [−] ,m _B [−]) <i>gal dcm</i> λ(DE3) pRARE (Cam ^R) Novagen, Wisconsin (USA)
Rosetta(DE3) pLysS	F [−] <i>ompT hsdSB</i> (r _B [−] ,m _B [−]) <i>gal dcm</i> λ(DE3) pLysSRARE (Cam ^R) Novagen, Wisconsin (USA)
T7 Express	<i>fhuA2 lacZ::T7 gene1 [lon] ompT gal sulA11 R(mcr-73::miniTn10–Tet^S)2 [dcm] R(zgb-210::Tn10–Tet^S) endA1</i> Δ(<i>mcrC-mrr</i>)114::IS10 NEB, Massachusetts (USA)

¹ **Agrobacterium tumefaciens**

GV3101	chromosomal background: C58, marker gene: <i>rif</i> , Ti-plasmid: cured, opine: nopaline Sylvestre Marillonet, IPB
--------	--

² **3.1.5 Plasmids**

Table 3.1.: *Plasmids used in this work.*

name	supplier/source
pACYCDuet-1	Merck, Darmstadt (Germany)
pCDFDuet-1	Merck, Darmstadt (Germany)
pET-20b(+)	Merck, Darmstadt (Germany)
pET-28a(+)	Merck, Darmstadt (Germany)
pET-32a(+)	Merck, Darmstadt (Germany)
pET-41a(+)	Merck, Darmstadt (Germany)
pQE30	QIAGEN, Hilden (Germany)
pUC19	Invitrogen, Karlsruhe (Germany)

3.1.6 Oligonucleotides and synthetic genes

Oligonucleotides and primers were ordered from Eurofins Genomics (Ebersberg, Germany). The purity grade was *high purity salt free* (HPSF). Synthetic genes or gene fragments were obtained from GeneArt® (Life Technologies, Darmstadt, Germany) or Eurofins Genomics (Ebersberg, Germany).

Table 3.2.: Primers used in this work. Recognition sites for endonucleases are underlined. Positions used for site directed mutagenesis are in lower case font.

name	sequence (5'→3')	cloning site
somt1	TTG <u>AAG ACA</u> AAA TGG CTT CTT CAT TAA ACA ATG GCC G	BpiI
somt2	TTG <u>AAG ACA</u> AGG ACA CCC CAA ATA CTG TGA GAT CTT CC	BpiI
somt3	TTG <u>AAG ACA</u> AGT CCT TAG GAA CAC CTT TCT GGG AC	BpiI
somt4	TTG <u>AAG ACA</u> AAA GCT CAA GGA TAG ATC TCA ATA AGA GAC	BpiI
pfomt1.fw	CAG AGA GGC cTA TGA GAT TGG CTT GC	
pfomt1.rv	GCA AGC CAA TCT CAT AgG CCT CTC TG	
pfomt2.fw	<u>CAT ATG</u> GAT TTT GCT GTG ATG AAG CAG GTC	NdeI
pfomt2.rv	<u>GAA TTC</u> AAT AAA GAC GCC TGC AGA AAG TG	EcoRI
pRha1.fw	CTC TAG <u>CAG ATC</u> <u>TCG</u> GTG AGC ATC ACA TCA CCA CAA TTC	BglII
pRha1.rv	CAA TTG <u>AGG ATC</u> <u>CCC</u> ATT TTA ACC TCC TTA GTG	BamHI
pUC1.fw	GCG TAT TGG Gag aTC TTC CGC TTC CTC	
pUC1.rv	GAG GAA GCG GAA GAt ctC CCA ATA CGC	

3.1.7 Instruments

CD-spectrometer	Jasco J-815 (Eaton, USA)
electrophoresis (horizontal)	Biometra Compact XS/S (Göttingen, Germany)
electrophoresis (vertical)	Biometra Compact M (Göttingen, Germany) Biometra Minigel-Twin (Göttingen, Germany)
FPLC	ÄKTA purifier (GE Healthcare, Freiburg, Germany)
GC/MS	GC-MS-QP2010 Ultra (Shimadzu, Duisburg, Germany)
HPLC	VWR-Hitachi LaChrom Elite (VWR, Darmstadt, Germany)
ITC	MicroCal iTC200 (Malvern, Worcestershire, UK)
plate-reader	SpectraMax M5 (Molecular Devices, Biberach, Germany)
NMR-spectrometer	Varian Unity 400 (Agilent, Böblingen, Germany) Varian VNMRS 600 (Agilent, Böblingen, Germany)

photospectrometer	Eppendorf Biophotometer Plus (Hamburg, Germany) JASCO V-560 (Eaton, USA) Colibri Microvolume Spectrometer (Biozym, Hess. Oldendorf, Germany)
centrifuges	Eppendorf 5424 (Hamburg, Germany) Hettich Mikro 120 (Kirchlengern, Germany) Beckman Avanti J-E, Beckman Allegra X-30R (Krefeld, Germany)
centrifuge rotors	Beckman JA-10, JA-16.250, JS-4.3 (Krefeld, Germany)

3.1.8 Software

All mathematical and statistical computations and graphics were done with the R software (versions 3.1.X, <http://cran.r-project.org/>). Visualizations of macromolecules were arranged using the PyMol Molecular Graphics System, version 1.7.0.0 (Schrödinger, New York, USA). Physicochemical calculations and calculations of different molecular descriptors were performed using Marvin Beans 15.4.13.0 (ChemAxon, Budapest, Hungary) and Molecular Operating Environment 2008.10 (Chemical Computing Group, Montreal, Canada).

3.2 Microbiology

3.3 Molecular Biology

Basic molecular biology methods like polymerase chain reaction (PCR), DNA restriction/ligation, DNA gel electrophoresis, preparation of competent cells and transformation were performed based on the protocols summarized by Sambrook and Russell [14]. Plasmid DNA was isolated using the QIAprep[®] Spin Miniprep Kit (QIAGEN, Hilden, Germany) according to the manufacturer's instructions. In vitro site-directed mutagenesis was set-up according to the protocol of the QuikChange[™] Site-Directed Mutagenesis kit [1] offered by Agilent Technologies (Santa Clara, USA). Nucleotide fragments obtained by PCR, restriction/ligation procedures or excision from electrophoresis gels were purified and concentrated using the Nucleospin Gel and PCR Clean-up kit provided by Machery-Nagel (Düren, Germany) according to the instructions provided by the manufacturer.

3.3.1 Golden Gate Cloning

The Golden Gate cloning procedure is a one-pot method, meaning the restriction digestion and ligation are carried out in the same reaction vessel at the same time [6, 4]. Consequently PCR-fragments, destination vector, restriction endonuclease and ligase are added together in this reaction. The methodology employs type II restriction enzymes, which together with proper design of the fragments allow for a ligation product lacking the original restriction sites.

For digestion/ligation reactions of fragments containing BpiI sites, 20 fmol of each fragment or vector, together with 5 U of BpiI and 5 U of T4 ligase were combined in a total volume of 15 µl 1× ligase buffer. For fragments to be cloned via BsaI sites, BpiI in the above reaction was substituted by 5 U BsaI.

The reaction mixture was placed in a thermocycler and incubated at 37 °C for 2 min and 16 °C for 5 min. These two first steps were repeated 50 times over. Finally, the temperature was raised to 50 °C (5 min) and 80 °C (10 min) to inactivate the enzymes.

3.3.2 Subcloning of genes

All subcloning procedures were performed according to section 3.3 and specifically subsection 3.3.1. Specific steps for the subcloning of any genes discussed can be found in the appendix (p.27). The *pfomt* gene was subcloned from the pQE-30 vector kindly provided by Thomas Vogt (Leibniz-Institute of Plant Biochemistry (IPB), Halle, Germany) into the pET-28a(+) vector. The *somt-2* gene was subcloned from the pQE-30 vector kindly provided by Martin Dippe (IPB, Halle, Germany) into the pET-28-MC vector.

3.4 Protein biochemistry

Stock solutions of antibiotics, IPTG or sugars were prepared according to the pET System Manual by Novagen [11], unless otherwise noted.

3.4.1 Protein production test (expression test)

The heterologous production of proteins in *E. coli* was assessed in a small scale protein production test, henceforth called expression test. Single colonies of *E. coli* transformed with the constructs to be studied were used to inoculate a 2 ml starter culture in lysogeny broth (LB)-medium containing the appropriate antibiotics. The concentrations of antibiotics used was as follows: 200 µg/ml ampicillin, 150 µg/ml kanamycin, 50 µg/ml chloramphenicol, 20 µg/ml tetracycline.

The starter culture was allowed to grow at 37 °C and 200 rpm over night. A 5 ml sampling culture of the medium to be studied containing the appropriate antibiotics was prepared. The media tested included LB, terrific broth (TB) and auto-induction media like ZYP-5052. The sampling culture was inoculated to an OD⁶⁰⁰ of 0.075 using the starter culture and incubated at different temperatures and 200 rpm in a shaking incubator. 1 mM isopropyl-D-thiogalactopyranosid (IPTG) was added when the OD⁶⁰⁰ reached 0.6-0.8, if appropriate for the studied construct. 1 ml samples were removed after different times of incubation (e.g. 4, 8, 12 hours), subfractionated (3.4.2) and analyzed via SDS-polyacrylamide gel electrophoresis (PAGE) (3.4.5).

Exact specifications of growth conditions (e.g. temperature, time, constructs) are discussed in the individual sections.

3.4.2 Protein subfractionation

The protein subfractionation procedure described herein was adapted from the protocol described in the pET Manual [11]. Overall 5 protein subfractions can be obtained, including *total cell protein*, *culture supernatant (medium) protein*, *periplasmic protein*, *soluble cytoplasmic protein* and *insoluble protein*.

The OD⁶⁰⁰ of the culture sample was measured and the cells harvested by centrifugation at 10000 × *g*, 4 °C for 5 minutes. The protein in the supernatant medium was concentrated by precipitation with trichloro acetic acid (TCA) (3.4.3) for SDS-PAGE analysis. The periplasmic protein was prepared (3.4.4) and also concentrated by TCA precipitation for SDS-PAGE. Cells were lysed by resuspending the cell pellet in (OD⁶⁰⁰ × *V* × 50) µl of bacterial protein extraction reagent (B-PER) and vortexing vigorously for 30 s. The suspension was incubated at room temperature (RT) for 30 min to assure complete lysis. To separate insoluble protein and cell debris from the soluble cytosolic protein, the suspension was centrifuged at 10000 × *g* and 4 °C for 10 min. Soluble cytoplasmic protein was contained in the supernatant, whereas insoluble protein remained in the pellet. For SDS-PAGE analysis of the insoluble protein, the pellet was resuspended in the same volume of B-PER. To obtain only the total cell protein fraction, the preparation of periplasmic and soluble cytosolic protein was omitted. Sample volumes of 10 µl of each fraction were used for SDS-PAGE analysis.

3.4.3 Protein sample concentration by TCA precipitation

Diluted protein samples were concentrated by TCA precipitation in microcentrifuge tubes. Therefore 0.1 volume (*V*) of 100 % (w/v) TCA in water was added to the clarified sample, which was then vortexed for 15 s and placed on ice for a minimum

of 15 min. The sample was centrifuged at $14000 \times g$, 4°C for 15 min. The supernatant was discarded and the pellet was washed twice with 0.2 V ice-cold acetone. The acetone was removed and the pellet set to air-dry in an open tube. After drying, the protein pellet was resuspended in 0.1 V phosphate buffered saline (PBS) containing $1 \times$ SDS-sample buffer by heating to 85°C and vigorous vortexing, to achieve a $10 \times$ concentration. After resuspension the sample was analyzed by SDS-PAGE or stored at -20°C until use.

3.4.4 Preparation of periplasmic protein

Target proteins may be directed to the periplasmic space by N-terminal signal sequences like *pelB* or *DsbA/C* [9]. The periplasma is, other than the cytosol, an oxidizing environment and often used for the production of proteins containing disulfide linkages. The preparation of periplasmic protein was accomplished by an osmotic shock protocol modified from Current Protocols in Molecular Biology [2]. The cell pellet was resuspended in the same volume as the culture sample of 30 mM tris-HCl, 20 % (w/v) sucrose, pH 8 and 1 mM ethylenediaminetetraacetic acid (EDTA) was added. The suspension was stirred for 10 min at RT and the cells were collected by centrifugation at $10000 \times g$, 4°C for 10 min. The supernatant was discarded and the cell pellet was resuspended in the same volume of ice-cold 5 mM MgSO_4 . The suspension was stirred for 10 min on ice, while the periplasmic proteins were released into the solution. The cells were collected by centrifugation as before. Periplasmic proteins were contained in the supernatant.

3.4.5 Discontinuous SDS-polyacrylamide gel electrophoresis (SDS-PAGE)

The analysis of samples via SDS-PAGE was realized via the discontinuous system first described by Laemmli, which allows separation of proteins based on their electrophoretic mobility, which in turn depends on their size [8]. The SDS-PAGE procedure was carried out according to standard protocols described by Sambrook and Russell [14]. Very dilute and/or samples with high ionic strength were concentrated and/or desalted by the TCA precipitation procedure described in subsection 3.4.3. Generally a 10 % (acrylamide/bisacrylamide) running gel combined with a 4 % stacking gel was used. Reducing SDS-PAGE sample buffer was added to the protein sample to be analyzed, whereafter the sample was heated to 95°C for 5 min, to allow for total unfolding of the protein. After cooling to RT the samples were transferred into the gel pockets for analysis. The *PageRuler™ Prestained Protein Ladder* (Life Technologies GmbH, Darmstadt, Germany) was used as a molecular weight (MW) marker and run alongside every analysis as a reference.

¹ Gels were stained using a staining solution of 0.25 % Coomassie Brilliant Blue G-
² 250 (w/v) in water:methanol:acetic acid (4:5:1) and destained by treatment with
³ water:methanol:acetic acid (6:3:1).

⁴ 3.4.6 Buffer change of protein samples

⁵ The buffer in protein samples was exchanged either by dialysis, or by centrifugal
⁶ filter concentrators (Amicon® Ultra Centrifugal Filter; Merck, Darmstadt, Ger-
⁷ many). Large volumes of highly concentrated protein solutions were preferably
⁸ dialyzed. Respectively, very dilute samples were concentrated and rebuffed using
⁹ centrifugal concentrators.

¹⁰ Dialysis was carried out at least twice against a minimum of 100 times the sample
¹¹ volume. Dialysis steps were carried out at RT for 2 hours, or over-night at 4 °C.
¹² Centrifugal concentrators were used according to the manufacturers instructions.

¹³ 3.4.7 Production of recombinant protein

¹⁴ Heterologous production of PFOMT

¹⁵ Phenylpropanoid and flavonoid O-methyl transferase (PFOMT) was produced as a
¹⁶ N-terminally (His)₆-tagged fusion protein. A 2 ml starter culture of LB containing
¹⁷ 100 µg/ml kanamycin was inoculated with a single colony of *E. coli* BL21(DE3)
¹⁸ transformed with pET28-pfomt and incubated at 37 °C, 220 rpm for 6 hours. The
¹⁹ main culture (N-Z-amine, yeast extract, phosphate (ZYP-5052) containing 150 µg/ml
²⁰ kanamycin) was inoculated with the starter culture such that OD⁶⁰⁰ was 0.05.
²¹ The culture was incubated in a shaking incubator at 37 °C, 220 rpm over night
²² (≈16 h). Due to the autoinducing nature of the ZYP-5052 medium, addition of
²³ IPTG was not necessary. Cells were harvested by centrifugation at 10000 × *g*, 4 °C
²⁴ for 10 min and the supernatant discarded. The pellet was resuspended in 50 mM
²⁵ Tris/HCl, 500 mM NaCl, 2.5 mM imidazole, 10 % glycerol pH 7 using a volume of
²⁶ ≈10 ml/g of cell pellet. The cells were lysed by sonication (70 % amplitude, 1 s on-
²⁷ off-cycle) for 30 seconds, which was repeated twice. The crude lysate was clarified
²⁸ by centrifugation at 15000 × *g*, 4 °C for 15 minutes followed by filtration through a
²⁹ 0.45 µm filter. Consequently, the His-tagged PFOMT was purified by immobilized
³⁰ metal affinity chromatography (IMAC) (3.4.9). The eluted PFOMT protein was
³¹ dialyzed (3.4.6) against 25 mM HEPES, 100 mM NaCl, 5 % glycerol pH 7 and stored
³² at -20 °C until use.

1 Heterologous production of SOMT-2

2 Soy O-methyl transferase (SOMT-2) was produced as a fusion protein with an N-
3 terminal His-tag. A starter LB-culture (≈ 2 ml) containing 100 $\mu\text{g/ml}$ kanamycin was
4 inoculated with a single colony of *E. coli* BL21(DE3) transformed with pET28MC-
5 somt and incubated at 37 °C, 220 rpm for 6 hours. The starter culture was used to
6 inoculate the main culture (LB-medium containing 100 $\mu\text{g/ml}$ kanamycin), such
7 that $\text{OD}^{600} \approx 0.05$. The culture was incubated at 37 °C, 220 rpm in a shaking in-
8 cubator until $\text{OD}^{600} \approx 0.6$. Expression was induced by addition of 1 mM IPTG.
9 Incubation continued at 37 °C, 220 rpm for 6 hours. Cells were harvested by cen-
10 trifugation ($10000 \times g$, 4 °C, 10 min) and used, or stored at -20 °C until use. SOMT-2
11 was produced only in inclusion bodies (IBs), which were prepared as laid out in
12 subsection 3.4.8.

BWEIGEL: Überprüfen!

13 3.4.8 Preparation of IB

14 When insoluble protein is produced in large amounts in *E. coli* it is accumulated
15 in so-called inclusion bodies (IBs). The accumulating IBs consist mainly of the
16 overproduced target protein, which is inherently quite pure already. IBs can be
17 prepared [12]

BWEIGEL: complete

18 3.4.9 Purification of His-tagged proteins using immobilized 19 metal affinity chromatography (IMAC)

20 N- or C-terminal oligo-histidine tags (His-tags) are a common tool to ease purifica-
21 tion of recombinantly produced proteins. The free electron pairs of the imidazol
22 nitrogens of histidines can complex divalent cations such as Mg^{2+} or Ni^{2+} , which
23 are usually immobilized on a matrix of nitrilo triacetic acid (NTA)-derivatives. The
24 affinity of the His-tag is correlated with its length and tagged proteins can simply
25 be eluted by increasing the concentration of competing molecules (e.g. imidazole).
26 His-tagged protein was purified by fast protein liquid chromatography (FPLC) via
27 Ni^{2+} - (HisTrap FF crude) or Co^{2+} -NTA (HiTrap Talon FF crude) columns, obtained
28 from GE Healthcare (Freiburg, Germany), following modified suppliers instructions.
29 First the column was equilibrated with 5 column volumes (CV) of binding buffer
30 (50 mM Tris/HCl, 500 mM NaCl, 10 % glycerol, 2.5 mM imidazole pH 7). The sample
31 (generally clarified lysate) was applied to the column using a flow of 0.75 ml/min.
32 Unbound protein was removed by washing with 3 CV binding buffer. Unspecificall
33 bound proteins were washed away by increasing the amount of elution buffer
34 (50 mM Tris/HCl, 500 mM NaCl, 10 % glycerol, 250 mM imidazole pH 7) to 10 %

¹ (constant for 3 to 5 CV). Highly enriched and purified target protein was eluted
² with 6 to 10 CV of 100 % elution buffer.

³ 3.5 Analytics

⁴ 3.5.1 In vitro determination of glucose


⁵ The glucose concentration in clarified, aqueous samples was determined by a
⁶ modified version of the glucose assay kit procedure provided by Sigma-Aldrich
⁷ [15]. Glucose oxidase (GOD) oxidizes D-glucose to gluconic acid, whereby hydrogen
⁸ peroxide is produced. The hydrogen peroxide can be detected and quantified by
⁹ horseradish peroxidase (HRP), which reduces the produced H_2O_2 and thereby
¹⁰ oxidizes its chromogenic substrate *o*-dianisidine. The oxidized form of *o*-dianisidine
¹¹ can then be measured photospectrometrically [3].

¹² The methodology employs a coupled photospectrometric assay using GOD and
¹³ HRP with *o*-dianisidine as reporter substrate. The assay was prepared in micro-titer
¹⁴ plate (MTP)-format. A reaction solution containing 12.5 U/ml GOD, 2.5 U/ml HRP
¹⁵ and 0.125 mg/ml *o*-dianisidine dihydrochloride in 50 mM sodium acetate pH 5.1
¹⁶ was prepared.

¹⁷ Sample solutions from culture supernatants were typically diluted in 9 volumes
¹⁸ of water. The reaction was started, by adding 50 μ l reaction solution to 25 μ l of
¹⁹ sample and was incubated at 37 °C and 200 rpm for 30 min in a shaking incubator.
²⁰ 50 μ l 6 M sulfuric acid was added to stop the reaction and achieve maximum color
²¹ development (full oxidation of any *o*-dianisidine charge transfer complexes).

²² The developed pink color was measured at 540 nm in a MTP-reader. A calibration
²³ curve of a standard D-glucose solutions (0 to 100 μ g/ml), that was always part of
²⁴ the experiments, was used to quantify the sample measurements.





² 3.5.2 HPLC measurements

¹4 **Evaluation of PFOMT towards** ²**the acceptance of long-chain SAM** ³**analogues**

⁴4.1 **Introduction**

⁵4.2 **Substrate binding studies using ITC**

⁶4.3 **Determination of the structure of *apo*-PFOMT**

⁷4.3.1 **PFOMT activity in deep eutectic solvent (DES) / Solubility-** ⁸**enhancing effects of DES**

⁹vielleicht eigenes kapitel DES?

¹⁰4.4 **Study of variants for long-chain alkylations**

¹¹4.4.1 **PFOMT-Paper (DIM)**

¹²4.4.2 **Dockings???**

¹³4.5 **Colclusion/Discussion**

¹ 5 Enzymatic methylation of Non- ² catechols

³ 5.1 Introduction

⁴ Non-catechols in nature (biosynthesis, mode of action?), chemical methylation???

⁵ 5.2 SOMT-2

⁶ 5.2.1 In vivo methylation studies using *N. benthamiana*

⁷ 5.2.2 In vivo studies in *E. coli*

⁸ 5.2.3 In vitro studies using recombinantly produced SOMT-2

⁹ 5.3 PFOMT

¹⁰ 5.3.1 Acidity and Nucleophilicity of phenolic hydroxyl-groups

¹¹ 5.3.2 pH-Profiles of PFOMT-catalysis

¹² 5.3.3 Influence of Mg^{2+} on PFOMT activity

¹³ 5.4 Consensus or Bioinformatic points-of-view
¹⁴ (COMT)???

¹⁵ 5.5 Conclusion/Discussion

¹ **6 Development of an whole cell** ² **methyl transferase screening sys-** ³ **tem**

⁴ **6.1 Introduction**

⁵ **6.2 Theoretical considerations / design of system**

⁶ **6.3 Detectability of *S*-adenosyl-textscL-homocysteine** ⁷ **(SAH)**

⁸ *S*-adenosyl-L-methionine (SAM)

⁹ **6.4 Usage of the *lsr*-promoter for true autoinduc-** ¹⁰ **tion**

¹¹ **6.5 Conclusion/Discussion**

¹ **7 DES in protein crystallography**

² **7.1 Introduction**

³ **7.2 Solubility enhancement of hydrophobic substances by addition of DES** ⁴

⁵ **7.3 Enzymatic *O*-methylation in DES**

⁶ **7.4 DES as precipitants in protein crystallization**

⁷ **7.5 Conclusion/Discussion**

8 Acknowledgements

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt
ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse
platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum
fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin,
felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin
tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.
Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus
semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam.
Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam,
in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor.
Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi
fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui
cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac
magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc.
Nam feugiat lacus vel est. Curabitur consectetur.



Appendix

1

2

¹A Figures




Figure A.1.: Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem.

B Tables

Table B.1.: Overview over the constructs produced for the present thesis. Each step during the production of the construct is given in the workflow steps column. Primers (*italic font*) or restriction sites used during each step are displayed in parenthesis.

construct name	description	entry constructs	destination	workflow steps (primers/cloning sites)
pBEW101	lsrA promoter pBEW102 with BamHI cloning site rhaP _{BAD} promoter	pBEW102 pBEW4b	pBEW103	amplification (<i>pRha1.fw/rv</i>), cloning (BglII, BamHI)
pBEW102				
pBEW103				
pBEW104				
pBEW106	pICH413038-somt	pET28MC-somt	pICH413038	amplification (<i>somt1/2/3/4</i>), golden gate cloning (Bpil)
pBEW107		pICH51266, pBEW106, pICH41421	pICH75044	golden gate cloning (BsaI)
pBEW1a				
pBEW1b				
pBEW2a				
pBEW2b				
pBEW3a				
pBEW3b				
pBEW4a				
pBEW4b				
pET28-pfomt	<i>pfomt</i> gene in pET-28a(+), endogenous NdeI site removed	pQE30-pfomt	pET-28a(+)	mutagenesis (<i>pfomt1.fw/rv</i>), amplification (<i>pfomt2.fw/rv</i>), cloning (NdeI, EcoRI)
pET20-somt	N-terminal pelB-tag fusion for periplasmic expression		pET20-b(+)	
pET28-somt			pET28-a(+)	
pET28MC-somt	N-terminal Trx-tag fusion N-terminal GST-tag fusion added BglII site pUC19 derivative with <i>lsrA</i> promoter	pUC19 lsr-XX-DAS	pET-32a(+)	mutagenesis (<i>pUC1.fw/rv</i>) cloning (NdeI, BglII)
pET32-somt			pET-41a(+)	
pET41-somt			-	
pUC19*			pUC19*	
pUCB1				
pUCB1-sfGFP-DAS+4				

¹ **C Affidavit**

² I hereby declare that this document has been written only by the undersigned and
³ without any assistance from third parties. Furthermore, I confirm that no sources
⁴ have been used in the preparation of this document other than those indicated in
⁵ the thesis itself.

⁶ Date:....., Location:....., Signature:.....

Bibliography

- [1] Agilent Technologies. *QuikChange II Site-Directed Mutagenesis Kit: Instruction Manual*. 2011.
- [2] F Ausubel et al. “Current Protocols in Molecular Biology”. In: (2008), p. 23.
- [3] a Claiborne and Irwin Fridovich. “of of of”. In: *Society* 18 (1979), pp. 2324–2329.
- [4] Carola Engler, Romy Kandzia, and Sylvestre Marillonnet. “A one pot, one step, precision cloning method with high throughput capability”. In: *PLoS ONE* 3.11 (Jan. 2008), e3647.
- [5] Mwafaq Ibdah et al. “A Novel Mg²⁺-dependent O-Methyltransferase in the Phenylpropanoid Metabolism of *Mesembryanthemum crystallinum*”. In: *Journal of Biological Chemistry* 278.45 (Nov. 2003), pp. 43961–43972.
- [6] Youichi Kondou et al. “cDNA Libraries”. In: *Methods* 729 (2011), pp. 183–197.
- [7] Jakub G. Kopycki et al. “Biochemical and Structural Analysis of Substrate Promiscuity in Plant Mg²⁺-Dependent O-Methyltransferases”. In: *Journal of Molecular Biology* 378.1 (Apr. 2008), pp. 154–164.
- [8] U K Laemmli. “Cleavage of structural proteins during the assembly of the head of bacteriophage T4.” In: *Nature* 227.5259 (1970), pp. 680–685.
- [9] Sc Makrides and Sc Makrides. “Strategies for achieving high-level expression of genes in *Escherichia coli*”. In: *Microbiol. Rev.* 60.3 (1996), pp. 512–538.
- [10] Janet Newman. “Novel buffer systems for macromolecular crystallization”. In: *Acta Crystallographica Section D: Biological Crystallography* 60.3 (2004), pp. 610–612.
- [11] Novagen. *pET System Manual*. 11th ed. Vol. 123. 2 Pt 1. Darmstadt: EMD Biosciences, 2014, p. 369.
- [12] Ira Palmer and Paul T. Wingfield. “Preparation and extraction of insoluble (Inclusion-body) proteins from *Escherichia coli*”. In: *Current Protocols in Protein Science* 1.SUPPL.70 (Nov. 2012), Unit6.3.

- ¹ [13] R Rudolph and H Lilie. “In vitro folding of inclusion body proteins.” In: *The*
² *FASEB journal : official publication of the Federation of American Societies for*
³ *Experimental Biology* 10.1 (1996), pp. 49–56.
- ⁴ [14] J Sambrook and D W Russell. *Molecular Cloning: A Laboratory Manual*.
⁵ 3rd ed. Cold Spring Harbor (NY, USA): Cold Spring Harbor Laboratory Press,
⁶ 2001.
- ⁷ [15] Sigma-Aldrich. *Technical bulletin UDP-Galactosyltransferase Assay Kit*.
- ⁸ [16] F William Studier. “Protein production by auto-induction in high density
⁹ shaking cultures.” In: *Protein expression and purification* 41.1 (May 2005),
¹⁰ pp. 207–234. arXiv: NIHMS150003.
- ¹¹ [17] Thomas Vogt. “Regiospecificity and kinetic properties of a plant natural
¹² product O-methyltransferase are determined by its N-terminal domain”. In:
¹³ *FEBS Letters* 561.1-3 (Mar. 2004), pp. 159–162.

¹ Acronyms

² **B-PER** bacterial protein extraction reagent

³ **CD** circular dichroism 9

⁴ **CV** column volumes

⁵ **EDTA** ethylenediaminetetraacetic acid 13

⁶ **FPLC** fast protein liquid chromatography 15

⁷ **GOD** glucose oxidase 16, 33

⁸ **HEPES** 2-[4-(2-hydroxyethyl)piperazin-1-yl]ethanesulfonic acid

⁹ **HRP** horseradish peroxidase 16

¹⁰ **IB** inclusion body 15

¹¹ **IMAC** immobilized metal affinity chromatography

¹² **IPB** Leibniz-Institute of Plant Biochemistry

¹³ **IPTG** isopropyl-D-thiogalactopyranosid 12, 14, 15

¹⁴ **LB** lysogeny broth 11, 12, 14, 15

¹⁵ **MES** 2-(N-morpholino)ethanesulfonic acid

¹⁶ **MMT** L-malic acid/MES/Tris 6

¹⁷ **MTP** micro-titer plate 16, 33

¹⁸ **MW** molecular weight 13

¹⁹ **NPS** nitrogen, phosphate, sulfate buffer

²⁰ **NTA** nitrilo triacetic acid 15

²¹ **PAGE** polyacrylamide gel electrophoresis 12, 13

²² **PBS** phosphate buffered saline 13

²³ **PCR** polymerase chain reaction 10, 11

²⁴ **PFOMT** phenylpropanoid and flavonoid O-methyl transferase 14, 33

- ¹ **RT** room temperature 12–14
- ² **SAM** S-adenosyl-L-methionine 21
- ³ **SDS** sodium dodecylsulfate 7, 12, 13
- ⁴ **SOMT-2** soy O-methyl transferase 15
- ⁵ **TB** terrific broth 12
- ⁶ **TCA** trichloro acetic acid 12, 13
- ⁷ **Ti-plasmid** tumor inducing plasmid 8, 33
- ⁸ **Tris** tris(hydroxymethyl)-aminomethane
- ⁹ **V** volume
- ¹⁰ **ZYP** N-Z-amine, yeast extract, phosphate 14, 33

¹ Glossary

² **GOD** Glucose oxidase is an enzyme.... ³¹

³ **MTP** Micro-titer plate. Small format rectangular plastic plate containing wells
⁴ to allow for storage of multiple small samples or the containment multiple
⁵ simultaneous reactions. Typical sizes include 24, 96 and 384-wells ³¹

⁶ **PFOMT** Phenylpropanoid and flavonoid O-methyl transferase from *Mesembryan-*
⁷ *themum crystallinum*, which was first described by Ibdah et al. in 2003 [5]
⁸ ³¹

⁹ **Ti-plasmid** Commonly found plasmids in *A. tumefaciens* and *A. rhizogenes* that
¹⁰ confer virulence ³²

¹¹ **ZYP-5052** Autoinduction medium developed by Studier [16]. The naming stems
¹² from the components N-Z-amine, yeast extract and phosphate. The numbering
¹³ designates the composition; e.g. 5052 refers to 0.5 % glycerol, 0.05 % glucose and
¹⁴ 0.2 % lactose. ³²