Biotransformations from and to methylated flavonoids

Hpw all went to shit

Benjamin Weigel Leibniz-Institute of Plant Biochemistry Department of Bioorganic Chemistry Weinberg 3 06120 Halle(Saale) April 20, 2015

Advisor: Prof. Dr. Ludger A. Wessjohann wessjohann@ipb-halle.de +49 (345) 5582-1301

noch nicht bekannt

Contents

I	Preface	1
1	Abstracts	3
	1.1 English Abstract	3
	1.2 Deutsche Zusammenfassung	3
I	Thesis	5
2	Introduction	7
	2.1 Natural products and secondary metabolites	7
	2.1.1 General	7
	2.1.2 Classes of natural products	7
	2.2 Alkylating reactions in nature	7
	2.2.1 Methylation	7
	2.2.2 Prenylation	7
	2.2.3 Glycosylation	8
	2.3 Usage and expansion of natures reaction toolbox	8
	2.3.1 Terpene synthases and elongases	8
	2.3.2 Methyl transferases	8
	2.3.3 Glycosyl transferases	8
	2.3.4 Other important enzymes in biotech research	8
	2.4 Conclusion	8
3	Material And Methods	9
	3.1 Materials	9
	3.1.1 Chemicals	9

	3.1.2 Instruments	10
	3.1.3 Bacterial strains	10
	3.1.4 Plasmids	12
	3.1.5 Oligonucleotides	12
	3.1.6 Software	12
	3.2 Microbiology	13
	3.3 Molecular Biology	13
	3.3.1 Golden Gate Cloning	13
	3.4 Protein biochemistry	14 14
	3.5 Analytics	14
4	Evaluation of PFOMT towards the acceptance of long-chain SAM	
	analogues	15
5	Enzymatic methylation of Non-catechols	17
6	Development of an whole cell methyl transferase screening sys-	
	tem	19
7	Acknowledgements	21
I	II Appendix	23
_	- Appendix	
A	Figures	25
В	Tables	27
С	Affidavit	29
	Acronyms	33
	•	
	Glossary	35

List of Figures

A.1	Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Aenean com-	
	modo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus	
	et magnis dis parturient montes, nascetur ridiculus mus. Donec quam	
	felis, ultricies nec, pellentesque eu, pretium quis, sem	25

List of Tables

3.1	Plasmids used in this work	12
3.2	Primers used in this work. Recognition sites for endonucleases are underlined	12
B.1	Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem	27

Preface

1 Abstracts

1.1 English Abstract

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetuer id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

1.2 Deutsche Zusammenfassung

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Thesis

2 Introduction

S ome introductionary text

2.1 Natural products and secondary metabolites

2.1.1 General

2.1.2 Classes of natural products

Terpenoids and Steroids

... here is some text

Polyketides and non-ribosomal peptides

... here is some text

Alkaloids

... here is some text

Phenylpropanoids

... here is some text

2.2 Alkylating reactions in nature

2.2.1 Methylation

2.2.2 Prenylation

- 2.2.3 Glycosylation
- 2.3 Usage and expansion of natures reaction toolbox
- 2.3.1 Terpene synthases and elongases
- 2.3.2 Methyl transferases
- 2.3.3 Glycosyl transferases
- 2.3.4 Other important enzymes in biotech research BMVOs

Esterases/Lipases

Oxidases

Lyases

Transaminases

2.4 Conclusion

3 Material And Methods

3.1 Materials

3.1.1 Chemicals

Enzymes and buffers used for molecular cloning were obtained from Thermo Scientific (Darmstadt, Germany), unless otherwise noted. Flavonoid HPLC standards were purchased from Extrasynthese (Genay, France). Deuterated solvents were aquired from Deutero GmbH (Kastellaun, Germany). Solvents, purchased from VWR (Poole, England), were distilled in-house before use.

All other chemicals were obtained from either Sigma-Aldrich (Steinheim, Germany), Applichem (Darmstadt, Germany), Carl Roth (Karlsruhe, Germany) or Merck (Darmstadt, Germany).

3.1.2 Instruments

circulary dichroism (CD)- Jasco J-815 (Eaton, USA)

spectrometer

electrophoresis (horizontal) Biometra Compact XS/S (Göttingen, Germany) electrophoresis (vertical) Biometra Compact M (Göttingen, Germany) Biometra Minigel-Twin (Göttingen, Germany)

fast protein liquid chromatogra-

phy (FPLC)

gas chromatography coupled GC-MS-QP2010 Ultra (Shimadzu, Duisburg, Germany)

mass-spectrometry (GC/MS)

high-performance liquid chro-

matography (HPLC)

Isothermal Titration Calorime-MicroCal iTC200 (Malvern, Worcestershire, UK)

VWR ???

try (ITC)

micro-titer plate (MTP) ???

MTP-reader SpectraMax M5 (Molecular Devices, Biberach, Germany)

nuclear magnetic resonance

Varian Unity 400 (Agilent, Böblingen, Germany) Varian VNMRS 600 (Agilent, Böblingen, Germany) (NMR)-spectrometer photospectrometer

Eppendorf Biophotometer Plus (Hamburg, Germany)

ÄKTA purifier (GE Healthcare, Freiburg, Germany)

JASCO V-560 (Eaton, USA)

Nanodrop???

Eppendorf 5424 (Hamburg, Germany) centrifuges

> Hettich Mikro 120 (Kirchlengern, Germany) Beckman Avanti J-E (Krefeld, Germany)

centrifuge rotors Beckman JA-10, JA-16.250, JS-4.3 (Krefeld, Germany)

3.1.3 Bacterial strains

E.coli

BL21(DE3) $F^- ompT \ hsdSB(r_R^-, m_R^-) \ gal \ dcm \ \lambda(DE3)$

Invitrogen, Karslruhe

C41(DE3) $F^- ompT \ hsdSB(r_B^-, m_B^-) \ gal \ dcm \ \lambda(DE3)$

Lucigen, Wisconsin (USA)

C43(DE3) $F^- ompT \ hsdSB(r_B^-, m_B^-) \ gal \ dcm \ \lambda(DE3)$

Lucigen, Wisconsin (USA)

DH5 α F⁻ Φ 80 $lacZ\Delta$ M15 Δ (lacZYA-argF) U169 recA1 endA1

 $hsdR17(r_K^-m_K^+)$ phoA supE44 λ^- thi-1 gyrA96 relA1

Invitrogen, Karlsruhe (Germany)

JM110 rpsL thr leu thi lacY galK galT ara tonA tsx dam

 $dcm \ glnV44 \ \Delta(lac\text{-}proAB) \ e14\text{--} [F' \ traD36 \ proAB^+ \ lacI^q$

 $lacZ\Delta M15$] $hsdR17(r_K^-m_K^+)$

Martin-Luther-University Halle-Wittenberg

JW1593 $rrnB \Delta lacZ4787 HsdR514 \Delta (araBAD)568 rph-1 \Delta ydgG$

(BW25113 derivative) (Kan^R)

Keio Collection, National Institute of Genetics (Japan)

MG1655 $F^- \lambda^- ilvG^- rfb-50 rph-1$

DSMZ, Hamburg (Germany)

One Shot TOP10 $F^- \Phi 80 lac Z \Delta M15 \Delta (mrr-hsdRMS-mcrBC) recA1 endA1$

 $mcrA \Delta lacX74 \ araD139 \ \Delta (ara-leu)7697 \ galU \ galK \ rpsL$

(Str^R) λ^- nupG

Invitrogen, Karlsruhe (Germany)

Origami(DE3) Δ (ara-leu)7697 Δ lacX74 Δ phoA Pvull phoR araD139 ahpC

galE galK rpsL F'[lac + lacI q pro] (DE3)gor522::Tn10 trxB

 (Kan^R, Str^R, Tet^R)

Novagen, Wisconsin (USA)

Rosetta(DE3) $F^- ompT \ hsdSB(r_R^-, m_R^-) \ gal \ dcm \ \lambda(DE3) \ pRARE \ (Cam^R)$

Novagen, Wisconsin (USA)

Rosetta(DE3) pLysS F^- ompT $hsdSB(r_B^-, m_B^-)$ gal dcm $\lambda(DE3)$ pLysSRARE

 (Cam^R)

Novagen, Wisconsin (USA)

T7 Express fhuA2 lacZ::T7 gene1 [lon] ompT gal sulA11 R(mcr-

73:: $miniTn10-Tet^S$)2 [dcm] $R(zgb-210::Tn10-Tet^S$) endA1

Δ(*mcrC-mrr*)114::*IS*10 NEB, Massachusetts (USA)

Agrobacterium tumefaciens

GV3101

chromosomal background: C58, marker gene: *rif*, tumor inducing plasmid (Ti-plasmid): cured, opine: nopaline Sylvestre Marillonet, IPB

3.1.4 Plasmids

Table 3.1.: Plasmids used in this work.

name	description
pACYC Duet-1	
pCDF Duet-1	
pET20b(+)	
pET28a(+)	
pET32a(+)	
pET41a(+)	
pQE30	
pUC19	

3.1.5 Oligonucleotides

Oligonucleotides and primers were ordered from Eurofins Genomics (Ebersberg, Germany). The purity grade was *high purity salt free* (HPSF).

Table 3.2.: Primers used in this work. Recognition sites for endonucleases are underlined.

name	sequence $(5'\rightarrow 3')$	cloning site
somt1	TTGAAGACAAAATGGCTTCTTCATTAAACAATGGCCG	BpiI
somt2	TTGAAGACAAGGACACCCCAAATACTGTGAGATCTTCC	BpiI
somt3	TTGAAGACAAGTCCTTAGGAACACCTTTCTGGGAC	BpiI
somt4	TT <u>GAAGAC</u> AAAAGCTCAAGGATAGATCTCAATAAGAGAC	BpiI

3.1.6 Software

All mathematical and statistical computations and graphics were done with the R software (versions 3.1.X, http://cran.r-project.org/). Visualizations of macromolecules were arranged using the PyMol Molecular Graphics System, version 1.7.0.0 (Schrödinger, New York (USA)).

Physicochemical calculations and calculations of different molecular descriptors were performed using Marvin Beans 15.4.13.0 (ChemAxon, Budapest (Hungary)) and Molecular Operating Environment 2008.10 (Chemical Computing Group, Montreal (Canada)).

3.2 Microbiology

3.3 Molecular Biology

Basic molecular biology methods like polymerase chain reaction (PCR), DNA restriction/ligation, DNA gel electrophoresis, preparation of competent cells and transformation were performed based on the protocols summarized by Sambrook and Russell [5].

Plasmid DNA was isolated using the QIAprep Spin Miniprep Kit (QIAGEN, Hilden (Germany)) according to the manufacturer's instructions.

In vitro site-directed mutatgenesis was set-up according to the protocol of the *QuikChange Site-Directed Mutagenesis* kit [6] offered by Agilent Technologies (Santa Clara, USA).

Nucleotide fragments obtained by PCR, restriction/ligation procedures or excision from electrophoresis gels were purified and concentrated using the *Nucleospin Gel and PCR Clean-up* kit provided by Machery-Nagel (Düren, Germany) according to the instructions provided by the manufacturer.

3.3.1 Golden Gate Cloning

The Golden Gate cloning procedure is a one-pot method, meaning the restriction digestion and ligation are carried out in the same reaction vessel at the same time [3, 1]. Consequently PCR-fragments, destination vector, restriction endonuclease and ligase are added together in this reaction. The methodology employs type II restriction enzymes, which together with proper design of the fragments allow for a ligation product lacking the original restriction sites.

For digestion/ligation reactions of fragments containing BpiI sites, 20 fmol of each fragment or vector, together with 5 U of BpiI and 5 U of T4 ligase were combined in a total volume of 15 μ l 1× ligase buffer. For fragments to be cloned via BsaI sites, BpiI in the above reaction was substituted by 5 U BsaI.

The reaction mixture was placed in a thermocycler and inbcubated at 37 $^{\circ}$ C for 2 min and 16 $^{\circ}$ C for 5 min. These two first steps were repeated 50 times over. Finally, the temperature was raised to 50 $^{\circ}$ C (5 min) and 80 $^{\circ}$ C (10 min) to inactivate the enzymes.

- 3.4 Protein biochemistry
- 3.5 Analytics

4 Evaluation of PFOMT towards the acceptance of long-chain SAM analogues

5 Enzymatic methylation of Noncatechols

testing the HPLC and again the HPLC.

Blöalala phenyl
propanoid and flavonoid O-methyl transferase (PFOMT) and
 $\mbox{\sc PFOMT}$

6 Development of an whole cell methyl transferase screening system

7 Acknowledgements

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetuer.

Appendix

A Figures




Figure A.1.: Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem.

B Tables

Table B.1.: Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem.

A	В	С	D	Е	F	G	Н	I
1	2	3	4	5	6	7	8	9
1	2	3	4	5	6	7	8	9
1	2	3	4	5	6	7	8	9

C Affidavit

I hereby declare that this document has been written only by the undersigned and
without any assistance from third parties. Furthermore, I confirm that no sources
have been used in the preparation of this document other than those indicated in
the thesis itself.
Date:, Location:, Signature:

Bibliography

- [1] Carola Engler, Romy Kandzia, and Sylvestre Marillonnet. "A one pot, one step, precision cloning method with high throughput capability". In: *PLoS ONE* 3.11 (2008).
- [2] Mwafaq Ibdah et al. "A novel Mg(2+)-dependent O-methyltransferase in the phenylpropanoid metabolism of Mesembryanthemum crystallinum." In: *The Journal of biological chemistry* 278.45 (Nov. 2003), pp. 43961–72.
- [3] Youichi Kondou et al. "cDNA Libraries". In: Methods 729 (2011), pp. 183-197.
- [4] Jakub G Kopycki et al. "Biochemical and structural analysis of substrate promiscuity in plant Mg2+-dependent O-methyltransferases." In: *Journal of molecular biology* 378.1 (Apr. 2008), pp. 154–64.
- [5] J. Sambrook and D.W. Russell. *Molecular Cloning: A Laboratory Manual.* 3rd ed. Cold Spring Harbor (NY, USA): Cold Spring Harbor Laboratory Press, 2001.
- [6] Agilent Technologies. QuikChange Site-Directed Mutagenesis Kit Instruction Manual. 2013.
- [7] Thomas Vogt. "Regiospecificity and kinetic properties of a plant natural product O-methyltransferase are determined by its N-terminal domain." In: *FEBS letters* 561.1-3 (Mar. 2004), pp. 159–62.

Acronyms

CD circulary dichroism. 10

FPLC fast protein liquid chromatography. 10

GC/MS gas chromatography coupled mass-spectrometry. 10

HPLC high-performance liquid chromatography. 10, 17

ITC Isothermal Titration Calorimetry. 10, 35

MTP micro-titer plate. 10, 35

NMR nuclear magnetic resonance. 10

PCR polymerase chain reaction. 13

PFOMT phenylpropanoid and flavonoid O-methyl transferase. 17, 35

Ti-plasmid tumor inducing plasmid. 12, 35

Glossary

Isothermal Titration Calorimetry (ITC) Fill in description here. 33

MTP Micro-titer plate. Small format rectangular plastic plate containing wells to allow for storage of multiple small samples or the containment multiple simultaneous reactions. Typical sizes include 24, 96 and 384-wells. 33

PFOMT Phenylpropanoid and flavonoid O-methyl transferase from *Mesembryan-themum crystallinum*, which was first described by Ibdah et al. in 2003 [2]. 33

Ti-plasmid Commonly found plasmids in *A. tumefaciens* and *A. rhizogenes* that confer virulence. 33