

Biotransformations from and to methylated flavonoids

Hpw all went to shit

Benjamin Weigel
Leibniz-Institute of Plant Biochemistry
Department of Bioorganic Chemistry
Weinberg 3
06120 Halle(Saale)
April 21, 2015

Advisor: Prof. Dr. Ludger A. Wessjohann
wessjohann@ipb-halle.de
+49 (345) 5582-1301

noch nicht bekannt

Contents

I Preface	1
<hr/>	
1 Abstracts	3
1.1 English Abstract	3
1.2 Deutsche Zusammenfassung	3
 II Thesis	 5
<hr/>	
2 Introduction	7
2.1 Natural products and secondary metabolites	7
2.1.1 General	7
2.1.2 Classes of natural products	7
2.2 Alkylating reactions in nature	7
2.2.1 Methylation	7
2.2.2 Prenylation	7
2.2.3 Glycosylation	8
2.3 Usage and expansion of natures reaction toolbox	8
2.3.1 Terpene synthases and elongases	8
2.3.2 Methyl transferases	8
2.3.3 Glycosyl transferases	8
2.3.4 Other important enzymes in biotech research	8
2.4 Conclusion	8
3 Material And Methods	9
3.1 Materials	9
3.1.1 Chemicals	9

3.1.2 Instruments	10
3.1.3 Bacterial strains	10
3.1.4 Plasmids	12
3.1.5 Oligonucleotides	12
3.1.6 Software	12
3.2 Microbiology	13
3.3 Molecular Biology	13
3.3.1 Golden Gate Cloning	13
3.4 Protein biochemistry	14
3.4.1 Production of recombinant protein	14
3.5 Analytics	14
3.5.1 In vitro determination of glucose	14
3.5.2 HPLC measurements	14
4 Evaluation of PFOMT towards the acceptance of long-chain SAM analogues	17
5 Enzymatic methylation of Non-catechols	19
6 Development of an whole cell methyl transferase screening system	21
7 Acknowledgements	23
 III Appendix	 25
<hr/>	
A Figures	27
B Tables	29
C Affidavit	31
Acronyms	35
Glossary	37

List of Figures

3.1 GOD assay	15
A.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem.	27

List of Tables

3.1 Plasmids used in this work.	12
3.2 Primers used in this work. Recognition sites for endonucleases are underlined.	12
B.1 Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem.	29



Preface

1 Abstracts

1.1 English Abstract

Test Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

1.2 Deutsche Zusammenfassung

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.



Thesis

2 Introduction

Some introductory text

2.1 Natural products and secondary metabolites

2.1.1 General

2.1.2 Classes of natural products

Terpenoids and Steroids

... here is some text

Polyketides and non-ribosomal peptides

... here is some text

Alkaloids

... here is some text

Phenylpropanoids

... here is some text

2.2 Alkylating reactions in nature

2.2.1 Methylation

2.2.2 Prenylation

2.2.3 Glycosylation

2.3 Usage and expansion of nature's reaction toolbox

2.3.1 Terpene synthases and elongases

2.3.2 Methyl transferases

2.3.3 Glycosyl transferases

2.3.4 Other important enzymes in biotech research

BMVOs

Esterases/Lipases

Oxidases

Lyases

Transaminases

2.4 Conclusion

3 Material And Methods

3.1 Materials

3.1.1 Chemicals

Enzymes and buffers used for molecular cloning were obtained from Thermo Scientific (Darmstadt, Germany), unless otherwise noted. Flavonoid HPLC standards were purchased from Extrasynthese (Genay, France). Deuterated solvents were aquired from Deutero GmbH (Kastellaun, Germany). Solvents, purchased from VWR (Poole, England), were distilled in-house before use.

All other chemicals were obtained from either Sigma-Aldrich (Steinheim, Germany), Applichem (Darmstadt, Germany), Carl Roth (Karlsruhe, Germany) or Merck (Darmstadt, Germany).

3.1.2 Instruments

circular dichroism (CD)-spectrometer	Jasco J-815 (Eaton, USA)
electrophoresis (horizontal)	Biometra Compact XS/S (Göttingen, Germany)
electrophoresis (vertical)	Biometra Compact M (Göttingen, Germany)
	Biometra Minigel-Twin (Göttingen, Germany)
fast protein liquid chromatography (FPLC)	ÄKTA purifier (GE Healthcare, Freiburg, Germany)
gas chromatography coupled mass-spectrometry (GC/MS)	GC-MS-QP2010 Ultra (Shimadzu, Duisburg, Germany)
high-performance liquid chromatography (HPLC)	VWR ???
Isothermal Titration Calorimetry (ITC)	MicroCal iTC200 (Malvern, Worcestershire, UK)
micro-titer plate (MTP)	???
MTP-reader	SpectraMax M5 (Molecular Devices, Biberach, Germany)
nuclear magnetic resonance (NMR)-spectrometer	Varian Unity 400 (Agilent, Böblingen, Germany)
	Varian VNMRs 600 (Agilent, Böblingen, Germany)
photospectrometer	Eppendorf Biophotometer Plus (Hamburg, Germany)
	JASCO V-560 (Eaton, USA)
	Nanodrop ???
centrifuges	Eppendorf 5424 (Hamburg, Germany)
	Hettich Mikro 120 (Kirchlengern, Germany)
	Beckman Avanti J-E (Krefeld, Germany)
centrifuge rotors	Beckman JA-10, JA-16.250, JS-4.3 (Krefeld, Germany)

3.1.3 Bacterial strains

E.coli

BL21(DE3)	F ⁻ <i>ompT hsdSB(r_B⁻,m_B⁻) gal dcm λ</i> (DE3) Invitrogen, Karlsruhe
C41(DE3)	F ⁻ <i>ompT hsdSB(r_B⁻,m_B⁻) gal dcm λ</i> (DE3) Lucigen, Wisconsin (USA)
C43(DE3)	F ⁻ <i>ompT hsdSB(r_B⁻,m_B⁻) gal dcm λ</i> (DE3) Lucigen, Wisconsin (USA)
DH5α	F ⁻ Φ80 <i>lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17(r_K⁻m_K⁺) phoA supE44 λ⁻ thi-1 gyrA96 relA1</i> Invitrogen, Karlsruhe (Germany)
JM110	<i>rpsL thr leu thi lacY galK galT ara tonA tsx dam dcm glnV44 Δ(lac-proAB) e14- [F' traD36 proAB⁺ lacI^q lacZΔM15] hsdR17(r_K⁻m_K⁺)</i> Martin-Luther-University Halle-Wittenberg
JW1593 (BW25113 derivative)	<i>rrnB ΔlacZ4787 HsdR514 Δ(araBAD)568 rph-1 ΔydgG</i> (Kan ^R) Keio Collection, National Institute of Genetics (Japan)
MG1655	F ⁻ λ ⁻ <i>ilvG⁻ rfb-50 rph-1</i> DSMZ, Hamburg (Germany)
One Shot TOP10	F ⁻ Φ80 <i>lacZΔM15 Δ(mrr-hsdRMS-mcrBC) recA1 endA1 mcrA ΔlacX74 araD139 Δ(ara-leu)7697 galU galK rpsL</i> (Str ^R) λ ⁻ <i>nupG</i> Invitrogen, Karlsruhe (Germany)
Origami(DE3)	Δ(<i>ara-leu</i>)7697 Δ <i>lacX74 ΔphoA Pvull phoR araD139 ahpC galE galK rpsL</i> F' [<i>lac + lacI q pro</i>] (DE3)gor522::Tn10 <i>trxB</i> (Kan ^R , Str ^R , Tet ^R) Novagen, Wisconsin (USA)
Rosetta(DE3)	F ⁻ <i>ompT hsdSB(r_B⁻,m_B⁻) gal dcm λ</i> (DE3) pRARE (Cam ^R) Novagen, Wisconsin (USA)
Rosetta(DE3) pLysS	F ⁻ <i>ompT hsdSB(r_B⁻,m_B⁻) gal dcm λ</i> (DE3) pLysSRARE (Cam ^R) Novagen, Wisconsin (USA)
T7 Express	<i>fhuA2 lacZ::T7 gene1 [lon] ompT gal sulA11 R(mcr-73::miniTn10-Tet^S)2 [dcm] R(zgb-210::Tn10-Tet^S) endA1 Δ(mcrC-mrr)114::IS10</i> NEB, Massachusetts (USA)

Agrobacterium tumefaciens

GV3101 chromosomal background: C58, marker gene: *rif*, tumor inducing plasmid (Ti-plasmid): cured, opine: nopaline
Sylvestre Marillonet, IPB

3.1.4 Plasmids**Table 3.1.:** Plasmids used in this work.

name	description
pACYC Duet-1	..
pCDF Duet-1	..
pET20b(+)	...
pET28a(+)	..
pET32a(+)	..
pET41a(+)	..
pQE30	..
pUC19	..

3.1.5 Oligonucleotides

Oligonucleotides and primers were ordered from Eurofins Genomics (Ebersberg, Germany). The purity grade was *high purity salt free* (HPSF).

Table 3.2.: Primers used in this work. Recognition sites for endonucleases are underlined.

name	sequence (5'→3')	cloning site
somt1	TTGAAGACA <u>AAA</u> ATGGCTTCTTCATTAAACAATGGCCG	BpI
somt2	TTGAAGACAAGGACACCCCAAATACTGTGAGATCTTCC	BpI
somt3	TTGAAGACAAGTCCTTAGGAACACCTTTCTGGGAC	BpI
somt4	TTGAAGACA <u>AAA</u> AGCTCAAGGATAGATCTCAATAAGAGAC	BpI

3.1.6 Software

All mathematical and statistical computations and graphics were done with the R software (versions 3.1.X, <http://cran.r-project.org/>). Visualizations of macromolecules were arranged using the PyMol Molecular Graphics System, version 1.7.0.0 (Schrödinger, New York (USA)).

Physicochemical calculations and calculations of different molecular descriptors were performed using Marvin Beans 15.4.13.0 (ChemAxon, Budapest (Hungary)) and Molecular Operating Environment 2008.10 (Chemical Computing Group, Montreal (Canada)).

3.2 Microbiology

3.3 Molecular Biology

Basic molecular biology methods like polymerase chain reaction (PCR), DNA restriction/ligation, DNA gel electrophoresis, preparation of competent cells and transformation were performed based on the protocols summarized by Sambrook and Russell [5].

Plasmid DNA was isolated using the QIAprep Spin Miniprep Kit (QIAGEN, Hilden (Germany)) according to the manufacturer's instructions.

In vitro site-directed mutagenesis was set-up according to the protocol of the *QuikChange Site-Directed Mutagenesis* kit [7] offered by Agilent Technologies (Santa Clara, USA).

Nucleotide fragments obtained by PCR, restriction/ligation procedures or excision from electrophoresis gels were purified and concentrated using the *Nucleospin Gel and PCR Clean-up* kit provided by Machery-Nagel (Düren, Germany) according to the instructions provided by the manufacturer.

3.3.1 Golden Gate Cloning

The Golden Gate cloning procedure is a one-pot method, meaning the restriction digestion and ligation are carried out in the same reaction vessel at the same time [3, 1]. Consequently PCR-fragments, destination vector, restriction endonuclease and ligase are added together in this reaction. The methodology employs type II restriction enzymes, which together with proper design of the fragments allow for a ligation product lacking the original restriction sites.

For digestion/ligation reactions of fragments containing BpI sites, 20 fmol of each fragment or vector, together with 5 U of BpI and 5 U of T4 ligase were combined in a total volume of 15 µl 1× ligase buffer. For fragments to be cloned via BsaI sites, BpI in the above reaction was substituted by 5 U BsaI.

The reaction mixture was placed in a thermocycler and incubated at 37 °C for 2 min and 16 °C for 5 min. These two first steps were repeated 50 times over. Finally, the temperature was raised to 50 °C (5 min) and 80 °C (10 min) to inactivate the enzymes.

3.4 Protein biochemistry

3.4.1 Production of recombinant protein

Heterologous production of PFOMT

3.5 Analytics

3.5.1 In vitro determination of glucose

The glucose concentration in clarified, aqueous samples was determined by a modified version of the glucose assay kit procedure provided by Sigma-Aldrich [6]. Glucose oxidase (GOD) oxidizes D-glucose to gluconic acid, whereby hydrogen peroxide is produced. The hydrogen peroxide can be detected and quantified by horseradish peroxidase (HRP), which reduces the produced H_2O_2 and thereby oxidizes its chromogenic substrate *o*-dianisidine. The oxidized form of *o*-dianisidine can then be measured photospectrometrically.

The methodology employs a coupled photospectrometric assay using GOD and HRP with *o*-dianisidine as reporter substrate. The assay was prepared in MTP-format. A reaction solution containing 12.5 U/ml GOD, 2.5 U/ml HRP and 0.125 mg ml^{-1} *o*-dianisidine dihydrochloride in 50 mM sodium acetate pH 5.1 was prepared.

Sample solutions from culture supernatants were typically diluted in 9 volumes of water. The reaction was started, by adding 50 μl reaction solution to 25 μl of sample and was incubated at 37 °C and 200 rpm for 30 min in a shaking incubator. 50 μl 6 M sulfuric acid was added to stop the reaction and achieve maximum color development (full oxidation of any *o*-dianisidine charge transfer complexes).

The developed pink color was measured at 540 nm in a MTP-reader. A calibration curve of a standard D-glucose solutions (0 to $100 \mu\text{g ml}^{-1}$), that was always part of the experiments, was used to quantify the sample measurements.

3.5.2 HPLC measurements




Figure 3.1.: GOD assay

4 Evaluation of PFOMT towards the acceptance of long-chain SAM analogues

5 Enzymatic methylation of Non-catechols

testing the HPLC and again the HPLC.


Blöälala phenylpropanoid and flavonoid O-methyl transferase (PFOMT) and PFOMT

6 Development of an whole cell methyl transferase screening system

7 Acknowledgements

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.



Appendix

A Figures




Figure A.1.: Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem.

B Tables

Table B.1.: Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aenean commodo ligula eget dolor. Aenean massa. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Donec quam felis, ultricies nec, pellentesque eu, pretium quis, sem.

A	B	C	D	E	F	G	H	I
1	2	3	4	5	6	7	8	9
1	2	3	4	5	6	7	8	9
1	2	3	4	5	6	7	8	9

C Affidavit

I hereby declare that this document has been written only by the undersigned and without any assistance from third parties. Furthermore, I confirm that no sources have been used in the preparation of this document other than those indicated in the thesis itself.

Date:....., Location:....., Signature:.....

Bibliography

- [1] Carola Engler, Romy Kandzia, and Sylvestre Marillonnet. “A one pot, one step, precision cloning method with high throughput capability.” In: *PloS one* 3.11 (Jan. 2008), e3647.
- [2] Mwafaq Ibdah et al. “A novel Mg(2+)-dependent O-methyltransferase in the phenylpropanoid metabolism of *Mesembryanthemum crystallinum*.” In: *The Journal of biological chemistry* 278.45 (Nov. 2003), pp. 43961–72.
- [3] Youichi Kondou et al. “cDNA Libraries”. In: *Methods* 729 (2011), pp. 183–197.
- [4] Jakub G Kopycki et al. “Biochemical and structural analysis of substrate promiscuity in plant Mg²⁺-dependent O-methyltransferases.” In: *Journal of molecular biology* 378.1 (Apr. 2008), pp. 154–64.
- [5] J Sambrook and D W Russell. *Molecular Cloning: A Laboratory Manual*. 3rd ed. Cold Spring Harbor (NY, USA): Cold Spring Harbor Laboratory Press, 2001.
- [6] Sigma-Aldrich. *Glucose (GO) Assay Kit - Technical Bulletin GAGO-20*.
- [7] Agilent Technologies. *QuikChange Site-Directed Mutagenesis Kit Instruction Manual*. 2013.
- [8] Thomas Vogt. “Regiospecificity and kinetic properties of a plant natural product O-methyltransferase are determined by its N-terminal domain.” In: *FEBS letters* 561.1-3 (Mar. 2004), pp. 159–62.

Acronyms

CD circular dichroism. 10

FPLC fast protein liquid chromatography. 10

GC/MS gas chromatography coupled mass-spectrometry. 10

GOD glucose oxidase. 14, 37

HPLC high-performance liquid chromatography. 10, 19

HRP horseradish peroxidase. 14

ITC Isothermal Titration Calorimetry. 10, 37

MTP micro-titer plate. 10, 14, 37

NMR nuclear magnetic resonance. 10

PCR polymerase chain reaction. 13

PFOMT phenylpropanoid and flavonoid O-methyl transferase. 19, 37

Ti-plasmid tumor inducing plasmid. 12, 37

Glossary

GOD Glucose oxidase is an enzyme.... . 35

Isothermal Titration Calorimetry (ITC) Fill in description here. 35

MTP Micro-titer plate. Small format rectangular plastic plate containing wells to allow for storage of multiple small samples or the containment multiple simultaneous reactions. Typical sizes include 24, 96 and 384-wells. 35

PFOMT Phenylpropanoid and flavonoid O-methyl transferase from *Mesembryanthemum crystallinum*, which was first described by Ibdah et al. in 2003 [Ibdah2003b]. 35

Ti-plasmid Commonly found plasmids in *A. tumefaciens* and *A. rhizogenes* that confer virulence. 35