

Parabolic PDEs

for a PDE $Au_{xx} + 2Bu_{xy} + Cu_{yy} + Du_x + Eu_y + F = 0$,
Parabolic PDEs satisfy $B^2 - AC = 0$

In astrophysics, conduction / diffusion are the standard examples:

$$\vec{F} = -k \vec{\nabla} \phi$$

\uparrow flux of temp. or concentration \uparrow coefficient (may not be constant) \longleftarrow temp. or concentration

Which can be rewritten as a conservation law:

$$\phi_t + \vec{\nabla} \cdot \vec{F} = 0, \text{ which turns into a diffusion eqn:}$$


$$\phi_t = -\vec{\nabla} \cdot \vec{F} = \vec{\nabla} \cdot (k \vec{\nabla} \phi) \quad (\text{general form})$$

In 1D w/ constant k ,

$$\frac{\partial \phi}{\partial t} = k \frac{\partial^2 \phi}{\partial x^2}$$

[both diffusion and conduction in astrophysics use non-constant k , but it adds to the algebra but conceptually stays the same],

Explicit expression:



$$\frac{\phi_i^{n+1} - \phi_i^n}{\Delta t} = k \frac{\phi_{i+1}^n - 2\phi_i^n + \phi_{i-1}^n}{\Delta x^2}$$

[reminder: $n, n+1 = \text{time}$
 $i, i \pm 1 = \text{space}$]

shuffling around terms:

$$\phi_i^{n+1} = \phi_i^n + \boxed{\frac{k \Delta t}{\Delta x^2}} (\phi_{i+1}^n - 2\phi_i^n + \phi_{i-1}^n)$$

↳ gives a clue to stability criterion

stability analysis: to not be numerically unstable,

$$\Delta t < \frac{1}{2} \frac{\Delta x^2}{k}$$

or
$$\Delta t = \frac{C}{2} \frac{\Delta x^2}{k}, \quad C < 1$$

very restrictive due to Δx^2 ! even worse in plasmas for conduction, since $k \propto T^{5/2}$, so

$$\Delta t \propto \frac{\Delta x^2}{T^{5/2}} \leftarrow \text{ouch!}$$

→ lends itself to an implicit solution!

Implicit discretization: (stable for all $c > 0$, but
(the "Implicit Euler method") not necessarily accurate)

$$\frac{\phi_i^{n+1} - \phi_i^n}{\Delta t} = k \frac{\phi_{i+1}^{n+1} - 2\phi_i^{n+1} + \phi_{i-1}^{n+1}}{\Delta x^2}$$

Define $\alpha \equiv \frac{k\Delta t}{\Delta x^2}$ and reshuffle:

$$\phi_i^{n+1} + 2\alpha\phi_i^{n+1} - \alpha\phi_{i+1}^{n+1} - \alpha\phi_{i-1}^{n+1} = \phi_i^n$$

or, rearranging:

$$-\alpha\phi_{i-1}^{n+1} + (1 + 2\alpha)\phi_i^{n+1} - \alpha\phi_{i+1}^{n+1} = \phi_i^n$$

which starts to look a lot like $Ax = b$

$$(\text{or } A\phi^{n+1} = \phi^n)$$

Which ends up looking like:

$$\begin{pmatrix} (1+2\alpha) & -\alpha & 0 & 0 & \dots \\ -\alpha & (1+2\alpha) & -\alpha & 0 & \dots \\ 0 & -\alpha & (1+2\alpha) & -\alpha & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \begin{pmatrix} \phi_{i-1}^{n+1} \\ \phi_i^{n+1} \\ \phi_{i+1}^{n+1} \\ \vdots \end{pmatrix} = \begin{pmatrix} \phi_{i-1}^n \\ \phi_i^n \\ \phi_{i+1}^n \\ \vdots \end{pmatrix}$$

This is a tridiagonal
matrix and is easily
solvable w/ libraries!

but, end points need to be modified for boundary conditions!
(see Zingales 10.19 and following)

Implicit Euler is only 1st order in time, how can we do better?

→ use time-centered $\nabla^2 \phi$ to solve! ("Crank-Nicholson method")

so, $\phi_t = k \phi_{xx}$ becomes:

$$\frac{\phi_i^{n+1} - \phi_i^n}{\Delta t} = k \nabla^2 \phi_i^{n+\frac{1}{2}} = k \cdot \frac{1}{2} (\nabla^2 \phi_i^{n+1} + \nabla^2 \phi_i^n)$$

averaging in time to get
2nd order!

turns into:

$$\phi_i^{n+1} - \phi_i^n = \frac{k}{2} \left[\underbrace{\frac{\phi_{i+1}^{n+1} - 2\phi_i^{n+1} + \phi_{i-1}^{n+1}}{\Delta x^2}}_{\text{time } n+1} + \underbrace{\frac{\phi_{i+1}^n - 2\phi_i^n + \phi_{i-1}^n}{\Delta x^2}}_{\text{time } n} \right]$$

Define $\alpha \equiv \frac{k\Delta t}{\Delta x^2}$ as before, group $n+1$ and n terms:

$$\underbrace{\phi_i^{n+1} - \frac{\alpha}{2} (\phi_{i+1}^{n+1} - 2\phi_i^{n+1} + \phi_{i-1}^{n+1})}_{\text{time } n+1} = \underbrace{\phi_i^n + \frac{\alpha}{2} (\phi_{i+1}^n - 2\phi_i^n + \phi_{i-1}^n)}_{\text{time } n}$$

doing some shuffling:

$$\underbrace{-\frac{\alpha}{2}\phi_{i-1}^{n+1} + (1+\alpha)\phi_i^{n+1} - \frac{\alpha}{2}\phi_{i+1}^{n+1}}_{\text{tri-diagonal matrix elements for } Ax=b} = \underbrace{\phi_i^n + \frac{\alpha}{2} (\phi_{i+1}^n - 2\phi_i^n + \phi_{i-1}^n)}_{= k \frac{\Delta t}{2} \nabla^2 \phi_i^n}$$

tri-diagonal matrix elements
for $Ax=b$

looks complicated, but
we know all of
this info already!