

Part I

SUMMARY of L'Hôpital's Rule:

Summary

February 21, 2026

Indeterminate Forms - Part I

- (1) $\frac{0}{0}$ refers to a limit of the form $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$, where $\lim_{x \rightarrow a} f(x) = 0$ and $\lim_{x \rightarrow a} g(x) = 0$
- (2) $\frac{\infty}{\infty}$ refers to a limit of the form $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$, where $\lim_{x \rightarrow a} f(x) = \pm\infty$ and $\lim_{x \rightarrow a} g(x) = \pm\infty$. (The signs do not need to match.)

How do we determine whether the limit of this form exists and, if it does, how do we evaluate the limit? We can apply L'Hôpital's Rule to limits of the form $\frac{0}{0}$ or $\frac{\infty}{\infty}$.

L'Hôpital's Rule (LHR)

Let $f(x)$ and $g(x)$ be functions that are differentiable near a .

If $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ and $g'(x) \neq 0$ for all x near a , then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)},$$

provided that $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$ exists.

L'Hôpital's rule applies also when $\lim_{x \rightarrow a} f(x) = \pm\infty$ and $\lim_{x \rightarrow a} g(x) = \pm\infty$.

Indeterminate Forms - Part II

- (3) $0 \cdot \infty$ refers to a limit of the form $\lim_{x \rightarrow a} f(x) \cdot g(x)$, where $\lim_{x \rightarrow a} f(x) = 0$, and $\lim_{x \rightarrow a} g(x) = \pm\infty$
- (4) $\infty - \infty$ refers to a limit of the form $\lim_{x \rightarrow a} (f(x) - g(x))$, where $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = \infty$

How do we determine whether the limit of this form exists and, if it does, how do we evaluate the limit? By performing convenient algebraic operations, we show that the limit is equivalent to a limit of the form $\frac{0}{0}$ or $\frac{\infty}{\infty}$, so that we can apply the LHR.

Indeterminate Forms - Part III

(5) 1^∞ refers to a limit of the form $\lim_{x \rightarrow a} f(x)^{g(x)}$, where $\lim_{x \rightarrow a} f(x) = 1$, and $\lim_{x \rightarrow a} g(x) = \infty$

(6) 0^0 refers to a limit of the form $\lim_{x \rightarrow a} f(x)^{g(x)}$, where $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$

(7) ∞^0 refers to a limit of the form $\lim_{x \rightarrow a} f(x)^{g(x)}$, where $\lim_{x \rightarrow a} f(x) = \infty$, and $\lim_{x \rightarrow a} g(x) = 0$

How do we determine whether the limit of this form exists and, if it does, how do we evaluate the limit? By performing the following steps

$$\lim_{x \rightarrow a} f(x)^{g(x)} = \lim_{x \rightarrow a} [e^{\ln(f(x))}]^{g(x)} = \lim_{x \rightarrow a} e^{g(x) \ln(f(x))} = e^{\lim_{x \rightarrow a} g(x) \ln(f(x))}$$

we end up with the limit of simpler form (in the exponent).

Part II

Recitation Questions

Problem 1

February 21, 2026

Problem 1 State the form of the limit. Determine whether the form is determinate or indeterminate. Evaluate each limit.

$$(a) \lim_{x \rightarrow 0} \frac{\sin(x) - \cos(x) + 1}{x^2 - x}$$

Explanation. The form is $\frac{0}{0}$; indeterminate. we can apply L.R.

$$\lim_{x \rightarrow 0} \frac{\sin(x) - \cos(x) + 1}{x^2 - x} = \lim_{x \rightarrow 0} \frac{\cos(x) + \sin(x)}{2x - 1} = \frac{\cos(0) + \sin(0)}{2(0) - 1} = \frac{1}{-1} = -1$$

$$(b) \lim_{x \rightarrow 0} \frac{e^x - 1 - x}{x^2}$$

Explanation. The form is $\frac{0}{0}$; indeterminate.

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x}{x^2} = \lim_{x \rightarrow 0} \frac{e^x - 1}{2x}, \text{ by L.R.}$$

The last limit also has the form $\frac{0}{0}$, so, we can apply L.R. again.

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x}{x^2} = \lim_{x \rightarrow 0} \frac{e^x - 1}{2x} = \lim_{x \rightarrow 0} \frac{e^x}{2} = \frac{1}{2}$$

$$(c) \lim_{x \rightarrow \infty} \frac{e^x}{x^4}$$

Explanation. The form is $\frac{\infty}{\infty}$; indeterminate.

$$\lim_{x \rightarrow \infty} \frac{e^x}{x^4} = \lim_{x \rightarrow \infty} \frac{e^x}{4x^3}, \text{ by L.R.}$$

The last limit also has the form $\frac{\infty}{\infty}$. It turns out that we can apply L.R. again, and again, and again:

$$\lim_{x \rightarrow \infty} \frac{e^x}{x^4} = \lim_{x \rightarrow \infty} \frac{e^x}{4x^3} = \lim_{x \rightarrow \infty} \frac{e^x}{12x^2} = \lim_{x \rightarrow \infty} \frac{e^x}{24x} = \lim_{x \rightarrow \infty} \frac{e^x}{24} = \infty$$

This result shows that the function e^x grows much faster than x^4 , as x goes to ∞ .

Problem 2

February 21, 2026

Problem 2 State the form of the limit. Determine whether the form is determinate or indeterminate. Evaluate each limit.

(a) $\lim_{x \rightarrow \infty} (\ln(1 + e^{-x}))^x$

Explanation. Since $\lim_{x \rightarrow \infty} (1 + e^{-x}) = 1 + 0 = 1$ and $\ln(1) = 0$, this limit is of the form 0^∞ . This is a determinate form. $\lim_{x \rightarrow \infty} (\ln(1 + e^{-x}))^x = 0$.

(b) $\lim_{x \rightarrow \infty} \left(\frac{1}{x} + 1 \right)^{\frac{1}{x}}$

Explanation. This limit is of the form 1^0 , which is a determinate form.

Thus, $\lim_{x \rightarrow \infty} \left(\frac{1}{x} + 1 \right)^{\frac{1}{x}} = 1$

(c) $\lim_{x \rightarrow \infty} \left(\frac{\arctan(x)}{x} \right)$

Explanation. Since $\lim_{x \rightarrow \infty} \arctan(x) = \frac{\pi}{2}$, this limit is of the form $\frac{\#}{\infty}$, which is a determinate form.

Thus, $\lim_{x \rightarrow \infty} \left(\frac{\arctan(x)}{x} \right) = 0$

(d) $\lim_{x \rightarrow \infty} (x - \ln(x))$

Explanation. This limit is of the form $\infty - \infty$, which is an indeterminate form. We can rewrite this as:

$$\lim_{x \rightarrow \infty} (x - \ln(x)) = \lim_{x \rightarrow \infty} \left(x \left(1 - \frac{\ln(x)}{x} \right) \right)$$

We can see that $\lim_{x \rightarrow \infty} \left(\frac{\ln(x)}{x} \right)$ is of the form $\frac{\infty}{\infty}$, so we can use L'Hopital's Rule.

$$\xrightarrow{L.R.} \lim_{x \rightarrow \infty} \left(\frac{1/x}{1} \right) = \lim_{x \rightarrow \infty} \left(\frac{1}{x} \right) = 0$$

Now we have:

$$\lim_{x \rightarrow \infty} \left(x \left(1 - \frac{\ln(x)}{x} \right) \right)$$

This limit has the form $\infty \cdot 1$. This is a determinate form, and, therefore,

$$\lim_{x \rightarrow \infty} (x - \ln(x)) = \infty$$

(e) $\lim_{x \rightarrow \infty} \left(x \ln \left(\frac{1}{x} \right) \right)$

Explanation. As x approaches ∞ , $\frac{1}{x}$ approaches 0 from the right. So

$$\lim_{x \rightarrow \infty} \ln \left(\frac{1}{x} \right) = -\infty$$

Therefore, the limit in question is of the form $\infty \cdot -\infty$, which is a determinate form.

Thus,

$$\lim_{x \rightarrow \infty} \left(x \ln \left(\frac{1}{x} \right) \right) = -\infty$$

(f) $\lim_{x \rightarrow 0^+} (\sin(x) \cot(x))$

Explanation. Since $\lim_{x \rightarrow 0^+} \cot(x) = \infty$, this limit is of the form $0 \cdot \infty$. This is an indeterminate form.

Note: $\cot(x) = \frac{\cos(x)}{\sin(x)}$. So

$$\lim_{x \rightarrow 0^+} (\sin(x) \cot(x)) = \lim_{x \rightarrow 0^+} \cos(x) = 1.$$

Problem 3

February 21, 2026

Problem 3 Circle the correct answer in each part:

(a) Consider the limit $\lim_{x \rightarrow 0} (\cos(x))^{\sin(x)}$.

(i) Evaluate the limit.

i. the limit DNE

ii. e

iii. 1

iv. ∞

v. $-\infty$

vi. 0

vii. none of the previous answers is correct

Explanation. The correct choice is (iii).

Evaluation of limit:

$$\lim_{x \rightarrow 0} \underbrace{(\cos(x))^{\sin(x)}}_{\text{form } 1^0} = 1$$

(ii) What Limit Law, rule or technique did you use to find this limit?

i. The Squeeze Theorem;

ii. L'Hôpital's Rule;

iii. The Product Law;

iv. evaluated the function at $x = 0$, since the function is continuous at $x = 0$;

v. none of the previous answers is correct

Explanation. The correct choice is (iv).

(b) Evaluate the limit $\lim_{x \rightarrow 4^-} \frac{\ln(x)}{x - 4}$.

(i) the limit DNE

(ii) e

(iii) 1

(iv) ∞

(v) $-\infty$

(vi) 0

(vii) none of the previous answers is correct

Explanation. The correct choice is (v).

Evaluation of limit: $\lim_{x \rightarrow 4^-} \frac{\ln(x)}{x - 4} = -\infty$,

since the limit is of the form $\frac{\#}{0}$, and since the numerator is positive and the denominator negative.

Problem 3

(c) Evaluate the limit $\lim_{x \rightarrow \infty} \frac{\ln(x)}{x-4}$.

- (i) the limit DNE
- (ii) e
- (iii) 1
- (iv) ∞
- (v) $-\infty$
- (vi) 0
- (vii) none of the previous answers is correct

Explanation. The correct choice is (vi).

Evaluation of limit:

$$\lim_{x \rightarrow \infty} \underbrace{\frac{\ln(x)}{x-4}}_{\text{form } \infty/\infty} \stackrel{L.H.}{=} \lim_{x \rightarrow \infty} \frac{1/x}{1} = 0$$

(d) Consider the limit

$$\lim_{h \rightarrow 0} \frac{(2+h)^3 - 8}{h} = f'(2).$$

Determine the function f .

- (i) such a function DNE;
- (ii) $f(x) = x^3$;
- (iii) $f(x) = (2+x)^3$;
- (iv) $f(x) = \frac{(2+x)^3}{x}$;
- (v) none of the previous answers is correct

Explanation. The correct choice is (ii).

(e) Consider the limit: $\lim_{x \rightarrow 0^+} \left(\frac{\sin(x)}{x} \right)^{|\ln(x)|}$.

Determine the form of this limit.

- (i) $\frac{0}{0}$;
- (ii) $\frac{\infty}{\infty}$;
- (iii) 1^0 ;

Problem 3

- (iv) 0^0 ;
- (v) 1^∞ ;
- (vi) ∞^∞ ;
- (vii) *none of the previous answers is correct*

Explanation. The correct choice is (v).

Problem 4

February 21, 2026

Problem 4 Determine the following limits. Use L'Hôpital's Rule if applicable.

(a) $\lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}}$

Explanation. This limit is of the form: $\frac{\infty}{\infty}$. L.R. is applicable, but we don't need to apply it.

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 + 1}} &= \lim_{x \rightarrow \infty} \frac{x}{\sqrt{x^2 \left(1 + \frac{1}{x^2}\right)}} \\ &= \lim_{x \rightarrow \infty} \frac{x}{|x| \sqrt{1 + \frac{1}{x^2}}} \\ &= \lim_{x \rightarrow \infty} \frac{x}{x \sqrt{1 + \frac{1}{x^2}}} \\ &= \lim_{x \rightarrow \infty} \frac{1}{\sqrt{1 + \frac{1}{x^2}}} \\ &= \frac{1}{\sqrt{1 + 0}} = 1 \end{aligned}$$

(b) $\lim_{x \rightarrow -\infty} x^2 e^x$

Explanation. This limit is of the form: $\infty \cdot 0$

$$\begin{aligned} \lim_{x \rightarrow -\infty} x^2 e^x &= \lim_{x \rightarrow -\infty} \frac{x^2}{e^{-x}} \left(\text{of the form } \frac{\infty}{\infty} \right) \\ &\stackrel{L.R.}{=} \lim_{x \rightarrow -\infty} \frac{2x}{-e^{-x}} \left(\text{of the form } \frac{\infty}{\infty} \right) \\ &\stackrel{L.R.}{=} \lim_{x \rightarrow -\infty} \frac{2}{e^{-x}} \\ &= 0 \end{aligned}$$

where "L.R." above an equals sign means that that equality is due to "L'Hôpital's Rule".

(c) $\lim_{x \rightarrow \infty} x^{\frac{1}{x}}$

Explanation. This limit is of the form: ∞^0

$$\begin{aligned}\lim_{x \rightarrow \infty} x^{\frac{1}{x}} &= \lim_{x \rightarrow \infty} e^{\ln\left(x^{\frac{1}{x}}\right)} \\ &= \lim_{x \rightarrow \infty} e^{\frac{1}{x} \ln x} \\ &= \lim_{x \rightarrow \infty} \frac{\ln x}{x} \quad \left(\text{limit is of the form } \frac{\infty}{\infty}\right) \\ &\stackrel{L.R.}{=} \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{1} \\ &= \lim_{x \rightarrow \infty} \frac{1}{x} \\ &= e^0 = 1\end{aligned}$$

(d) $\lim_{x \rightarrow \infty} \left(1 + \frac{2}{x}\right)^x$

Explanation. This limit is of the form: 1^∞

$$\begin{aligned}\lim_{x \rightarrow \infty} \left(1 + \frac{2}{x}\right)^x &= \lim_{x \rightarrow \infty} e^{\ln\left(\left(1 + \frac{2}{x}\right)^x\right)} \\ &= \lim_{x \rightarrow \infty} e^{x \ln\left(1 + \frac{2}{x}\right)} \\ &= \lim_{x \rightarrow \infty} \frac{\ln\left(1 + \frac{2}{x}\right)}{1/x} \quad \left(\text{limit is of the form } \frac{0}{0}\right) \\ &\stackrel{L.R.}{=} \lim_{x \rightarrow \infty} \frac{-2x^{-2} \cdot \frac{1}{(1+2/x)}}{-x^{-2}} \\ &= \lim_{x \rightarrow \infty} \left(2 \left(\frac{1}{1 + \frac{2}{x}}\right)\right) \\ &= e^2\end{aligned}$$

(e) $\lim_{\theta \rightarrow 0^+} (\sin(\theta))^{\tan(\theta)}$

Explanation. This limit is of the form: 0^0

Problem 4

$$\begin{aligned}
 \lim_{\theta \rightarrow 0^+} (\sin(\theta))^{\tan \theta} &= \lim_{\theta \rightarrow 0^+} e^{\ln((\sin(\theta))^{\tan(\theta)})} \\
 &= \lim_{\theta \rightarrow 0^+} e^{\tan(\theta) \ln(\sin(\theta))} \\
 &= e^{\lim_{\theta \rightarrow 0^+} \frac{\ln(\sin(\theta))}{\cot(\theta)}} \quad \left(\text{limit is of the form } \frac{\infty}{\infty} \right) \\
 &\stackrel{L.R.}{=} e^{\lim_{\theta \rightarrow 0^+} \frac{\cos(\theta) \cdot \frac{1}{\sin(\theta)}}{-\csc^2(\theta)}} \\
 &= e^{\lim_{\theta \rightarrow 0^+} \left(\frac{\cos(\theta)}{\sin(\theta)} \cdot \frac{-\sin^2(\theta)}{1} \right)} \\
 &= e^{\lim_{\theta \rightarrow 0^+} (-\cos(\theta) \cdot \sin(\theta))} \\
 &= e^0 = 1
 \end{aligned}$$

Problem 5

February 21, 2026

Problem 5 Let f and g be two functions such that $\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} g(x) = \infty$.

We say that f **grows faster** than g as x goes to ∞ if $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \infty$, or, equivalently, if

$$\lim_{x \rightarrow \infty} \frac{g(x)}{f(x)} = 0.$$

If the limit exists, namely, if $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = M$, for some positive number M ,

then we say that the functions f and g have **comparable growth rates**.

In other words, we compare the growth rates of functions f and g by computing the limit of the form $\frac{\infty}{\infty}$: $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$.

For each of the following pairs of functions, determine which of the pair grows faster or state that they have comparable growth rates. Justify your answer by computing the limit $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$.

(a) $b^x; x^x; b > 1$

Explanation. $\lim_{x \rightarrow \infty} \frac{b^x}{x^x} = \lim_{x \rightarrow \infty} \left(\frac{b}{x}\right)^x = 0$. Since this limit is of the form 0^∞ , which is a determinate form. Therefore, x^x grows faster.

(b) $x^x; \left(\frac{x}{e}\right)^x$

Explanation. $\lim_{x \rightarrow \infty} \frac{x^x}{(x/e)^x} = \lim_{x \rightarrow \infty} \frac{x^x}{\frac{x^x}{e^x}} = \lim_{x \rightarrow \infty} x^x \cdot \frac{e^x}{x^x} = \lim_{x \rightarrow \infty} e^x = \infty$
Therefore, x^x grows faster.

(c) $x^3; x^3 \cdot \ln(x)$

Explanation. $\lim_{x \rightarrow \infty} \frac{x^3}{x^3 \cdot \ln(x)} = \lim_{x \rightarrow \infty} \frac{1}{\ln(x)} = 0$. Therefore, $x^3 \cdot \ln(x)$ grows faster.

(d) $a^x; b^x; 0 < a < b$

Explanation. $\lim_{x \rightarrow \infty} \frac{a^x}{b^x} = \lim_{x \rightarrow \infty} \left(\frac{a}{b}\right)^x = 0$ Therefore, b^x grows faster

(e) $\log_a(x); \log_b(x); 1 < a < b$

Explanation. $\lim_{x \rightarrow \infty} \frac{\log_a(x)}{\log_b(x)}$

By L'Hopital's Rule: $\lim_{x \rightarrow \infty} \frac{\frac{1}{x \ln(a)}}{\frac{1}{x \ln(b)}} = \lim_{x \rightarrow \infty} \left(\frac{1}{x \ln(a)} \cdot \frac{x \ln(b)}{1} \right) = \frac{\ln b}{\ln a}$

Therefore, $\log_a(x)$ and $\log_b(x)$ grow at comparable rates.

(f) $\ln^3(x); x^{1/2}$

Explanation. $\lim_{x \rightarrow \infty} \frac{\ln^3(x)}{x^{1/2}}$

By L'Hopital's Rule: $\lim_{x \rightarrow \infty} \frac{\ln^3(x)}{x^{1/2}} = \lim_{x \rightarrow \infty} \frac{3 \ln^2(x) \cdot (1/x)}{(1/2)x^{-1/2}} = \lim_{x \rightarrow \infty} \frac{6 \cdot \ln^2(x)}{x^{1/2}}$

By L'Hopital's Rule: $\lim_{x \rightarrow \infty} \frac{12 \ln(x) \cdot (1/x)}{(1/2)x^{-1/2}} = \lim_{x \rightarrow \infty} \frac{24 \cdot \ln(x)}{x^{1/2}}$

By L'Hopital's Rule: $\lim_{x \rightarrow \infty} \frac{24}{x^{1/2}} = 0$ Therefore, $x^{1/2}$ grows faster

(g) $x; \ln(x)\sqrt{x}$

Explanation. $\lim_{x \rightarrow \infty} \frac{x}{\ln(x)\sqrt{x}} = \lim_{x \rightarrow \infty} \frac{x^{1/2}}{\ln(x)}$

By L'Hopital's Rule: $\lim_{x \rightarrow \infty} \frac{(1/2)x^{-1/2}}{x^{-1}} = \lim_{x \rightarrow \infty} (1/2)x^{1/2} = \infty$

Therefore, x grows faster

(h) *Challenge:* $x^{40}; 1.004^x$ (*Hint: Use the substitution $x = \ln(t)$.*)

Explanation.

$$\begin{aligned}
 \lim_{x \rightarrow \infty} \frac{x^{40}}{1.004^x} &= \lim_{t \rightarrow \infty} \frac{[\ln(t)]^{40}}{1.004^{\ln(t)}} \\
 &= \lim_{t \rightarrow \infty} \frac{\ln(t)^{40}}{e^{\ln(1.004) \ln(t)}} \\
 &= \lim_{t \rightarrow \infty} \frac{\ln(t)^{40}}{t^{\ln(1.004)}} \\
 &= \lim_{t \rightarrow \infty} \left(\frac{\ln(t)}{t^{1.004/40}} \right)^{40} \\
 &= \left[\lim_{t \rightarrow \infty} \left(\frac{\ln(t)}{t^{1.004/40}} \right) \right]^{40} \\
 &\stackrel{L.R.}{=} \left[\lim_{t \rightarrow \infty} \frac{\frac{1}{t}}{(1.004/40)t^{(1.004/40)-1}} \right]^{40} \\
 &= \left[\lim_{t \rightarrow \infty} \frac{1}{(1.004/40)t^{((1.004/40)-1+1)}} \right]^{40} \\
 &= 0^{40} \\
 &= 0
 \end{aligned}$$

Therefore, 1.004^x grows faster.

(i) Put the following functions in order of growth rate.

$$x^3 \cdot \ln(x), \ln^3(x), x^x, 1.004^x, x^{40}, x^3$$

Explanation.

$$\ln^3(x) \ll x^3 \ll x^3 \cdot \ln(x) \ll x^{40} \ll 1.004^x \ll x^x$$