

Problem 4

Problem 4


Problem 1 Simplify each of the following expressions.

(a) $\cos^{-1}(\sin(\pi/2))$

Explanation. By the unit circle, $\sin(\pi/2) = 1$, and so we are looking for $\cos^{-1}(1)$. The range of $\cos^{-1}$ is $[0, \pi]$, and so, by properties of inverse functions, $\cos^{-1}(1) = 0$.

(b) $\tan(\sin^{-1}(x/4))$

Explanation. Let $\theta = \sin^{-1}(x/4)$, then $\sin(\theta) = x/4$. We can then draw the corresponding right triangle:



Calling the adjacent side y , by the Pythagorean Theorem we obtain

$$4^2 = x^2 + y^2 \implies y = \sqrt{16 - x^2}$$

Remark: Since θ is in the range of $\sin^{-1}$, it follows that $-\pi/2 < \theta < \pi/2$. Therefore, $\cos(\theta) = y/4 > 0$. Therefore, $y > 0$.

Problem 4

Then

$$\begin{aligned}\tan\left(\sin^{-1}(4/x)\right) &= \tan(\theta), \\ &= \frac{x}{y}, \\ &= \frac{x}{\sqrt{16-x^2}}.\end{aligned}$$

Note: $\tan \theta$ has the same sign as x , since $y > 0$.