STAT340 - Topic 4: Estimation

Estimation

- Suppose you want to apply a RV model, there are usually **parameters** you need to estimate to **fit** the model to the data
  - E.g. estimating p for a binomial, or μ for a normal, etc.
- Estimates are produced using **estimator** functions, i.e. **statistics** of your sample that ideally should converge to the target parameter
  - Law of large numbers useful here
  - Method of moments can also be used
- Remember **estimators are RVs** themselves, since they're a function of data, thus also random, with a distribution, expectation, variance, etc..

Estimation

- Suppose you want to apply a RV model, there are usually **parameters** you need to estimate to **fit** the model to the data
  - E.g. estimating p for a binomial, or μ for a normal, etc.
- Estimates are produced using **estimator** functions, i.e. **statistics** of your sample that ideally should converge to the target parameter
  - Law of large numbers useful here
  - Method of moments can also be used
- Remember **estimators are RVs** themselves, since they're a function of data, thus also random, with a distribution, expectation, variance, etc..

Generally 2 common types of estimates:

- **Point estimates** are single number estimates of the parameter, i.e. a single "best guess" based on available data
- **Interval estimates**, confidence intervals, are ranges that try to "capture" the true value a certain amount of the time

Previous coin example cont.

Continuing from coin flip example from testing chapter, suppose we wish to estimate chance of heads p for a given coin. Experiment:

- Flip it n times and let X be total number of heads,
- We know $X \sim \text{Binomial}(n, p)$ for known n but unknown p.

Previous coin example cont.

Continuing from coin flip example from testing chapter, suppose we wish to estimate chance of heads p for a given coin. Experiment:

- Flip it n times and let X be total number of heads,
- We know $X \sim \text{Binomial}(n, p)$ for known n but unknown p.

- Define the **estimator** $\hat{p} = X/n$ (which is well-defined), note:

$$\text{E}[\hat{p}] = \text{E}\left[\frac{X}{n}\right] = \frac{\text{E}[X]}{n} = \frac{np}{n} = p$$

$$\text{Var}[\hat{p}] = \text{Var}\left[\frac{X}{n}\right] = \frac{\text{Var}[X]}{n^2} = \frac{np(1-p)}{n^2} = \frac{p(1-p)}{n}$$

Previous coin example cont.

Continuing from coin flip example from testing chapter, suppose we wish to estimate chance of heads p for a given coin. Experiment:

- Flip it n times and let X be total number of heads,
- We know $X \sim \text{Binomial}(n, p)$ for known n but unknown p.

- Define the **estimator** $\hat{p} = X/n$ (which is well-defined), note:

$$\text{E}[\hat{p}] = \text{E}\left[\frac{X}{n}\right] = \frac{\text{E}[X]}{n} = \frac{np}{n} = p$$

$$\text{Var}[\hat{p}] = \text{Var}\left[\frac{X}{n}\right] = \frac{\text{Var}[X]}{n^2} = \frac{np(1-p)}{n^2} = \frac{p(1-p)}{n}$$

- Since $\text{E}[\hat{p}] = p$, we say the estimator $\hat{p}$ is **unbiased**
- Also $\text{Var}[\hat{p}] = \frac{p(1-p)}{n} \to 0$ as $n \to \infty$, so larger sample means more precision

Bias of estimator

Let $\hat{\theta}$ be an estimator for the parameter θ, then its **bias** is defined as:

$$\mathrm{Bias}(\hat{\theta}, \theta) = \mathrm{E}[\hat{\theta}] - \theta$$

I.e. it's the systematic difference between what the estimator produces and the true value

Bias of estimator

Let $\hat{\theta}$ be an estimator for the parameter θ, then its **bias** is defined as:

$$\text{Bias}(\hat{\theta}, \theta) = \text{E}[\hat{\theta}] - \theta$$

I.e. it's the systematic difference between what the estimator produces and the true value

Since $\text{E}[\hat{p}] - p = 0$, $\hat{p}$ is unbiased. Let $p = 0.5$. We can find the distribution of $\hat{p}$ for different n values, with either theory or MC:


```
# theoretical pmf table of p-hat with p=0.5 for some n
p.hat.theory = \(n,p=0.5) data.frame(p.hat=0:n/n,prob=dbinom(0:n,n,p))

# MC-simulated distribution of p-hat with p=0.5 for some n
p.hat.mc = \(n,p=0.5,M=1e5) rbinom(M,n,p)/n
```

Let's plot both of these for several values of n, noting the unbiasedness of the estimator for each value of n, as well as how the variance scales with n


$\hat{p}$ distribution for $n = 50$

```
N=50 ; ggplot(p.hat.theory(N)) + geom_col(aes(p.hat,prob),color="black",fill="white") +
  geom_density(data=tibble(X=p.hat.mc(N)),aes(x=X,y=..density../N),adjust=2.5,color="red",size=1)
```




$\hat{p}$ distribution for $n = 200$

```
N=200 ; ggplot(p.hat.theory(N)) + geom_col(aes(p.hat,prob),color="black",fill="white") +
  geom_density(data=tibble(X=p.hat.mc(N)),aes(x=X,y=..density../N),adjust=2,color="red",size=1)
```




$\hat{p}$ distribution for $n = 800$

```
N=800 ; ggplot(p.hat.theory(N)) + geom_linerange(aes(p.hat,ymin=0,ymax=prob),color="grey50") +
  geom_density(data=tibble(X=p.hat.mc(N)),aes(x=X,y=..density../N),adjust=1.5,color="red",size=1)
```



$\hat{p}$ distribution for $n = 3200$

```
N=3200 ; ggplot(p.hat.theory(N)) + geom_linerange(aes(p.hat,ymin=0,ymax=prob),color="grey50") +
  geom_density(data=tibble(X=p.hat.mc(N)),aes(x=X,y=..density../N),adjust=1.5,color="red",size=1)
```



Exercise

Suppose you conducted a simple random sample of 150 people to survey if they prefer candidate A or candidate B for an upcoming local election where voting is mandatory and there's no 3rd option. Out of the 150 people surveyed, 90 preferred A, 60 preferred B.

Estimate the proportion of all residents that would support candidate A over B. Do you expect this estimate to be biased? Estimate the SD of this estimate.

Exercise

Suppose you conducted a simple random sample of 150 people to survey if they prefer candidate A or candidate B for an upcoming local election where voting is mandatory and there's no 3rd option. Out of the 150 people surveyed, 90 preferred A, 60 preferred B.

Estimate the proportion of all residents that would support candidate A over B. Do you expect this estimate to be biased? Estimate the SD of this estimate.

$\hat{p} = 90/150 = 0.6$. This estimate should be unbiased.

Estimate variance by substituting $\hat{p}$ for p, i.e:

$$\mathrm{Var}[\hat{p}] = \frac{p(1-p)}{n} = \frac{0.6(1-0.6)}{150} = 0.0016$$

So SD $= 0.04$

Method of moments

Extending this further, the **method of moments** (MOM) uses the moments of a RV, i.e. $\mathrm{E}[X], \mathrm{E}[X^2], \mathrm{E}[X^3], \ldots$ to estimate parameters.

Recall $\mathrm{Var}[X] = \mathrm{E}[X^2] - (\mathrm{E}[X])^2$, so variance also called the **second central moment**.

Method of moments

Extending this further, the **method of moments** (MOM) uses the moments of a RV, i.e. $\mathrm{E}[X], \mathrm{E}[X^2], \mathrm{E}[X^3], \ldots$ to estimate parameters.

Recall $\mathrm{Var}[X] = \mathrm{E}[X^2] - (\mathrm{E}[X])^2$, so variance also called the **second central moment**.

Example: suppose we have X_i iid from an exponential distribution with some unknown λ. We know from theory that:

$$\bar{X} \xrightarrow{n} \mathrm{E}[X] = \frac{1}{\lambda}$$

Therefore it stands to reason that $\hat{\lambda} = 1/\bar{X}$ should be a good estimator for λ

Method of moments

Extending this further, the **method of moments** (MOM) uses the moments of a RV, i.e. $\mathrm{E}[X], \mathrm{E}[X^2], \mathrm{E}[X^3], \ldots$ to estimate parameters.

Recall $\mathrm{Var}[X] = \mathrm{E}[X^2] - (\mathrm{E}[X])^2$, so variance also called the **second central moment**.

Example: suppose we have X_i iid from an exponential distribution with some unknown λ. We know from theory that:

$$\bar{X} \xrightarrow{n} \mathrm{E}[X] = \frac{1}{\lambda}$$


Therefore it stands to reason that $\hat{\lambda} = 1/\bar{X}$ should be a good estimator for λ

Example: suppose we sample from $X \sim \mathrm{Exp}(\lambda = 5)$ (ofc normally we don't know λ). Imagine we repeat the following experiment a large number of times: draw $n = 200$ observations, then apply $\hat{\lambda} = 1/\bar{X}$. What is the sampling distribution of $\hat{\lambda}$?

```
# estimate lambda across many samples using 1/mean
lambda.hats = replicate(1e4,1/mean(rexp(200,5)))
```

Exponential example (cont.)


```
hist(lambda.hats) ; abline(v=mean(lambda.hats),col="red",lwd=4)
```



Exponential example, using SD


Note: since $\sigma^2 = 1/\lambda^2$, also possible to use (sample SD)$^{-1}$ to estimate λ, though generally higher moments have more variance, so may be worse, but let's test it!

```
# estimate lambda across many samples using 1/sd
lambda.hats2 = replicate(1e4,1/sd(rexp(200,5)))
hist(lambda.hats2) ; abline(v=mean(lambda.hats2),col="red",lwd=4)
```



Exponential example, replotting both statistics

```
enframe(c("1/mean"=lambda.hats,"1/sd"=lambda.hats2)) %>% mutate(Estimator=gsub("\\d+$","",name)) %>%
  ggplot(aes(value,color=Estimator)) + geom_density(adjust=2,linewidth=1) + xlab("lambda hat") +
  ggtitle("Comparison of 1/mean, 1/sd estimators for exponential lambda parameter")
```



MOM exercise (gamma dist)

Can also combine moments in expression for more complex problems. Ex:

A Gamma RV takes two parameters, $\alpha, \theta > 0$ (shape & scale parameters).
If $X \sim \mathrm{Gamma}(\alpha, \theta)$ we have $\mathrm{E}[X] = \alpha\theta$ and $\mathrm{Var}[X] = \alpha\theta^2$.
Use MOM to define usable $\hat{\theta}$, $\hat{\alpha}$ for the two parameters.

MOM exercise (gamma dist)

Can also combine moments in expression for more complex problems. Ex:

A Gamma RV takes two parameters, $\alpha, \theta > 0$ (shape & scale parameters).
If $X \sim \mathrm{Gamma}(\alpha, \theta)$ we have $\mathrm{E}[X] = \alpha\theta$ and $\mathrm{Var}[X] = \alpha\theta^2$.
Use MOM to define usable $\hat{\theta}, \hat{\alpha}$ for the two parameters.


Ans:

1. Since $\mathrm{Var}[X]/\,\mathrm{E}[X] = \theta$, we can use $\hat{\theta} = S^2/\bar{X}$
2. Since $(\mathrm{E}[X])^2/\,\mathrm{Var}[X] = \alpha$, we can use $\hat{\alpha} = (\bar{X})^2/S^2$

Other estimators are possible using higher moments or other methods, but these are the simplest using the information available. Demo on next slide.


MOM exercise (gamma dist) cont.


```r
# pick some arbitrary shape/scale parameters and plot gamma (alpha=3,theta=2)
plot(\(x)dgamma(x,shape=3,scale=2),from=0,to=20,main="PDF of Gamma(shape=3,scale=2)")
```



MOM exercise (gamma dist) cont.

```r
# draw many samples of n=100 and check results of estimators are good (alpha=3,theta=2)
estimates = replicate(1e4, { x=rgamma(100,shape=3,scale=2); c(var(x)/mean(x),mean(x)^2/var(x)) })
alpha.hats = estimates[2,] ; theta.hats = estimates[1,]
hist(alpha.hats) ; abline(v=mean(alpha.hats),col="red",lwd=4)
hist(theta.hats) ; abline(v=mean(theta.hats),col="red",lwd=4)
```





Interval estimates

Often, want to know how good a point estimate is, can use **confidence interval** instead

Review exercise: continuing from previous exercise where 90/150 people surveyed preferred candidate A to B. Give a 95% confidence interval for this result.

Interval estimates

Often, want to know how good a point estimate is, can use **confidence interval** instead

Review exercise: continuing from previous exercise where 90/150 people surveyed preferred candidate A to B. Give a 95% confidence interval for this result.

Ans: $\hat{p} = 0.6$ with estimated SD, or **standard error** (SE) of 0.04. To get 95% confidence interval, need about $2\times$ SE on either side of estimate, so about $(0.52, 0.68)$

Interval estimates

Often, want to know how good a point estimate is, can use **confidence interval** instead

Review exercise: continuing from previous exercise where 90/150 people surveyed preferred candidate A to B. Give a 95% confidence interval for this result.


Ans: $\hat{p} = 0.6$ with estimated SD, or **standard error** (SE) of 0.04. To get 95% confidence interval, need about $2\times$ SE on either side of estimate, so about $(0.52, 0.68)$

MC version: resimulate with $p = \hat{p}$, then find 95% central band of new resimulated point estimates

```
new.X = rbinom(1e5,150,90/150)
new.phats = new.X/150
(ci = quantile(new.phats,c(.025,.975)))
```

```
 2.5% 97.5%
 0.52  0.68
```

```
hist(new.phats)
abline(v=ci, col="red", lwd=3)
```



Wait what? MC interval explained

Why does this work? Suppose we are trying to estimate a parameter θ with our observed sample statistic S. After fitting with S, we resimulate to get a new set of statistics S^*. One can check these are generally approximately true with typical sample sizes:

1. $\mathrm{E}[S^*] = S$
2. $\mathrm{Var}[S^*] \approx \mathrm{Var}[S]$
3. S^* symmetric about $\mathrm{E}[S^*]$

Wait what? MC interval explained

Why does this work? Suppose we are trying to estimate a parameter θ with our observed sample statistic S. After fitting with S, we resimulate to get a new set of statistics S^*. One can check these are generally approximately true with typical sample sizes:

1. $\mathrm{E}[S^*] = S$
2. $\mathrm{Var}[S^*] \approx \mathrm{Var}[S]$
3. S^* symmetric about $\mathrm{E}[S^*]$

Together, these properties imply that the **distribution of S^* around S is *similar* to the distribution of S around θ**. Thus, resimulating S^* around S estimates the variability of S, which is used to compute the CI.

This is a highly simplified argument. For more mathematical details, see Karl Rohe's writeup here. Visual demo on next page…

Demo of MC interval

Suppose we sample $n = 100$ from a known normal RV with unknown $\mu = 10, \sigma^2 = 2^2$. We observe the following sample statistics:

```
x = rnorm(100, 10, 2) ; c(mean(x), sd(x))
```

```
[1] 10.217775  1.796399
```

Demo of MC interval

Suppose we sample $n = 100$ from a known normal RV with unknown $\mu = 10, \sigma^2 = 2^2$. We observe the following sample statistics:

```
x = rnorm(100, 10, 2) ; c(mean(x), sd(x))
```

```
[1] 10.217775  1.796399
```

Thus, $S = \bar{X} \approx 10.22$. Then, we resimulate many times using S to get distribution of S^*:

```
new.means = replicate(1e4,mean(rnorm(100,mean(x),sd(x)))) ; c(ms<-mean(new.means),ss<-sd(new.means))
```

```
[1] 10.2178332  0.1789841
```

Demo of MC interval

Suppose we sample $n = 100$ from a known normal RV with unknown $\mu = 10, \sigma^2 = 2^2$. We observe the following sample statistics:

```
x = rnorm(100, 10, 2) ; c(mean(x), sd(x))
```

```
[1] 10.217775  1.796399
```

Thus, $S = \bar{X} \approx 10.22$. Then, we resimulate many times using S to get distribution of S^*:


```
new.means = replicate(1e4,mean(rnorm(100,mean(x),sd(x)))) ; c(ms<-mean(new.means),ss<-sd(new.means))
```

```
[1] 10.2178332  0.1789841
```

Compare distribution of $S = \bar{X}$ around $\mu = 10$ with distribution of S^* around $S \approx 10.22$


```
# adjust=3.5 smoothes out S* a bit more, but distorts the density curve,
# hence quantiles of S* later plotted with ±z=2.2 to match distorted density
ggplot() + geom_function(aes(color="S"),fun=\(x)dnorm(x,10,2/10),xlim=c(9.2,10.8),linewidth=1.5) +
  geom_density(data=tibble(X=new.means),aes(X,color="S*"),adjust=3.5,linewidth=1.5,key_glyph="path")
  geom_linerange(aes(x=c(10,mean(x))),ymin=0,ymax=1.99) + labs(color="Statistic",x=NULL,y=NULL) +
  stat_function(fun=\(x)dnorm(x,10,2/10),geom="area",xlim=c(9.2,9.6),fill="#1d9d79",alpha=.5) +
  stat_function(fun=\(x)dnorm(x,10,2/10),geom="area",xlim=c(10.4,10.8),fill="#1d9d79",alpha=.5) +
  stat_function(fun=\(x)dnorm(x,ms,ss*1.12),geom="area",xlim=ms+ss*c(-4,-2.2),fill="#d76420",alpha=.5
  stat_function(fun=\(x)dnorm(x,ms,ss*1.12),geom="area",xlim=ms+ss*c(2.2,4),fill="#d76420",alpha=.5)
  scale_x_continuous(breaks=seq(9.2,10.8,.2),minor_breaks=NULL,expand=0) +
  scale_y_continuous(expand=0,limits=c(0,2.1))
```

Demo of MC interval (cont.)



- Green curve is theoretical distribution of $S = \bar{X} \sim \text{Normal}(10, 2^2/100)$ about $\mu = 10$
- Orange curve is simulated distribution of S^* about observed sample mean $\bar{X} \approx 10.22$
- Note two curves have similar shape/width, just shifted
- 95% of time, $\bar{X} \in (9.6, 10.4)$, so that quantiles of S^*, taken as CI, contains $\mu = 10$

Demo of MC interval (cont.)



- 5% of the time, $\bar{X} \notin (9.6, 10.4)$, and quantiles of S^* taken as CI won't contain $\mu = 10$
- Here, in this "unlucky" rerun, observed sample mean $\bar{X} \approx 10.5$
- This is outside the central 95% range of S, so the quantiles of S^* don't contain μ

Similar argument/demo approximately applies to other RVs too, not just normal

Generalized MC confidence interval steps:

1. Identify RV **model** & **parameter(s)** for data

Generalized MC confidence interval steps:

1. Identify RV **model** & **parameter(s)** for data

2. Pick point **estimator** function (e.g. using LLN, MOM, or other method)

Generalized MC confidence interval steps:

1. Identify RV **model** & **parameter(s)** for data

2. Pick point **estimator** function (e.g. using LLN, MOM, or other method)

3. Apply estimator to get initial **point estimate** (S)

Generalized MC confidence interval steps:

1. Identify RV **model** & **parameter(s)** for data

2. Pick point **estimator** function (e.g. using LLN, MOM, or other method)

3. Apply estimator to get initial **point estimate** (S)

4. Assume true parameter equals point estimate, then **resimulate** many times, each time:
    - Generate **new dataset of same size as original**,
    - Compute new point estimate (S^*) of newly generated data.

Generalized MC confidence interval steps:

1. Identify RV **model** & **parameter(s)** for data

2. Pick point **estimator** function (e.g. using LLN, MOM, or other method)

3. Apply estimator to get initial **point estimate** (S)

4. Assume true parameter equals point estimate, then **resimulate** many times, each time:
    - Generate **new dataset of same size as original**,
    - Compute new point estimate (S^*) of newly generated data.

5. Take appropriate quantiles of S^* distribution as **confidence interval**.

Generalized MC confidence interval steps:

1. Identify RV **model** & **parameter(s)** for data

2. Pick point **estimator** function (e.g. using LLN, MOM, or other method)

3. Apply estimator to get initial **point estimate** (S)

4. Assume true parameter equals point estimate, then **resimulate** many times, each time:
    - Generate **new dataset of same size as original**,
    - Compute new point estimate (S^*) of newly generated data.

5. Take appropriate quantiles of S^* distribution as **confidence interval**.

6. Interpret results in context of original question!
    - E.g. “We are 95% confident that true [parameter] is in [interval].”

Exercise:

Suppose we have the following $n = 30$ observations from a Poisson distribution:

> 6, 7, 8, 12, 6, 12, 13, 9, 9, 4, 6, 5, 9, 7, 10, 8, 10, 16, 7, 10, 12, 6, 9, 5, 6, 7, 2, 7, 11, 7

1. What's a good point estimator function?
2. Give a point estimate of the rate parameter λ.
3. Find a 95% confidence interval for λ.

Answer in steps:

Exercise:

Suppose we have the following $n = 30$ observations from a Poisson distribution:

> 6, 7, 8, 12, 6, 12, 13, 9, 9, 4, 6, 5, 9, 7, 10, 8, 10, 16, 7, 10, 12, 6, 9, 5, 6, 7, 2, 7, 11, 7

1. What's a good point estimator function?
2. Give a point estimate of the rate parameter λ.
3. Find a 95% confidence interval for λ.

Answer in steps:

1. Sample mean works great here. Variance also works, but will have higher SE.

Exercise:

Suppose we have the following $n = 30$ observations from a Poisson distribution:

> 6, 7, 8, 12, 6, 12, 13, 9, 9, 4, 6, 5, 9, 7, 10, 8, 10, 16, 7, 10, 12, 6, 9, 5, 6, 7, 2, 7, 11, 7

1. What's a good point estimator function?
2. Give a point estimate of the rate parameter λ.
3. Find a 95% confidence interval for λ.

Answer in steps:

1. Sample mean works great here. Variance also works, but will have higher SE.
2. Find sample mean $\bar{X} \approx 8.2$.

Exercise:

Suppose we have the following $n = 30$ observations from a Poisson distribution:

> 6, 7, 8, 12, 6, 12, 13, 9, 9, 4, 6, 5, 9, 7, 10, 8, 10, 16, 7, 10, 12, 6, 9, 5, 6, 7, 2, 7, 11, 7

1. What's a good point estimator function?
2. Give a point estimate of the rate parameter λ.
3. Find a 95% confidence interval for λ.

Answer in steps:

1. Sample mean works great here. Variance also works, but will have higher SE.
2. Find sample mean $\bar{X} \approx 8.2$.

```
# 3. repeatedly generate n=30 new poisson points and find new lambda hats, then take quantiles
new.lambda.hats = replicate(1e4,mean(rpois(30,mean(x))))
quantile(new.lambda.hats,c(.025,.975))
```

```
    2.5%    97.5% 
7.200000 9.233333 
```

Applied example: mark & recapture

More interesting example. In ecology, mark & recapture is a common technique to estimate the population of a species of animals where a 1st sample is marked and released, then later a 2nd sample is taken and the number already marked is counted. Assume in both samples, any animal in population is **equally likely** to be captured.

Applied example: mark & recapture

More interesting example. In ecology, mark & recapture is a common technique to estimate the population of a species of animals where a 1st sample is marked and released, then later a 2nd sample is taken and the number already marked is counted. Assume in both samples, any animal in population is **equally likely** to be captured. Let:

- N be total number of animals in population
- n be number of animals marked in 1st sample
- K be number of animals captured in 2nd sample
- k be number of animals in 2nd sample that are marked

Applied example: mark & recapture

More interesting example. In ecology, mark & recapture is a common technique to estimate the population of a species of animals where a 1st sample is marked and released, then later a 2nd sample is taken and the number already marked is counted. Assume in both samples, any animal in population is **equally likely** to be captured. Let:

- N be total number of animals in population
- n be number of animals marked in 1st sample
- K be number of animals captured in 2nd sample
- k be number of animals in 2nd sample that are marked

Under assumption, $k/K = n/N$, so an estimator could be $\hat{N} = nK/k$. This is called the Lincoln–Petersen estimator.

Ex: You're studying walleyes in Lake Mendota. You catch & mark 500 fish. A few weeks later, you catch another 500, 12 of which were already marked. Give a point estimate for the total walleye population N in the lake.

Applied example: mark & recapture

More interesting example. In ecology, mark & recapture is a common technique to estimate the population of a species of animals where a 1st sample is marked and released, then later a 2nd sample is taken and the number already marked is counted. Assume in both samples, any animal in population is **equally likely** to be captured. Let:

- N be total number of animals in population
- n be number of animals marked in 1st sample
- K be number of animals captured in 2nd sample
- k be number of animals in 2nd sample that are marked

Under assumption, $k/K = n/N$, so an estimator could be $\hat{N} = nK/k$. This is called the Lincoln–Petersen estimator.

Ex: You're studying walleyes in Lake Mendota. You catch & mark 500 fish. A few weeks later, you catch another 500, 12 of which were already marked. Give a point estimate for the total walleye population N in the lake. **Ans:** using estimator, $\hat{N} = 500^2/12 \approx 20833$.

Applied example: mark & recapture (cont.)

Next, use MC to find a 95% confidence interval for your estimate (follow the checklist!)

(*answer in steps*) First, simulate one new "observed" k for $N = \hat{N}$:

Applied example: mark & recapture (cont.)

Next, use MC to find a 95% confidence interval for your estimate (follow the checklist!)

(*answer in steps*) First, simulate one new "observed" k for $N = \hat{N}$:

Different ways to simulate:

```
# suppose first 1-500 of fish are marked, then can simulate n using sample()
sim = \(N,n=500,K=500) sum(sample(N,K)<=n)
sim(20833)
```

```
[1] 13
```

```
# can also approximate with rbinom(), i.e. each fish modeled as n/N chance of being marked
sim2 = \(N,n=500,K=500,M=1) rbinom(M,K,n/N)
sim2(20833)
```

```
[1] 8
```


```
# can even approximate with rpois(), i.e. draw fish with averate rate of n*K/N
sim3 = \(N,n=500,K=500,M=1) rpois(M,n*K/N)
sim3(20833)
```

```
[1] 11
```

Can we check if all three of these give similar results?

Applied example: mark & recapture (cont.)

```
# can check all three are comparable
tibble(x=c(replicate(1e4,sim(20833)),sim2(20833,M=1e4),sim3(20833,M=1e4)),
       type=rep(c("binom","sample","pois"),each=1e4)) %>%
  ggplot(aes(x)) + facet_grid(type~.) + geom_bar()
```



Applied example: mark & recapture (cont.)

Next, repeat this many times, each time computing $\hat{N}$, then take quantiles as CI

Applied example: mark & recapture (cont.)

Next, repeat this many times, each time computing $\hat{N}$, then take quantiles as CI

```
# using sample()
N.hats = 500^2/replicate(1e4,sim(20833))
quantile(N.hats,c(.025,.975))
```

```
    2.5%    97.5%
13157.89 41666.67
```

```
# using rbinom()
N.hats = 500^2/sim2(20833,M=1e4)
quantile(N.hats,c(.025,.975))
```

```
    2.5%    97.5%
13157.89 41666.67
```

```
# using rpois()
N.hats = 500^2/sim3(20833,M=1e4)
quantile(N.hats,c(.025,.975))
```

```
    2.5%    97.5%
13157.89 41666.67
```

All methods give same 95% CI for N of $(13158, 41667)$.

Assigned readings

Please read the following:

- Subsection 22.7 in Brian's notes on CLT based confidence intervals (basically the same as what we did previously in 240)
- Short page on MSE of estimators & bias-variance decomposition