# STAT340 - Topic 5: Prediction

# Simple linear regression (SLR) review

Quick overview of SLR (one predictor, one response) from 240:

- Let $X, Y$ be numeric RVs with **clear linear relationship**; recall theoretical correlation:

$$\rho_{X,Y} = \frac{\mathrm{Cov}(X, Y)}{\sqrt{\mathrm{Var}[X]\,\mathrm{Var}[Y]}}$$

  In sample, we observe **sample correlation coefficient**:

$$r = \frac{\frac{1}{n-1}\sum(x_i - \bar{x})(y_i - \bar{y})}{s_x s_y}$$

# Simple linear regression (SLR) review

Quick overview of SLR (one predictor, one response) from 240:

- Let $X, Y$ be numeric RVs with **clear linear relationship**; recall theoretical correlation:

$$\rho_{X,Y} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}[X]\,\text{Var}[Y]}}$$

  In sample, we observe **sample correlation coefficient**:

$$r = \frac{\frac{1}{n-1}\sum(x_i - \bar{x})(y_i - \bar{y})}{s_x s_y}$$

- Note $n-1$ used in numerator instead of $n$, similar to sample SDs $s_x$, $s_y$
- $\rho, r$ always in $[-1, 1]$, where:
  - Sign (positive/negative/zero) indicates **direction of slope** of best fit line, and
  - Absolute value indicates **strength of correlation** (closer to ±1 means stronger)
- **Note:** correlation coefficient ***assumes*** linear relationship; ***must*** be checked with plot!
- Interactive correlation demo: https://bwu62.shinyapps.io/correlation

# SLR model

Let $(x_i, y_i)$ be sample of size $n$ of apparently linear points. **Model equation** for SLR:

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

- $\beta_0, \beta_1$ are true intercept & slope (treated as constants like $x_i$)
- $\varepsilon_i \overset{\text{iid}}{\sim} \text{Normal}(0, \sigma^2)$ are **errors** of each point (the only variable on RHS!)

# SLR model


Let $(x_i, y_i)$ be sample of size $n$ of apparently linear points. **Model equation** for SLR:

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

- $\beta_0, \beta_1$ are true intercept & slope (treated as constants like $x_i$)
- $\varepsilon_i \overset{\text{iid}}{\sim} \text{Normal}(0, \sigma^2)$ are **errors** of each point (the only variable on RHS!)

## MC simulated demo:

```
b0  = -3              # true intercept
b1  = 0.5             # true slope
n   = 10              # sample size
m   = 10              # min X range
M   = 20              # max X range
sig = 2               # true sigma
x   = runif(n,m,M)    # generate Xs
e   = rnorm(n,0,sig)  # generate errors
y   = b0 + b1*x + e   # combine for Ys
```




# OLS fitting

Typically **ordinary least squares** (OLS) used to fit, i.e. **sum of squared errors** minimized. Given some $\beta_0', \beta_1'$, define OLS **loss function**:

$$\ell(\beta_0', \beta_1') = \sum_{i=1}^{n} (y_i - (\beta_0' + \beta_1' x_i))^2$$

To minimize $\ell(\beta_0', \beta_1')$, set partial derivatives vs $\beta_0', \beta_1'$ to 0 and solve (full derivation)




# OLS fitting

Typically **ordinary least squares** (OLS) used to fit, i.e. **sum of squared errors** minimized. Given some $\beta_0', \beta_1'$, define OLS **loss function**:

$$\ell(\beta_0', \beta_1') = \sum_{i=1}^{n} (y_i - (\beta_0' + \beta_1' x_i))^2$$

To minimize $\ell(\beta_0', \beta_1')$, set partial derivatives vs $\beta_0', \beta_1'$ to 0 and solve (full derivation)




Best fit coefficients:

$$\hat{\beta}_1 = r \cdot \frac{s_y}{s_x}$$
$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

- **Note:** equivalent to line with slope $\hat{\beta}_1$ passing through $(\bar{x}, \bar{y})$
- Can be shown these estimators are unbiased with least variance

# Inference

After fitting, our **fitted** or **predicted values** $\hat{y}_i$ are found by pluggin $x_i$ back:

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$$

Then our **residuals** $\hat{\varepsilon}_i$ are the realizations of $\varepsilon_i \sim \text{Normal}(0, \sigma^2)$, defined:

$$\hat{\varepsilon}_i = y_i - \hat{y}_i = y_i - (\hat{\beta}_0 + \hat{\beta}_1 x_i)$$

# Inference

After fitting, our **fitted** or **predicted values** $\hat{y}_i$ are found by pluggin $x_i$ back:

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$$

Then our **residuals** $\hat{\varepsilon}_i$ are the realizations of $\varepsilon_i \sim \text{Normal}(0, \sigma^2)$, defined:

$$\hat{\varepsilon}_i = y_i - \hat{y}_i = y_i - (\hat{\beta}_0 + \hat{\beta}_1 x_i)$$

Using these, we estimate $\sigma^2$ using the **mean squared error** of the residuals:

$$\hat{\sigma}^2 = \frac{\ell(\hat{\beta}_0, \hat{\beta}_1)}{n-2} = \frac{\sum \hat{\varepsilon}_i^2}{n-2} = \frac{1}{n-2} \sum_i (y_i - (\hat{\beta}_0 + \hat{\beta}_1 x_i))^2$$

Can be shown $\hat{\sigma}^2$ is unbiased estimator for $\sigma^2$. Note: $\sigma^2$ is important since this represents the **amount of noise** in the data. Higher $\sigma^2 \rightarrow$ more noise $\rightarrow$ less precise results.

Remember to check if $\hat{\varepsilon}_i \sim \text{Normal}(0, \sigma^2)$ with residual plots!

# SE of coefficients

Slope SE can be easily derived now. First, note:

$$\begin{aligned}\sum(x_i - \bar{x})(y_i - \bar{y}) &= \sum(x_i - \bar{x})y_i - \bar{y}\cancel{\sum(x_i - \bar{x})} \\ &= \sum(x_i - \bar{x})(\beta_0 + \beta_1 x_i + \varepsilon_i)\end{aligned}$$

# SE of coefficients

Slope SE can be easily derived now. First, note:

$$\begin{aligned}\sum(x_i-\bar{x})(y_i-\bar{y}) &= \sum(x_i-\bar{x})y_i - \bar{y}\cancel{\sum(x_i-\bar{x})} \\ &= \sum(x_i-\bar{x})(\beta_0+\beta_1 x_i+\varepsilon_i)\end{aligned}$$

$$\begin{aligned}\mathrm{Var}[\hat{\beta}_1] &= \mathrm{Var}\left[\frac{\sum(x_i-\bar{x})(y_i-\bar{y})}{\sum(x_i-\bar{x})^2}\right] = \mathrm{Var}\left[\frac{\sum_i(x_i-\bar{x})(\cancel{\beta_0+\beta_1 x_i}+\varepsilon_i)}{\sum(x_i-\bar{x})^2}\right] \\ &= \frac{\sum(x_i-\bar{x})^2\mathrm{Var}(\varepsilon_i)}{\left(\sum(x_i-\bar{x})^2\right)^2} = \frac{\sigma^2}{\sum_i(x_i-\bar{x})^2}\end{aligned}$$

# SE of coefficients

Slope SE can be easily derived now. First, note:

$$\begin{aligned}\sum (x_i - \bar{x})(y_i - \bar{y}) &= \sum (x_i - \bar{x}) y_i - \bar{y} \cancel{\sum (x_i - \bar{x})} \\ &= \sum (x_i - \bar{x})(\beta_0 + \beta_1 x_i + \varepsilon_i)\end{aligned}$$

$$\begin{aligned}\mathrm{Var}[\hat{\beta}_1] &= \mathrm{Var}\left[\frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sum (x_i - \bar{x})^2}\right] = \mathrm{Var}\left[\frac{\sum_i (x_i - \bar{x})(\cancel{\beta_0 + \beta_1 x_i} + \varepsilon_i)}{\sum (x_i - \bar{x})^2}\right] \\ &= \frac{\sum (x_i - \bar{x})^2 \mathrm{Var}(\varepsilon_i)}{\left(\sum (x_i - \bar{x})^2\right)^2} = \frac{\sigma^2}{\sum_i (x_i - \bar{x})^2}\end{aligned}$$

Then, using our sample we can estimate $\mathrm{SE}(\hat{\beta}_1) = \sqrt{\dfrac{\hat{\sigma}^2}{s_x^2 (n-1)}}$. Similar process gives:

$\mathrm{Var}[\hat{\beta}_0] = \sigma^2 \left[\dfrac{1}{n} + \dfrac{\bar{x}^2}{\sum_i (x_i - \bar{x})^2}\right]$, so estimate $\mathrm{SE}(\hat{\beta}_0) = \sqrt{\hat{\sigma}^2 \left[\dfrac{1}{n} + \dfrac{\bar{x}^2}{s_x^2 (n-1)}\right]}$

# Inference on coefficients

Coefficients $\hat{\beta}_0, \hat{\beta}_1$ follow T-distributions with $n-2$ degrees of freedom.

- For hypothesis testing, e.g. $H_0 : \beta_1 = 0$ vs $H_a : \beta_1 \neq 0$, use:

$$t_{\text{obs}} = \frac{\hat{\beta}_1 - 0}{\text{SE}(\hat{\beta}_1)} \sim T_{n-2}$$

# Inference on coefficients

Coefficients $\hat{\beta}_0, \hat{\beta}_1$ follow T-distributions with $n-2$ degrees of freedom.

- For hypothesis testing, e.g. $H_0 : \beta_1 = 0$ vs $H_a : \beta_1 \neq 0$, use:

$$t_{\text{obs}} = \frac{\hat{\beta}_1 - 0}{\text{SE}(\hat{\beta}_1)} \sim T_{n-2}$$

- For confidence intervals, e.g. C% interval for $\beta_1$, use:

$$C\% \text{ or } (1-\alpha) \text{ interval} = \hat{\beta}_1 \pm t_{\alpha/2,\, n-2} \cdot \text{SE}(\hat{\beta}_1)$$

  - $t_{\alpha/2,\, n-2}$ is T critical value, e.g. `qt(0.975,df=n-2)` for 95%
  - Can also use `confint()` function

# Confidence/prediction interval for predictions


Suppose we take new observation at $X = x_{\text{new}}$, can make two different intervals at there:

- **Confidence interval** for mean predicted response $\hat{y}$ (i.e. for the regression line), which has:

$$\text{SE}(\hat{y}) = \sqrt{\hat{\sigma}^2 \left[ \frac{1}{n} + \frac{(x_{\text{new}} - \bar{x})^2}{s_x^2 (n-1)} \right]}$$

- **Prediction interval** for next individual data point $y_{\text{new}}$ (i.e. for the data points), which has:

$$\text{SE}(y_{\text{new}}) = \sqrt{\hat{\sigma}^2 \left[ 1 + \frac{1}{n} + \frac{(x_{\text{new}} - \bar{x})^2}{s_x^2 (n-1)} \right]}$$




- Can also use `predict(lm.object, interval="confidence")` or `"prediction"`

# Cumulative SLR exercise!

Suppose you have the following observations:

```
x = c(14.4, 15.3, 11.2, 10.7, 11.9, 13, 18.4, 15.3, 17.3, 10.1, 18.4, 18.1, 16.6, 15.7, 10.3,
19, 10.9, 16.3, 17, 17.2, 14.3, 12.3, 15.8, 17.8, 15.2, 20, 11.1, 18.3, 15.9, 12.1)
y = c(50.9, 50.3, 41.7, 25, 34.6, 50.4, 65.5, 48.8, 62.2, 41, 64.4, 71.5, 66.8, 58.5, 38.8,
63.7, 36.5, 62.4, 68, 66.4, 49.7, 31.3, 51.3, 64.3, 56.3, 76.7, 38.8, 59.3, 55.1, 33.4)
```

Do the following (try both manually and using R functions):


1. Plot the data, check if SLR is appropriate, and find the correlation $r$,
2. Fit a best fit line, plot it, and find the coefficients $\hat{\beta}_1, \hat{\beta}_0$,
3. Find the fitted values $\hat{y}_i$ and residuals $\hat{\varepsilon}_i$, and check model assumptions,
4. Estimate SD of errors $\sigma$ and SEs of coefficients $\hat{\beta}_1, \hat{\beta}_0$,
5. Find p-value for testing $\beta_1 = 0$ vs not, and give 95% CIs for $\beta_1, \beta_0$,
6. At $X = 18$, give and interpret 95% confidence and prediction intervals.

Answers using R functions shown first:

# Exercise answers using R functions

```
# plot data, best fit line, and find correlation
ggplot(tibble(x,y),aes(x,y)) + geom_point() + geom_smooth(method=lm,se=F)
cor(x,y)
```

```
[1] 0.9196951
```



# Exercise answers using R functions (cont.)

```r
# fit with lm(), print summary with coefficients, SEs, and residual SE
# also shows coefficient p-values for testing each =0 vs ≠0
model = lm(y ~ x, data = tibble(x,y))
summary(model)
```

```
Call:
lm(formula = y ~ x, data = tibble(x, y))

Residuals:
     Min       1Q   Median       3Q      Max
-10.2063  -4.5742   0.2184   4.3378   8.6965

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  -9.9455     5.1557  -1.929   0.0639 .
x             4.1831     0.3375  12.395 6.92e-13 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.381 on 28 degrees of freedom
Multiple R-squared:  0.8458,    Adjusted R-squared:  0.8403
F-statistic: 153.6 on 1 and 28 DF,  p-value: 6.916e-13
```

# Exercise answers using R functions (cont.)


```r
# fitted values
fitted(model) %>% unname %>% round(2)
```

```
 [1] 50.29 54.06 36.90 34.81 39.83 44.43 67.02 54.06 62.42 32.30 67.02 65.77 59.49 55.73 33.14 69.53
[17] 35.65 58.24 61.17 62.00 49.87 41.51 56.15 64.51 53.64 73.72 36.49 66.60 56.57 40.67
```

```r
# residuals
resid(model) %>% unname %>% round(2)
```

```
 [1]   0.61  -3.76   4.80  -9.81  -5.23   5.97  -1.52  -5.26  -0.22   8.70  -2.62   5.73   7.31
[14]   2.77   5.66  -5.83   0.85   4.16   6.83   4.40  -0.17 -10.21  -4.85  -0.21   2.66   2.98
[27]   2.31  -7.30  -1.47  -7.27
```

```r
# check model
par(mfrow=c(1,2))  ;  plot(model,which=1:2)
```



# Exercise answers using R functions (cont.)

```r
# confidence intervals for coefficients
confint(model, level = 0.95)
```

```
                2.5 %    97.5 %
(Intercept) -20.506486 0.6153899
x             3.491761 4.8743932
```

```r
# confidence/prediction intervals for prediction at new X=18
predict(model, newdata = data.frame(x=18), interval = "confidence")
```

```
       fit      lwr     upr
1 65.34984 62.45838 68.2413
```

```r
predict(model, newdata = data.frame(x=18), interval = "prediction")
```

```
       fit      lwr     upr
1 65.34984 53.95459 76.7451
```

Answers using manual computation shown on next slides:

# Exercise answers computing manually

```r
# define useful values
n = length(x) ; mx = mean(x) ; my = mean(y) ; sx = sd(x) ; sy = sd(y) ; r = cor(x,y)
# best fit slope and intercept
c(b1 <- r*sy/sx, b0 <- my-b1*mx)
```

```
[1]  4.183077 -9.945548
```

```r
# fitted and residual values
fitted = b0 + b1*x  ;  resid = y - fitted
# residual standard error
(sigma.hat = sqrt(sum(resid^2)/(n-2)))
```

```
[1] 5.380917
```

```r
# estimated SE of coefficients
(se.b1 = sigma.hat*sqrt(1/(sx^2*(n-1))))
```

```
[1] 0.3374895
```

```r
(se.b0 = sigma.hat*sqrt(1/n+mx^2/(sx^2*(n-1))))
```

```
[1] 5.155683
```

```r
# test true slope =0 vs ≠0
(b1.stat = b1/se.b1)
```

```
[1] 12.39469
```

```r
(b1.pval = 2*pt(-abs(b1.stat), df=n-2))
```

```
[1] 6.916425e-13
```

# Exercise answers computing manually (cont.)

```r
# confidence intervals for both coefficients
(t.crit = qt(0.975, df=n-2))
```

```
[1] 2.048407
```

```r
(b1 + c(-1,1)*t.crit*se.b1)
```

```
[1] 3.491761 4.874393
```

```r
(b0 + c(-1,1)*t.crit*se.b0)
```

```
[1] -20.5064861   0.6153899
```

```r
# confidence/prediction intervals for prediction at new X=18
(y.new = b0 + b1*18)
```

```
[1] 65.34984
```

```r
(se.conf = sigma.hat*sqrt(1/n+(18-mx)^2/(sx^2*(n-1))))
```

```
[1] 1.411564
```

```r
(se.pred = sigma.hat*sqrt(1+1/n+(18-mx)^2/(sx^2*(n-1))))
```

```
[1] 5.562983
```

```r
(y.new + c(-1,1)*t.crit*se.conf)
```

```
[1] 62.45838 68.24130
```

```r
(y.new + c(-1,1)*t.crit*se.pred)
```

```
[1] 53.95459 76.74510
```

# Verify SEs with MC

Can verify the coefficient SEs with MC:

```
# use same X vector
x = c(14.4,15.3,11.2,10.7,11.9,13,18.4,15.3,17.3,10.1,18.4,18.1,16.6,15.7,10.3,
      19,10.9,16.3,17,17.2,14.3,12.3,15.8,17.8,15.2,20,11.1,18.3,15.9,12.1)
n = length(x)

# previous y points generated with:
b1 = 4 ; b0 = -8 ; sigma = 6

# generate many new samples, fit, and extract coefficients
betas = replicate(1e4, coef(lm(b0 + b1*x + rnorm(n,0,sigma) ~ x)) )
b0s = betas[1,] ; b1s = betas[2,]
c(mean(b0s), mean(b1s))
```

```
[1] -7.986806  3.998902
```

```
# empirical SEs
c(sd(b0s), sd(b1s))
```

```
[1] 5.7415962 0.3755352
```

```
# theoretical values
sigma * sqrt(1/n + mx^2/(sx^2*(n-1)))
```

```
[1] 5.748852
```

```
sigma * sqrt(1/(sx^2*(n-1)))
```

```
[1] 0.3763182
```

# How "good" is fit?

Often how "good" a fit is can have 2 different interpretations:

- Is the model **appropriate** for the data? I.e. is the linear model assumption correct?
  - Answered by checking residual plots: are they normal, 0-mean, constant variance?

# How "good" is fit?

Often how "good" a fit is can have 2 different interpretations:


- Is the model **appropriate** for the data? I.e. is the linear model assumption correct?
  - Answered by checking residual plots: are they normal, 0-mean, constant variance?
- Alternatively, can ask how **predictive** is model? I.e. is $X$ predicting $Y$ well?
  - Often answered with **coefficient of determination** $R^2$, which $= (r)^2$ for SLR
  - Formal definition: recall data composed of "signal" and "noise" components; after fitting, **Total** variability of reponse, $\text{TSS} = \sum(y_i - \bar{y})^2$ (also SST) partitioned into:
    i. Portion explained by **Model** $\text{MSS} = \sum(\hat{y}_i - \bar{y})^2$ (also ESS or SSR)
    ii. **Residuals** portion not explained by model $\text{RSS} = \sum(y_i - \hat{y}_i)^2$ (also SSE or SSR)

# How "good" is fit?

Often how "good" a fit is can have 2 different interpretations:

- Is the model **appropriate** for the data? I.e. is the linear model assumption correct?
  - Answered by checking residual plots: are they normal, 0-mean, constant variance?
- Alternatively, can ask how **predictive** is model? I.e. is $X$ predicting $Y$ well?
  - Often answered with **coefficient of determination** $R^2$, which $= (r)^2$ for SLR
  - Formal definition: recall data composed of "signal" and "noise" components; after fitting, **Total** variability of reponse, $\text{TSS} = \sum(y_i - \bar{y})^2$ (also SST) partitioned into:
    i. Portion explained by **Model** $\text{MSS} = \sum(\hat{y}_i - \bar{y})^2$ (also ESS or SSR)
    ii. **Residuals** portion not explained by model $\text{RSS} = \sum(y_i - \hat{y}_i)^2$ (also SSE or SSR)

$$R^2 = \frac{\text{TSS} - \text{RSS}}{\text{TSS}} = \frac{\text{MSS}}{\text{TSS}}$$
$$= 1 - \frac{\text{RSS}}{\text{TSS}}$$



# $R^2$ example

For previous fit,

```
# print portion of summary table
summary(model)  %>% capture.output %>% tail(-8) %>% cat(sep="\n")
```

```
Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)  -9.9455     5.1557  -1.929   0.0639 .
x             4.1831     0.3375  12.395 6.92e-13 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.381 on 28 degrees of freedom
Multiple R-squared:  0.8458,    Adjusted R-squared:  0.8403
F-statistic: 153.6 on 1 and 28 DF,  p-value: 6.916e-13
```

```
# compute R² from formula
1-sum(resid(model)^2)/(sy^2*(n-1))
```

```
[1] 0.845839
```

```
# compare with correlation²
r^2
```

```
[1] 0.845839
```

Interpretation: about 85% of variation in Y explained by variation in X

# Multiple linear regression (MLR)

Generally speaking, MLR same as SLR but with more terms. Model equation:

$$y_i = \beta_0 + \beta_1 x_{i,1} + \beta_2 x_{i,2} + \cdots + \beta_p x_{i,p} + \varepsilon_i$$

- $\beta_0, \beta_1, ..., \beta_p$ are true intercept & slopes (still constants like all $x_{i,j}$)
- $\varepsilon_i \overset{\text{iid}}{\sim} \text{Normal}(0, \sigma^2)$ are **errors** of each point (still the only variable on RHS!)
- Still usually fit with OLS, i.e. make squares of distances from points as small as possible
- Derivation arguably simpler than SLR (details here)

# Multiple linear regression (MLR)

Generally speaking, MLR same as SLR but with more terms. Model equation:

$$y_i = \beta_0 + \beta_1 x_{i,1} + \beta_2 x_{i,2} + \cdots + \beta_p x_{i,p} + \varepsilon_i$$

- $\beta_0, \beta_1, ..., \beta_p$ are true intercept & slopes (still constants like all $x_{i,j}$)
- $\varepsilon_i \overset{\text{iid}}{\sim} \text{Normal}(0, \sigma^2)$ are **errors** of each point (still the only variable on RHS!)
- Still usually fit with OLS, i.e. make squares of distances from points as small as possible
- Derivation arguably simpler than SLR (details here)

Only some key differences in MLR:

- Degrees of freedom: $n - (p + 1)$, one for each coefficient (note $p = 1$ gives SLR's $n - 2$)
- Coefficients interpreted as expected change in $Y$ **while holding other variables constant**
  - MLR naturally attempts to attribute changes in $Y$ associated with each variable
  - Useful to account/control for other influencing variables
  - Downside: predictors should not be strongly correlated, i.e. **multicollinearity** (can use statistics like VIF to detect)

# MLR exercise

Fitting in R: simply add more variables to `y ~ var1 + var2 + ...` formula in `lm()` call

Exercise: recall `mtcars` which contains fuel efficiency data for several cars, fit an MLR model predicting `qsec` using `disp`, `hp`, and `wt`. Show summary, interpret coefficients values + significances briefly, check residuals, and find 95% confidence intervals for coefficients.

# MLR exercise

Fitting in R: simply add more variables to `y ~ var1 + var2 + ...` formula in `lm()` call

Exercise: recall `mtcars` which contains fuel efficiency data for several cars, fit an MLR model predicting `qsec` using `disp`, `hp`, and `wt`. Show summary, interpret coefficients values + significances briefly, check residuals, and find 95% confidence intervals for coefficients.

```
model.mtcars = lm(qsec ~ disp + hp + wt, data = mtcars)
summary(model.mtcars)  %>% capture.output %>% tail(-8) %>% cat(sep="\n")
```

```
Coefficients:
             Estimate Std. Error t value Pr(>|t|)
(Intercept) 17.965050   0.849663  21.144  < 2e-16 ***
disp        -0.006622   0.004166  -1.590  0.12317
hp          -0.022953   0.004603  -4.986 2.88e-05 ***
wt           1.485283   0.429172   3.461  0.00175 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.062 on 28 degrees of freedom
Multiple R-squared:  0.6808,    Adjusted R-squared:  0.6466
F-statistic: 19.91 on 3 and 28 DF,  p-value: 4.134e-07
```


Interpretation, e.g. for `disp`: for each unit increase in displacement, qsec-time expected to drop by 0.0066 on average, **holding other variables constant**. Controlling for other variables, displacement not significant at $\alpha = 0.05$ ($p = 0.12$)


# MLR exercise (cont.)

```
confint(model.mtcars)
```

```
                  2.5 %       97.5 %
(Intercept) 16.22459542 19.705504902
disp        -0.01515581  0.001911775
hp          -0.03238272 -0.013524148
wt           0.60616454  2.364401365
```

```
plot(model.mtcars, which=1:2)
```



# MLR exercise (cont.)

For MLR, $R^2$ still defined as $1 - \mathrm{RSS}/\mathrm{TSS}$, but no longer interpretable as square of a correlation coefficient, since correlation only defined for two variables.

For previous exercise, compute and interpret $R^2$ using residuals.

# MLR exercise (cont.)

For MLR, $R^2$ still defined as $1 - \mathrm{RSS}/\mathrm{TSS}$, but no longer interpretable as square of a correlation coefficient, since correlation only defined for two variables.

For previous exercise, compute and interpret $R^2$ using residuals.

```
# extract from summary output
summary(model.mtcars)$r.squared
```

```
[1] 0.6808292
```

```
# compute manually
1-sum(resid(model.mtcars)^2)/(var(mtcars$qsec)*(nrow(mtcars)-1))
```

```
[1] 0.6808292
```

Interpretation: about 68% of variation in qsec-time explained by variations in displacement, horsepower, and weight of cars.

# Overall model F-test

Another way of assessing model significance: look at F-statistic, which is defined:

$$F_{\text{obs}} = \frac{\text{MSS}/p}{\text{RSS}/(n-(p+1))}$$

and tests against the null model in the null hypothesis:

$$H_0 : \beta_1 = \beta_2 = \cdots = \beta_p = 0,$$

vs $H_a$ : at least one $\beta_i \neq 0$, i.e. tests entire model useless vs not useless.

# Overall model F-test

Another way of assessing model significance: look at F-statistic, which is defined:

$$F_{\text{obs}} = \frac{\text{MSS}/p}{\text{RSS}/(n-(p+1))}$$

and tests against the null model in the null hypothesis:

$$H_0 : \beta_1 = \beta_2 = \cdots = \beta_p = 0,$$

vs $H_a$ : at least one $\beta_i \neq 0$, i.e. tests entire model useless vs not useless.


F-statistic follows F-distribution with $p, n-(p+1)$ DFs. For 340, just interpreting output of summary is sufficient:

```
summary(model.mtcars)  %>% capture.output %>% tail(4) %>% cat(sep="\n")
```

```
Residual standard error: 1.062 on 28 degrees of freedom
Multiple R-squared:  0.6808,    Adjusted R-squared:  0.6466
F-statistic: 19.91 on 3 and 28 DF,  p-value: 4.134e-07
```

$F = 19.91$ on 3, 28 DF, $p = 4.1 \times 10^{-7} \ll 0.05$, very significant vs null model

# $F$-distribution examples



# Additional predictor types

In MLR, there are often many other types of predictors we can consider adding to the model. We will examine 3 main types:

- Categorical variables
  - Examples: Control/treatment, male/female, high/medium/low, etc.
- Interactions
  - When two variables (numeric or categorical) have additional combined effects, i.e. “synergy”
- Non-linear terms
  - When a function of a variable is better modeled as linear vs response

Additional predictors can be added to model with same syntax `y ~ var1 + var2 + ...`

# Categorical predictors

For categorical predictors with 2 or more **levels** (or categories), the following is done:

1. One level is **taken as the reference leve**, (or baseline),
2. The difference between **every other level *vs the reference*** is then individually modeled as a **slope on an indicator variable** (i.e. 1/0) for that level,
3. These slopes are shown as coefficients (note: the **reference is NOT shown** as a slope!)

# Categorical predictors

For categorical predictors with 2 or more **levels** (or categories), the following is done:

1. One level is **taken as the reference leve**, (or baseline),
2. The difference between **every other level *vs the reference*** is then individually modeled as a **slope on an indicator variable** (i.e. 1/0) for that level,
3. These slopes are shown as coefficients (note: the **reference is NOT shown** as a slope!)

Example: for `iris` dataset below, `Species` is categorical with 3 levels:

```
str(iris)
```

```
'data.frame':   150 obs. of  5 variables:
 $ Sepal.Length: num  5.1 4.9 4.7 4.6 5 5.4 4.6 5 4.4 4.9 ...
 $ Sepal.Width : num  3.5 3 3.2 3.1 3.6 3.9 3.4 3.4 2.9 3.1 ...
 $ Petal.Length: num  1.4 1.4 1.3 1.5 1.4 1.7 1.4 1.5 1.4 1.5 ...
 $ Petal.Width : num  0.2 0.2 0.2 0.2 0.2 0.4 0.3 0.2 0.2 0.1 ...
 $ Species     : Factor w/ 3 levels "setosa","versicolor",..: 1 1 1 1 1 1 1 1 1 1 ...
```

```
# model predicting Sepal.Length from all other variables:
# alternative: lm(Sepal.Length ~ . , data=iris) where . means all other columns
model.iris = lm(Sepal.Length ~ Sepal.Width + Petal.Length + Petal.Width + Species, data=iris)
```

# Iris example (cont.)

```
summary(model.iris)
```

```
Call:
lm(formula = Sepal.Length ~ Sepal.Width + Petal.Length + Petal.Width
+
    Species, data = iris)

Residuals:
     Min       1Q   Median       3Q      Max
-0.79424 -0.21874  0.00899  0.20255  0.73103

Coefficients:
                   Estimate Std. Error t value Pr(>|t|)
(Intercept)         2.17127    0.27979   7.760 1.43e-12 ***
Sepal.Width         0.49589    0.08607   5.761 4.87e-08 ***
Petal.Length        0.82924    0.06853  12.101  < 2e-16 ***
Petal.Width        -0.31516    0.15120  -2.084  0.03889 *
Speciesversicolor -0.72356    0.24017  -3.013  0.00306 **
Speciesvirginica  -1.02350    0.33373  -3.067  0.00258 **
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.3068 on 144 degrees of freedom
Multiple R-squared:  0.8673,    Adjusted R-squared:  0.8627
F-statistic: 188.3 on 5 and 144 DF,  p-value: < 2.2e-16
```

- Note: setosa used as reference (not shown), other species vs setosa fitted as coefficients
- I.e. **all else equal**, versicolor sepal length lower vs setosa by 0.72cm **on average**
- Both species **significantly different** vs setosa (note: versicolor vs virginica difference NOT modeled & may not be significant)
- To change reference, mutate variable with `fct_relevel()`

  (*demo*)

# Categorical variables (cont.)

Sometimes, categorical variables may already be coded as 1/0 indicator variables, in this case add them in as normal and the coefficient shown is the **difference of 1 vs 0 levels**. Example with `mtcars` again:

```
# update previous mtcars model by adding am, vs categorical variables
# alternative: model.mtcars2 = update(model.mtcars, . ~ . + am + vs)
model.mtcars2 = lm(qsec ~ disp + hp + wt + am + vs, data = mtcars)
summary(model.mtcars2)  %>% capture.output %>% tail(-8) %>% cat(sep="\n")
```

```
Coefficients:
             Estimate Std. Error t value Pr(>|t|)
(Intercept) 17.237081   1.083732  15.905 6.49e-15 ***
disp        -0.003235   0.003509  -0.922 0.364987
hp          -0.014683   0.003945  -3.722 0.000961 ***
wt           0.949956   0.358770   2.648 0.013584 *
am          -0.802919   0.462681  -1.735 0.094517 .
vs           1.787328   0.463080   3.860 0.000673 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.799 on 26 degrees of freedom
Multiple R-squared:  0.8323,    Adjusted R-squared:  0.8001
F-statistic: 25.81 on 5 and 26 DF,  p-value: 2.6e-09
```

I.e. **all else equal**, manual transmission faster vs automatic in qsec time by 0.8s **on average**, but this observed difference is only **weakly significant** ($p \approx 0.09$).

# Practice exercise

```
str(penguins)
```

```
'data.frame':   344 obs. of  8 variables:
 $ species    : Factor w/ 3 levels "Adelie","Chinstrap",..: 1 1 1 1 1 1 1 1 1 1 ...
 $ island     : Factor w/ 3 levels "Biscoe","Dream",..: 3 3 3 3 3 3 3 3 3 3 ...
 $ bill_len   : num  39.1 39.5 40.3 NA 36.7 39.3 38.9 39.2 34.1 42 ...
 $ bill_dep   : num  18.7 17.4 18 NA 19.3 20.6 17.8 19.6 18.1 20.2 ...
 $ flipper_len: int  181 186 195 NA 193 190 181 195 193 190 ...
 $ body_mass  : int  3750 3800 3250 NA 3450 3650 3625 4675 3475 4250 ...
 $ sex        : Factor w/ 2 levels "female","male": 2 1 1 NA 1 2 1 2 NA NA ...
 $ year       : int  2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 ...
```

With the build-in `penguins` dataset (shown above), do the following:

1. Use species, bill length, bill depth to linearly predict flipper length.
2. Give a full & clear interpretation of **each coefficient** (both slope & significance).
3. Is the overall model significant? Give a statistic & p-value.
4. What amount of the variation in flipper length is explained by your model?
5. Predict flipper length for an Adelie penguin with 40mm bill length and 20mm bill depth.
6. Run model diagnostics (i.e. check residual assumptions).

# Penguins exercise answers

```r
model.penguins = lm(flipper_len ~ species + bill_len + bill_dep, data=penguins)
summary(model.penguins)  %>% capture.output %>% tail(-8) %>% cat(sep="\n")
```

```
Coefficients:
                 Estimate Std. Error t value Pr(>|t|)
(Intercept)      127.6915     5.1937  24.586  < 2e-16 ***
speciesChinstrap  -1.3062     1.4391  -0.908    0.365
speciesGentoo     27.5648     1.9606  14.060  < 2e-16 ***
bill_len           0.7004     0.1197   5.851 1.16e-08 ***
bill_dep           1.9127     0.3162   6.050 3.85e-09 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.541 on 337 degrees of freedom
  (2 observations deleted due to missingness)
Multiple R-squared:  0.8465,    Adjusted R-squared:  0.8447
F-statistic: 464.7 on 4 and 337 DF,  p-value: < 2.2e-16
```

- Bill length found to be **highly significant** ($p \approx 10^{-8}$). **All else equal**, each 1mm increase in bill length associated with 0.70mm increase in flipper length **on average**.
- **All else equal**, Gentoo penguins' flipper lengths found to be **highly significantly** different vs Adelies ($p < 10^{-15}$) averaging 27.6mm longer, but Chinstraps not found to be significant ($p \approx 0.37$) averaging only -1.3mm shorter.

# Penguins exercise answers (cont.)

```r
model.penguins = lm(flipper_len ~ species + bill_len + bill_dep, data=penguins)
summary(model.penguins)  %>% capture.output %>% tail(-8) %>% cat(sep="\n")
```

```
Coefficients:
                 Estimate Std. Error t value Pr(>|t|)
(Intercept)      127.6915     5.1937  24.586  < 2e-16 ***
speciesChinstrap  -1.3062     1.4391  -0.908    0.365
speciesGentoo     27.5648     1.9606  14.060  < 2e-16 ***
bill_len           0.7004     0.1197   5.851 1.16e-08 ***
bill_dep           1.9127     0.3162   6.050 3.85e-09 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.541 on 337 degrees of freedom
  (2 observations deleted due to missingness)
Multiple R-squared:  0.8465,    Adjusted R-squared:  0.8447
F-statistic: 464.7 on 4 and 337 DF,  p-value: < 2.2e-16
```

- Overall model highly significant vs null ($F = 464.7$ on 4, 337 DFs, $p < 10^{-15}$).
- About 85% of variation in flipper length explained by model.

# Penguins exercise answers (cont.)

```r
# prediction using predict() function
predict(model.penguins, tibble(species="Adelie",bill_len=40,bill_dep=20))
```

```
       1
193.9631
```


```r
# manual prediction
coef(model.penguins)
```

```
     (Intercept) speciesChinstrap    speciesGentoo         bill_len         bill_dep
     127.6914684       -1.3062319       27.5648475        0.7004416        1.9126995
```

```r
sum(coef(model.penguins) * c(1, 0, 0, 40, 20))
```

```
[1] 193.9631
```

```r
plot(model.penguins, which=1:2)
```



*(more notes added soon)*