

SIMPLE CONNECTIVITY OF THE LIDMAN–PICCIRILLO PIECE: THE COMPUTATION, STEP BY STEP

BERND JOHANNES WUEBBEN

ABSTRACT. This is an expository companion to Part II, the computation, of the manuscript [M] (*An exotic $S^2 \times S^2$ and an exotic $\mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$*). It re-derives, at a much finer grain than a journal format permits, the geometric and group-theoretic inputs of the relation system used there to prove $\pi_1(V) = 1$ for the specified piece $V = V'_{0,0}$: the marked surface model and its conventions, all three corrected transport relations worked end to end, the meridian basings, all four Lagrangian-framing direction words (including the pushoff-basing correction and the monodromy-drift word), the drilled-fiber relation, the epimorphism onto $\pi_1(V)$, and the assembled relation sheet. It closes with the Baldrige–Kirk T^4 configuration as a fully worked calibration against a published group family, using finite fingerprints rather than an asserted presentation isomorphism, and with a protocol for diffing an independently derived relation sheet against ours. Nothing here is new relative to [M]; the point is that every step be checkable in writing, with no gap a reader must bridge alone.

CONTENTS

1. How to read this document	1
2. Conventions, all of them	2
3. The fiber octagon and the developing engine	2
4. The cut-square model	3
5. Membranes: the mechanism, slowly	4
6. The three corrected relations, worked end to end	5
7. Meridians, fillings, and the four direction words	6
8. The drilled-fiber relation and the surjection onto $\pi_1(V)$	8
9. The assembled relation system for the fixed piece	9
10. The same method on Baldrige–Kirk’s T^4 : a fully worked calibration	10
11. How to diff a relation sheet against ours	11

1. HOW TO READ THIS DOCUMENT

The manuscript [M] separates geometric arguments from reproducible computations. This document follows the same division at tutorial speed. It takes one route through the derivation, in the order in which the objects are needed, and works every corrected relation and every direction word explicitly. Each claim of a crossing count or a derived word carries a pointer to its geometric argument or engine record in [M] and its ancillary files

Date: August 16, 2026.

2020 Mathematics Subject Classification. 57K40 (primary); 57M05, 20F05, 20F10, 20-08 (secondary).

Key words and phrases. Fundamental group, surface bundles, Luttinger surgery, monodromy relations, relation systems, verification.

(all public, together with every harness and run log, at <https://github.com/bwuebben/exotic-s2xs2>); the reader who wants to distrust a step can follow the pointer, and the reader who only wants to *understand* the step should not need to.

The theorem concerns one explicitly parametrized manifold: throughout, $V = V'_{0,0}$ is the result of the two Lidman–Piccirillo +1 surgeries with the base-direction push offs derived below. The other eight exponent pairs that occur in the run records are formal diagnostics. No fundamental-group claim for another parametrization is made here.

Throughout, [M] is the manuscript *An exotic $S^2 \times S^2$ and an exotic $\mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$* (section numbers refer to the version distributed with this companion; Part I is the reduction, Part II the computation), and [BK] Baldridge and Kirk, *Constructions of small symplectic 4-manifolds using Luttinger surgery*, J. Differential Geom. 82 (2009) 317–361, arXiv:math/0703065.

2. CONVENTIONS, ALL OF THEM

Convention 2.1.

- (1) **Composition.** Paths and loops compose left to right: $\gamma_1\gamma_2$ traverses γ_1 first. A relation written $ugu^{-1} = w$ names the relator $ugu^{-1}w^{-1}$.
- (2) **Fiber generators.** x, y, r, s are the edge loops of the rotation-symmetric octagon model of the genus-2 fiber F , based at the vertex point p , with the surface relator $[x, y][r, s]$. In Lidman–Piccirillo’s curve names, y realizes the class of b and s the class of d .
- (3) **Monodromies.** $h := T_a \circ T_b: x \mapsto y^{-1}, y \mapsto yx$ is the trefoil monodromy on the left handle (machine-certified: it fixes $[x, y]$ exactly, h^6 is the boundary twist, $h^3 = -\text{id}$ on H_1 ; [M] §3.1). The bundle monodromies are $\tilde{\psi} := h * \text{id}$ (so ψ_0 fixes the *right* handle, in particular the curves e and s , pointwise) and $\tilde{\varphi} :=$ the handle swap ($x \leftrightarrow r, y \leftrightarrow s$).
- (4) **Base generators.** $\bar{\alpha}, \bar{\beta}$ are the based base loops of Section 4; the relation system’s generators are $A := [\bar{\alpha}]^{-1}$ and $B := [\bar{\beta}]^{-1}$. With the left-to-right composition rule this makes conjugation by B implement one *upward* transport around $\bar{\beta}$:

$$BgB^{-1} = \tilde{\psi}(g) \cdot [\text{correction}] \quad \text{for a fiber loop } g,$$

and likewise A with $\tilde{\varphi}$. The opposite choices are the ε -flips of [M] Remark 3.7; they are swept, but every displayed word below uses the stated convention.

- (5) **Checked signs.** Five signs enter the displayed relation sheet: $\varepsilon_3, \varepsilon_4, \varepsilon_5$ (the meridian signs of the three corrected relations) and $\varepsilon_A, \varepsilon_B$ (the filling exponents). The fixed- V computation checks every assignment as part of the 4,096-case robustness family; no conclusion below depends on resolving them.
- (6) **Meridians.** M and N denote the meridians of the surgery tori T_α and T_β , *based along specific whiskers* fixed in Section 6; “meridian” always means this based loop, not a free-homotopy class.

3. THE FIBER OCTAGON AND THE DEVELOPING ENGINE

There is a geometric step before any word is read from the octagon. The five curves a, b, c, d, e in Lidman–Piccirillo’s marked surface form an ordered five-chain whose regular neighborhood fills the genus-2 surface up to two disks. An isomorphism from its oriented ribbon graph to the five-chain in the octagon therefore extends over the surface. It may be chosen equivariantly with respect to the involution that reverses the chain, and the

extensions over the two complementary disks send its fixed points to the octagon vertex p and center O . This is [M] Lemma 7.1. Thus the polygonal curves below are ambient representatives of the Lidman–Piccirillo curves, not merely curves with the same homology classes.

After that identification is fixed, each curve and path is specified by a finite combinatorial description: the corner wedge V_i it departs from, the ordered list of octagon edges it crosses, and the arrival wedge V_j . A small engine (`develop.py`, ancillary) converts that description into a based word by developing it in the universal cover and reducing with Dehn’s algorithm; [M] Appendix A.1 specifies the five-line algorithm exactly. Its validations V1–V5 compare the calculation with known answers (edge-parallels return their generators; the vertex-link loop develops to 1; the pushoff descriptions return $a \simeq x^{-1}$, $b \simeq y$, $e \simeq r^{-1}$, $d \simeq s$ swap-equivariantly; the two lasso arcs differ by a whiskered copy of c ; the frozen chord coordinates realize the intersection table and place the κ_3 path off c). The engine verifies the word associated with a stated description. It does not supply the marked-surface identification or choose the source curve.

Three engine outputs are used constantly below, so we fix them now (run record `develop_out.txt`, blocks D3–D5):

- the invariant curve c has based word $(rx)^{-1}$ (at its V_2 -whisker), and the curve e develops to r^{-1} (at V_7);
- *position table, s-edge*: traversing s from p , the crossing with c (call it c_s) comes *before* the crossing with e (call it s_e). This ordering is forced by disjointness: $c \cap e = \emptyset$, and e ’s corner arc meets the s -edge near its end. (It is also checked mechanically: the engine’s chord-diagram test enumerates all candidate orderings and shows this one is the unique planar realization of the model’s intersection pattern.) Write s_1 for the initial segment of s up to c_s , and s_2 for the initial segment up to s_e ; thus $s_1 \subset s_2$.
- *position table, y-edge*: y crosses c once, at c_y ; write y_1 for the initial segment of y up to c_y . The loops x and r cross neither c nor e .

4. THE CUT-SQUARE MODEL

The base T_0 (once-punctured torus) is the square $[0, 1]^2$ in coordinates (u_1, u_2) , minus a puncture disk O at $(3/4, 3/4)$, with the two edge identifications

$$(\xi, (t, 1)) \sim (\psi_0 \xi, (t, 0)), \quad (\xi, (1, u)) \sim (\varphi_0 \xi, (0, u))$$

on the total space $F \times [0, 1]^2 / \sim$. In words: the horizontal edge pair $\{u_2 = 0 \equiv 1\}$ is the α -cut, and crossing it *upward* applies ψ_0 to the fiber; the vertical edge pair $\{u_1 = 0 \equiv 1\}$ is the β -cut, and crossing it *rightward* applies φ_0 .

The basepoint is (p, q) with $q = (1/4, 1/4)$. The based base loops are $\bar{\beta}$ = the vertical circle through q (travel upward from q , wrap at $u_2 = 1 \equiv 0$, return to q ; it crosses the α -cut exactly once and the β -cut never) and $\bar{\alpha}$ = the horizontal circle through q (crosses the β -cut exactly once, the α -cut never). These realize the minimal possible crossing pattern.

The surgery tori sit *over the cuts*:

$$T_\alpha = c \times \{\alpha\text{-cut circle}\}, \quad T_\beta = e \times \{\beta\text{-cut circle}\}.$$

Why these are well-defined tori: traveling once along the α -cut circle (a horizontal circle) crosses the β -cut once, applying φ_0 , and $\varphi_0(c) = c$ (setwise: the restriction $\varphi_0|_c$ is a free half-rotation of the circle c , a fact that returns in Section 7); traveling once along the β -cut

circle crosses the α -cut once, applying ψ_0 , and $\psi_0(e) = e$ *pointwise*, so T_β even inherits the product framing. Note the consequence used constantly below:

a loop or membrane can meet T_α only over the α -cut, and only at fiber points on c ; it can meet T_β only over the β -cut, and only at fiber points on e .




FIGURE 1. *Illustration only; every crossing count in the text is a position-table or recorded combinatorial fact.* Left: the cut square. Right: the membrane square of Section 5.

5. MEMBRANES: THE MECHANISM, SLOWLY

Fix a fiber loop g based at p , and transport it once around $\bar{\beta}$. This sweeps the map of a square

$$\mathfrak{M}_g(u, v) = (\text{transport of } g(u) \text{ along } \bar{\beta} \text{ to time } v), \quad (u, v) \in [0, 1]^2,$$

whose four sides are: bottom = g ; top = $\tilde{\psi}(g)$ (the loop after one full transport); left and right = $\bar{\beta}$ itself, the track of the basepoint p (the monodromies fix p , so that track closes). Reading the boundary of the square from its base corner gives the identity

$$g \cdot \bar{\beta} \cdot \tilde{\psi}(g)^{-1} \cdot \bar{\beta}^{-1} = 1 \quad \text{in } \pi_1(R),$$

which after substituting $B = [\bar{\beta}]^{-1}$ is exactly the monodromy relation $BgB^{-1} = \tilde{\psi}(g)$. *The square is the proof of the relation:* everything that happens to the relation after surgery happens to this square.

Now remove the surgery tori. If the square's image is disjoint from $T_\alpha \cup T_\beta$, the relation survives verbatim in the complement. If the square meets a torus transversally in interior points $z_1, \dots, z_k$, puncture it there: the punctured square exhibits the boundary word as a product of small loops around the punctures, each conjugated by an in-square path from the base corner. Each small loop maps to a meridian of the torus it surrounds. Hence in the complement

$$BgB^{-1} = \left(\prod_i \ell_i \mu_i^{\pm 1} \ell_i^{-1} \right) \cdot \tilde{\psi}(g),$$

one conjugated meridian per crossing, where ℓ_i is the in-square path to the i -th puncture. Two facts keep this bookkeeping tame:

- (1) *The domain is a disk*, so the choice of in-square route to a puncture changes $\ell_i \mu^{\pm 1} \ell_i^{-1}$ by at most powers of the same meridian, i.e., not at all. (This is why the membrane calculus is stable; the only genuinely non-obvious lasso appears when a meridian must be *re-based* to a different whisker, Section 6.)
- (2) *The crossings are computable*. By the boxed principle of Section 4, the square $\mathfrak{M}_g$ meets T_α only where its base track crosses the α -cut (for $\bar{\beta}$, at the single wrap time $v = 3/4$), and its fiber point lies on c : i.e., at the points $(g \cap c) \times \{\text{wrap}\}$. Likewise for T_β with the β -cut and e . The crossing count of any membrane is therefore (number of base-track crossings with the cut) $\times$ (number of fiber crossings of g with the torus' fiber curve), both entries read off the position table.

6. THE THREE CORRECTED RELATIONS, WORKED END TO END

There are eight monodromy relations: x, y, r, s over $\bar{\alpha}$ and over $\bar{\beta}$. We now compute all eight crossing counts and derive the corrected forms. The outcome, stated first:

$$AsA^{-1} = N^{\varepsilon_3} y, \quad ByB^{-1} = M^{\varepsilon_4} (yx), \quad BsB^{-1} = (\delta M^{\varepsilon_5} \delta^{-1}) s \quad \text{with } \delta = r^{-1},$$

and the other five relations are clean.

6.1. y over $\bar{\beta}$ (the template). Base track: $\bar{\beta}$ crosses the α -cut once (the wrap), the β -cut never $\Rightarrow$ only T_α is possible. Fiber crossings: $y \cap c = \{c_y\}$, one point. So $\mathfrak{M}_y$ meets T_α in the single transverse point $z = (c_y, \text{wrap})$ and meets T_β not at all. Puncture there. The in-square lasso from the base corner is: along the bottom to $u(c_y)$ (that is the partial loop y_1), then vertically up to the wrap. Now *define* the based meridian of T_α along that very whisker:

$$M := y_1 \cdot (\text{meridian circle of } T_\alpha \text{ at } c_y) \cdot y_1^{-1}.$$

With this basing the conjugated-meridian factor is $M^{\pm 1}$ (the lasso has been absorbed into the definition), and since $\tilde{\psi}(y) = h(y) = yx$:

$$ByB^{-1} = M^{\varepsilon_4} (yx).$$

The sign ε_4 encodes the membrane-orientation convention; it is swept.

6.2. s over $\bar{\beta}$ (the one genuine lasso). Same base track, so again only T_α , at the wrap. Fiber crossings: $s \cap c = \{c_s\}$, one point. So one puncture, at $z_0 = (c_s, \text{wrap})$, and $\tilde{\psi}(s) = s$ (ψ_0 fixes the right handle pointwise). The in-square lasso is $s_1 \cdot$ (vertical transport). But M is based at c_y , not at c_s : to express the correction in terms of M , the meridian must be slid *along the torus* from the crossing to M 's basing point: concretely, along the curve c from c_s to c_y . The slide path, developed by the engine, is

$$\delta = s_1 \cdot (\text{arc of } c \text{ from } c_s \text{ to } c_y) \cdot y_1^{-1} = r^{-1}$$

(engine record D3). Hence

$$BsB^{-1} = (\delta M^{\varepsilon_5} \delta^{-1}) s, \quad \delta = r^{-1}.$$

Two sanity points. First, c has two arcs from c_s to c_y ; the other gives $\delta' = x$, and the engine verifies the consistency identity (V4) $\delta' \delta^{-1} = xr \sim$ a whiskered copy of c , precisely the “slid all the way around” discrepancy one should find. In the group the choice is immaterial: conjugating M by a whiskered c is conjugation by a pushoff class on the boundary 3-torus $\partial \nu(T_\alpha)$, which commutes with the meridian there. The harness fixes $\delta = r^{-1}$. Second,

the placement of the correction (left of $\tilde{\psi}(g)$) is an orientation convention; [M]’s 256-case placement check, with R_3 included, gives a trivial relation group in every case.

6.3. s **over** $\bar{\alpha}$. Base track: $\bar{\alpha}$ crosses the β -cut once, the α -cut never $\Rightarrow$ only T_β is possible. Fiber crossings: $s \cap e = \{s_e\}$, one point. One puncture; $\tilde{\varphi}(s) = y$. The in-square lasso is $s_2 \cdot$ (horizontal transport), and defining

$$N := s_2 \cdot (\text{meridian circle of } T_\beta \text{ at } s_e) \cdot s_2^{-1}$$

absorbs it, exactly as in §6.1:

$$AsA^{-1} = N^{\varepsilon_3} y.$$

6.4. **The five clean relations.** Over $\bar{\beta}$, only T_α is reachable and the fiber condition is “crosses c ”: x and r do not, so $BxB^{-1} = y^{-1}$ and $BrB^{-1} = r$ hold verbatim. Over $\bar{\alpha}$, only T_β is reachable and the condition is “crosses e ”: x, y, r do not, so $AxA^{-1} = r$, $AyA^{-1} = s$, $ArA^{-1} = x$ hold verbatim. That is the complete count: three corrections, each at one crossing, five clean relations, and every entry of the count is a position-table fact, not a picture-reading.

7. MERIDIANS, FILLINGS, AND THE FOUR DIRECTION WORDS

A $(1/k)$ -Luttinger filling along a curve γ on a surgery torus T kills the class $\mu \cdot (\text{pushoff of } \gamma)^k$, where (this is [BK]’s explicitly stated discipline, p. 3) *the meridian μ and the Lagrangian-framed pushoff must be based through the same whisker*. Our fillings ($k = \pm 1$ swept as $\varepsilon_A, \varepsilon_B$; directions $\alpha'c^n$ on T_α and $\beta'e^m$ on T_β , $m, n \in \{-1, 0, 1\}$ in the diagnostic grid) therefore need four *direction words*: the based pushoffs of the fiber and base directions of each torus, at the same whiskers that base M and N . The fixed manifold $V = V'_{0,0}$ uses the two base directions; the fiber directions parameterize the nearby formal systems and check the framing calculation. The four derivations, in increasing order of subtlety:

The identification of these topological push offs with the Lagrangian-framing push offs is geometric. It follows from the Weinstein neighborhood models and the two Moser normalizations proved in [M] Section 8.7; the development engine does not establish that framing identification.

7.1. **Fiber direction of T_β .** The pushoff of e at the s_2 -basing. The engine develops e (departing V_7) to r^{-1} ; re-whiskering to the s_2 -basing conjugates by s :

$$\text{dir}_{T_\beta}^{\text{fib}} = s r^{-1} s^{-1}.$$

(The in-fiber annulus pushoff is the Lagrangian framing here; no base motion, hence no crossing issues.)

7.2. **Fiber direction of T_α .** The pushoff of c at the y_1 -basing: the engine’s derived word for c ,

$$\text{dir}_{T_\alpha}^{\text{fib}} = (rx)^{-1}.$$

7.3. Base direction of T_α (the drift word). This is the input with no analogue in any trivial-monodromy calibration. A Seifert comparison on $\text{MT}(\varphi_0) \times S^1$, surgered along the drift section itself, tests it against independently presented graph-manifold groups ([M] Appendix B.2; Appendix A.3). The comparison uses abelianization and low-index subgroup fingerprints; it distinguishes the coherent arc pairing from the alternatives but is not a proof that two finitely presented groups are isomorphic. The base-direction curve of T_α travels once along the α -cut circle. But T_α 's natural parametrization is not a product: going once around the cut circle applies $\varphi_0|_c$, a *free half-rotation* of c . A closed curve on T_α in the base direction must therefore *drift along half of c* to close up. Following the based-pushoff formula with whisker $\sigma = y_1 c_1$: the transported whisker is $\tilde{\varphi}(y_1)$, and the two identities

$$\varphi_0(c_y) = c_s, \quad \tilde{\varphi}(y_1) = s_1$$

(the swap carries the y -edge data to the s -edge data) make the fiber part of the closed-up curve depends on which half-arc closes the section. For the y_1 -based parametrization of the specified piece, choose the x -arc. The word r^{-1} remains the transport conjugator in the Bs -correction of §6.2; it is not the closing arc of this section. Thus

$$\text{dir}_{T_\alpha}^{\text{base}} = \lambda = Ax.$$

Check 7.1 (The drift identity). Squaring must give the full rotation. The clean relation $Ax A^{-1} = r$ gives the exact identity

$$\lambda^2 = (Ax)^2 = r A^2 x.$$

With the Part I source orientation $[c] = [x] + [r]$, its abelianization is $2\lambda = 2[A] + [c]$. The developed fiber-direction word $(rx)^{-1}$ has class $-[c]$: the based orientation used for the Part II exponent n is opposite to the Part I homology orientation. Since n ranges over all integers, that orientation choice does not change the family.

The other minimal representative satisfies

$$Ar^{-1} = Ax(rx)^{-1}.$$

It is therefore the adjacent $n = 1$ member, not the reference $n = 0$ member. The side-resolved Seifert comparison corroborates the full dictionary: the y_1 side pairs the transport conjugator r^{-1} with the push-off label Ax , whereas the y_2 side pairs x with Ar^{-1} . The section used here is fixed by the explicit y_1 -side geometric choice, not by that finite-fingerprint comparison.

7.4. Base direction of T_β (the pushoff-basing correction). The base-direction curve of T_β is the β -cut circle at fiber s_e , pushed off; its whisker (shared with N) is s_2 . Two homotopies are needed to express the based pushoff in the generators, and they have different crossing behavior: this distinction is the content of the correction, and the place where the first version of [M] erred.

Homotopy 1 (the column slide) — clean. Slide the pushoff circle from the cut column to $\tilde{\beta}$'s column $u_1 = 1/4$, at the constant fiber point s_e . The swept annulus lies at fiber s_e throughout; it could meet T_α only if $s_e \in c$, and $c \cap e = \emptyset$; it stays off the β -cut, so it misses T_β too. Clean. (This is the step the original justification “avoids T_α since $c \cap e = \emptyset$ ” correctly covered, but it is not the only step.)

Homotopy 2 (the whisker transport) — one crossing. What remains is the loop $L = s_2 \cdot V_{s_e} \cdot s_2^{-1}$, where V_{s_e} is the vertical circle at fiber s_e . To express it via B , transport the whisker s_2 once around $\tilde{\beta}$: the membrane is $\mathfrak{A} = \mathfrak{M}_{s_2}$, the restriction of the Bs -membrane of §6.2 to $u \leq u(s_e)$, a square with sides s_2 , V_{s_e} , $\tilde{\psi}(s_2) = s_2$ (pointwise), $\tilde{\beta}$. Its crossing

count, by the same two-factor principle: it misses T_β (base column 1/4; interior fiber points $s_2(u)$, $u < u(s_e)$, miss e); it meets T_α at (fiber points of $s_2 \cap c$) $\times$ (the wrap). And here the position table bites: *the c -crossing precedes the e -crossing on the s -edge*, so $c_s \in s_2$, and the membrane meets T_α in exactly the point $z_0 = (c_s, \text{wrap})$ of §6.2: the same point, with the same local data, because $\mathfrak{A}$ is an oriented restriction of $\mathfrak{M}_s$. The local intersection number is ± 1 , so no interior homotopy avoids it.

Puncture and compute, exactly as before. The punctured square gives

$$s_2 V_{s_e} s_2^{-1} \bar{\beta}^{-1} = \ell \mu^{\varepsilon_0} \ell^{-1},$$

with ℓ the in-square lasso and ε_0 the same crossing sign that the Bs -membrane assigns to z_0 , i.e., ε_0 is the sweep parameter ε_5 . The established Bs -reduction (§6.2) says precisely that $B \ell \mu^k \ell^{-1} B^{-1} = \delta M^k \delta^{-1}$ for all k . Therefore

$$L = \ell \mu^{\varepsilon_5} \ell^{-1} \cdot \bar{\beta} \implies L^{-1} = B \cdot (\ell \mu^{-\varepsilon_5} \ell^{-1}) = B \cdot B^{-1} (\delta M^{-\varepsilon_5} \delta^{-1}) B = (\delta M^{-\varepsilon_5} \delta^{-1}) B,$$

and since the harness's direction convention has the uncorrected value B (that is, the pushoff circle traversed as $\bar{\beta}^{-1}$), the correct based word is

$$\boxed{\text{dir}_{T_\beta}^{\text{base}} = (\delta M^{-\varepsilon_5} \delta^{-1}) B, \quad \delta = r^{-1}.}$$

Note what did and did not happen: the B -conjugation cancelled exactly, so the correction sits on the left with the *plain* lasso δ ; and the meridian sign came out *anti-coupled* to the Bs -correction sign, a relative statement between two corrections at the same geometric crossing, hence covariant under the global orientation flip that the sign sweep blankets. No new swept convention was introduced. The correction abelianizes to zero, so all homology bookkeeping ([M] Lemma 4.2) is untouched. Every decision was rerun with this word. In addition, all eight candidate resolutions (both arcs, both signs, both placements) were checked at the fixed surgery cell before the exact derivation was pinned down; with the five relation-sheet signs this gives 256 systems, each trivial after R_3 is included. The simple-connectivity proof for the specified V therefore does not hinge on that local sign-and-placement choice, although the displayed relation must still record the correct geometric derivation ([M] Sections 8.5 and 11).

8. THE DRILLED-FIBER RELATION AND THE SURJECTION ONTO $\pi_1(V)$

One more clean transport relation is needed. It comes from an explicit basis of the drilled fiber, not from an exact presentation of the torus complement. Cutting the octagon along the single chord of e gives two disks; contracting a spanning tree in each disk leaves a three-petal rose represented by x, y, r . Thus

$$\pi_1(F \setminus \nu e, p) = F_3 \langle x, y, r \rangle.$$

Cutting along the two disjoint chords of c gives three disks. Contracting a tree in each of the upper, middle, and lower regions, while retaining the x - and r -edge pairs and one arc across the middle region, gives

$$\pi_1(F \setminus \nu c, p) = F_3 \langle x, r, \kappa_3 \rangle, \quad \kappa_3 = s^{-1} r^{-1} y x.$$

The middle-region representative of κ_3 runs from the V_7 wedge to the V_3 wedge and is geometrically disjoint from both c -chords. This tree collapse is the proof of the basis statement. Engine record D5 independently checks the frozen coordinates, the disjointness, and the displayed based word.

The transport annulus of κ_3 around B misses T_α because $\kappa_3 \cap c = \emptyset$, and misses T_β because $\bar{\beta}$ does not cross the β -cut. It therefore supplies the true relation

$$R_3: \quad B \kappa_3 B^{-1} = \tilde{\psi}(\kappa_3) = r^{-1} s^{-1} x.$$

For the last equality, $\tilde{\psi}(\kappa_3) = s^{-1} r^{-1} y x y^{-1}$, and the surface relation $[x, y][r, s] = 1$ reduces this word to $r^{-1} s^{-1} x$.

Now let G be the group on x, y, r, s, A, B, M, N defined by the relations in Section 9, for the fixed manifold $V = V'_{0,0}$ and a fixed assignment of the five signs. There is a natural epimorphism

$$G \twoheadrightarrow \pi_1(V).$$

Indeed, put $C = R \setminus (\nu T_\alpha \cup \nu T_\beta)$. General position makes every loop in R disjoint from the two codimension-two tori, so $\pi_1(C) \rightarrow \pi_1(R)$ is onto. The kernel is normally generated by the two meridians: make a null-homotopy transverse to the tori and remove small disks around its intersection points. Hence the six lifts x, y, r, s, A, B together with M, N generate $\pi_1(C)$. Attaching the two filling pieces gives a surjection $\pi_1(C) \rightarrow \pi_1(V)$ and kills the two filling slopes. The surface relation, the clean and corrected transport relations, the filling relations, and R_3 all hold by the geometric derivations above. The free group map on the eight generators therefore factors through G , proving the epimorphism.

No isomorphism $G \cong \pi_1(V)$ is asserted or needed. Since a quotient of the trivial group is trivial, $G = 1$ proves $\pi_1(V) = 1$. A nontrivial computation of G alone would be inconclusive; a nontriviality claim about the manifold group would require a separate proof that its proposed presentation is exact.

9. THE ASSEMBLED RELATION SYSTEM FOR THE FIXED PIECE

At $(m, n) = (0, 0)$, the group G has generators x, y, r, s, A, B, M, N and defining relations:

	relation	derived in
R0	$[x, y][r, s]$	§2
A1–A3	$AxA^{-1} = r, AyA^{-1} = s, ArA^{-1} = x$	§6.4
B1–B2	$BxB^{-1} = y^{-1}, BrB^{-1} = r$	§6.4
R3	$B(s^{-1}r^{-1}yx)B^{-1} = r^{-1}s^{-1}x$	§8
C1	$AsA^{-1} = N^{\varepsilon_3}y$	§6.3
C2	$ByB^{-1} = M^{\varepsilon_4}(yx)$	§6.1
C3	$BsB^{-1} = (r^{-1}M^{\varepsilon_5}r)s$	§6.2
F1	$M(Ax)^{\varepsilon_A} = 1$	§7.3
F2	$N((r^{-1}M^{-\varepsilon_5}r)B)^{\varepsilon_B} = 1$	§7.4

For the neighboring formal exponent pairs, the filling words acquire the fiber factors $((rx)^{-1})^n$ and $(sr^{-1}s^{-1})^m$, respectively; those systems are diagnostics, not additional manifold identifications. Known-answer probes for the fixed system give $H_1 = \mathbb{Z}^2$ after dropping F1–F2, $H_1 = \mathbb{Z}$ after dropping either one, and $H_1 = 0$ with both fillings, as [M] Lemma 4.2 requires.

The proof-level script `fixed_v_certify.g` varies independently the five signs, the left/right placement of all three transport corrections, both conjugating arcs for the Bs correction, and both arc/sign/placement choices in the T_β push-off correction. Coset enumeration

terminates with $|G| = 1$ in all $2^5 \cdot 2^3 \cdot 2 \cdot 2^3 = 4,096$ cases. In particular the geometrically derived member is trivial, and the epimorphism of Section 8 gives $\pi_1(V) = 1$. The script separately repeats all 4,096 cases after replacing the y_1 -side word Ax by the adjacent y_2 -side word Ar^{-1} ; those groups are also trivial. The first run is the one used for the specified piece.

Separately, Knuth–Bendix computations decide the nine formal exponent systems with R_3 (288/288 trivial relation groups) and an alternative based diagram (576/576). These are useful diagnostics, but the theorem makes no simple-connectivity claim for those other parametrizations. The distributed logs record reproducible software checks, not standalone formal completion traces.

10. THE SAME METHOD ON BALDRIDGE–KIRK’S T^4 : A FULLY WORKED CALIBRATION

Everything above, minus the monodromy, can be tested against a published group calculation. Take [BK] §2: $T^4 = \hat{H} \times \hat{K}$ with fiber $\hat{H}$ ($\pi_1 = \langle x, y \mid [x, y] \rangle$) and base $\hat{K}$ ($\pi_1 = \langle a, b \rangle$, based loops $\bar{a}, \bar{b}$), and the disjoint Lagrangian tori

$$T_1 = X \times A_1, \quad T_2 = Y \times A_2,$$

X, Y parallel pushoffs of x, y in the fiber, $A_1 = \{u_2 = 0.3\}$, $A_2 = \{u_2 = 0.7\}$ parallel copies of a in the base. Trivial monodromy; the membrane calculus of Sections 5–6 applies verbatim with cuts along a and b .

Crossing counts. The membranes of $\bar{a}$ -transport are clean ($\bar{a}$ is parallel to A_1, A_2 : no base crossing). For $\bar{b}$ -transport: the base track crosses A_1 once and A_2 once; the fiber conditions are $x \cap Y$, $y \cap X$ ($X \cap Y$ is one point). Result: exactly two corrected relations,

$$byb^{-1} = \mu_1^{\pm 1} y \text{ (from } T_1), \quad bxb^{-1} = \mu_2^{\pm 1} x \text{ (from } T_2),$$

each meridian based along its in-membrane whisker, the exact analogue of M, N .

The lasso datum. The fillings kill $\mu_i \cdot (\text{direction word})^{\pm 1}$ with meridian and pushoff based through the same whisker. For T_1 the direct basing is clean. For T_2 it is *not*: sliding the fiber-pushoff circle Y' down the base whisker (from $u_2 \approx 0.7$ to the basepoint row) transports the whole circle Y' , and Y' contains the point $Y \cap X$; the slide therefore crosses $T_1 = X \times A_1$ once, at $(Y \cap X) \times A_1$. Exactly the Homotopy-2 phenomenon of §7.4: testing the slide on the circle’s *basepoint* alone misses it. The repair is one lasso: the T_2 filling uses the b -conjugated meridian (equivalently, [BK]’s own $\ell_2 = bab^{-1}$ records the same wrap). The derived lasso pair is $(d_1, d_2) = (1, b)$; the $T_1 \leftrightarrow T_2$ mirror $(b, 1)$ is equivalent by an exact symmetry of the configuration.

Reference groups and outcome. [BK] Lemma 2: double ± 1 surgery gives $\pi_1 = (\mathbb{Z}^2 \rtimes_A \mathbb{Z}) \times \mathbb{Z}$ with $|\text{tr } A| \in \{1, 3\}$ by the relative signs, non-abelian in *every* convention; single surgery gives $H_3(\mathbb{Z}) \times \mathbb{Z}$. Here a computational match means equality of the abelianization and of the multisets of abelian invariants of subgroups through index 5, supplemented by explicit non-abelian finite quotients. Sweeping all 64 sign/placement conventions per lasso pair (harness `decide.t4.g`, run record ancillary):

$$\begin{aligned} (1, b) \text{ and its mirror } (b, 1) : & \quad 64/64 \text{ match the reference fingerprint,} \\ & \quad \text{with the exact tr 3/tr 1 split;} \\ \text{all seven other pairs :} & \quad 0/64 \text{ (six with } \textit{half} \text{ their failures abelian);} \\ & \quad (b^{-1}, b^{-1}) \text{ with wrong non-abelian groups throughout.} \end{aligned}$$

Single surgery has the expected $H_1 = \mathbb{Z}^3$, a quotient onto the Heisenberg group modulo 3, and the low-index fingerprint of $H_3(\mathbb{Z}) \times \mathbb{Z}$ in all eight conventions; the no-filling probe returns $H_1 = \mathbb{Z}^4$. [BK]’s own published words agree case by case with the words used

here. These are discriminating finite checks, not proofs that the compared presentations are isomorphic. Three lessons: (i) the membrane/basing/filling calculus passes a comparison against a community-vetted computation; (ii) a *single* missed whisker crossing is reliably fatal and reliably *detected*, and abelian output is its signature (this is also why any “can’t be abelian” theorem on T^4 is evidence about *wrong words*, not about methods that derive the right ones); (iii) the one class of input this calibration cannot exercise is monodromy drift (§7.3); T^4 has none, which is why that word received its own comparison: the Seifert comparison of [M] (Appendix B.2; surgery along the drift section of $\text{MT}(\varphi_0) \times S^1$). There the coherent words match the abelianization and low-index subgroup fingerprints through index 6 at three coefficients, while seven wrong-word families are separated, including the whisker-conjugate family every earlier instrument left undecided.

11. HOW TO DIFF A RELATION SHEET AGAINST OURS

Suppose an independently derived based relation sheet for $\pi_1(V)$, using the same eight generators, is at hand. To compare it with Section 9 relation by relation, quotient out the four benign discrepancies first:

- (1) *global inversions* of any generator (direction conventions; our ε -flips);
- (2) *the five swept signs* $\varepsilon_3, \varepsilon_4, \varepsilon_5, \varepsilon_A, \varepsilon_B$;
- (3) *basing conjugation*: any relator may be conjugated, and any meridian re-whiskered ($\mu \mapsto w\mu w^{-1}$), without geometric content;
- (4) *the arc identity*: $\delta = r^{-1}$ vs. $\delta' = x$ differ by a whiskered copy of c , immaterial in the group (§6.2).

A relator that fails to match modulo (1)–(4) is *the* disagreement, localized to one geometric datum (one crossing count, one lasso, one framing word), and each of ours carries the derivation above plus a verification pointer in [M]. Two structural checks then apply. An exact candidate presentation must have the known abelianization recorded in Section 9. More basically, a relation group built from generators of the manifold group and relations true there surjects onto π_1 , as in Section 8. A trivial relation group proves the manifold group trivial. A nontrivial relation group alone proves nothing about the manifold group; a contrary nontriviality claim also requires an independent proof that the proposed presentation is exact.

Verification index. Position tables and derived words: `develop_out.txt` (D3 = lassos, D4 = positions, D5 = κ_3), validations V1–V5 ([M] Appendix A.1). Fixed- V decision: `fixed_v_certify.g` and `fixed_v_certify_out.txt`. Diagnostic decisions: `placement_out.txt`, `kb_certify_out.txt`, and `kb_diag2_full_out.txt`. T^4 : `decide_t4.g`, `t4_out.txt`. All at <https://github.com/bwuebben/exotic-s2xs2>.

BERND JOHANNES WUEBBEN, NEW YORK, NY. wuebben@gmail.com