

Introduction to Machine Learning

Christos Dimitrakakis

September 11, 2024

Outline

The problems of Machine Learning (1 week)

Introduction

Estimation

Answering a scientific problem

Pandas and dataframes

Single variable models

Two variable models

Statistics, validation and model selection

Course summary

Course Contents

Reading for this week

Reading

The problems of Machine Learning (1 week)

Introduction

Estimation

Answering a scientific problem

Pandas and dataframes

Single variable models

Two variable models

Statistics, validation and model selection

Course summary

Course Contents

Reading for this week

Reading

The problems of Machine Learning (1 week)

Introduction

Estimation

Statistics, validation and model selection

Course summary

Reading for this week

Machine Learning And Data Mining

The nuts and bolts

- ▶ Models
- ▶ Algorithms
- ▶ Theory
- ▶ Practice

Workflow

- ▶ Scientific question
- ▶ Formalisation of the problem
- ▶ Data collection
- ▶ Analysis and model selection

Types of statistics / machine learning problems

- ▶ Classification
- ▶ Regression
- ▶ Density estimation
- ▶ Reinforcement learning

The nuts and bolts

- ▶ Models
- ▶ Algorithms
- ▶ Theory
- ▶ Practice

Machine learning

Data Collection

- ▶ Downloading a clean dataset from a repository
- ▶ Performing a survey
- ▶ Scraping data from the web
- ▶ Deploying sensors, performing experiments, and obtaining measurements.


Modelling (what we focus on this course)

- ▶ Simple: the bias of a coin
- ▶ Complex: a language model.
- ▶ The model depends on the data and the problem

Algorithms and Decision Making


- ▶ We want to use models to make decisions.
- ▶ Decisions are made every step of the way.
- ▶ Decisions are automated algorithmically.

The main problems in machine learning and statistics




Matter


Dark



Protein Folding




Climate Modelling




Economic Policy

Prediction




- ▶ Will it rain tomorrow?
- ▶ How much will bitcoin be worth next year?
- ▶ When is the next solar eclipse?

Inference



- ▶ Does my poker opponent have two aces?
- ▶ What is the law of gravitation?

Decision Making



./fig/artemis.gif

- ▶ What data should I collect?
- ▶ Which model should I use?
- ▶ Should I fold, call, or raise in my poker game?
- ▶ How can I get a spaceship to the moon and back?

The need to learn from data

Problem definition

- ▶ What problem do we need to solve?
- ▶ How can we formalise it?
- ▶ What properties of the problem can we learn from data?

Data collection

- ▶ **Why** do we need data?
- ▶ **What** data do we need?
- ▶ How **much** data do we want?
- ▶ **How** will we collect the data?

Modelling and decision making

- ▶ How will we **compute** something useful?
- ▶ How can we use the model to make **decisions**?

Course Material

Moodle

- ▶ Assignments and project
- ▶ Additional reading material
- ▶ Asking questions

Course Github <https://github.com/olethrosdc/machine-learning-neuch/tree/main/BSc>

- ▶ .org files for notes, PDF for slides
- ▶ source code for examples

Course literature

- ▶ An Introduction to Statistical Learning with Python https://hastie.su.domains/ISLP/ISLP_website.pdf.download.html

book chapters will be mentioned in the course

Assignment, teaching and questions

Assignments and project

- ▶ Individual Weekly assignments in the first half
- ▶ Group project in the second half
- ▶ Project presentation
- ▶ No exam.

Other questions

- ▶ Use Moodle for technical/administrative questions: That way everybody gets the same information.
- ▶ Use email for personal problems or extra help, if the moodle is not enough.
- ▶ Complicated questions can be answered at the next lecture

Office hours

- ▶ Fridays 13:00-14:00: book with an email to avoid clashes.
- ▶ Email me for an appointment outside those hours.

The problems of Machine Learning (1 week)

Introduction

Estimation

Answering a scientific problem

Pandas and dataframes

Single variable models

Two variable models

Statistics, validation and model selection

Course summary

Course Contents

Reading for this week

Reading

Problem definition

- ▶ Example: Health, weight and height

Example (Health questions regarding height and weight)

- ▶ What is a normal height and weight?
- ▶ How are they related to health?
- ▶ What variables affect height and weight?

Data collection

Think about which variables we need to collect to answer our research question.

Necessary variables

The variables we need to know about

- ▶ Weight
- ▶ Height
- ▶ Dependent: (health/vote/opinion/salary)

Auxiliary variables

Measurable factors related to the variables of interest

Possible confounders

Hidden factors that might affect variables

Class data and variables

- ▶ The class enters their data into the excel file.



- ▶ Pay attention to the variables we wish to measure

Privacy

- ▶ Is the use of a pseudonym sufficient to hide your identity?

Types of learning problems

Unsupervised learning (unconditional estimation)

- ▶ Predict the **gender** of an unknown individual.
- ▶ Predict the **height**.
- ▶ Predict the **height and weight**?

Supervised learning problems (conditional estimation)

- ▶ Classification: Can we predict gender from height/weight?
- ▶ Regression: Can we predict weight from height and gender?
- ▶ In both cases we predict **output** variables from **input** variables

Variables

- ▶ **Input** variables: aka features, predictors, independent variables
- ▶ **Output** variables: aka response, dependent variables, labels, or targets.
- ▶ The input/output dichotomy only exists in **some prediction problems**.

Variables

The class data looks like this

| First Name | Gender | Height | Weight | Age | Nationality | Smoking |
|------------|--------|--------|--------|-----|-------------|---------|
| Lee | M | 170 | 80 | 20 | Chinese | 10 |
| Fatemeh | F | 150 | 65 | 25 | Turkey | 0 |
| Ali | Male | 174 | 82 | 19 | Turkish | 0 |
| Joan | N | 5'11 | 180 | 21 | Brtish | 4 |

- ▶ \mathbf{X} : Everybody's data
- ▶ \mathbf{x}_t : The t -th person's data
- ▶ $x_{t,k}$: The k -th feature of the t -th person.
- ▶ \mathbf{x}_k : Everybody's k -th feature

We will not use special symbols to distinguish random variables from their realisations

Raw versus neat data

- ▶ Neat data: $\mathbf{x}_t \in \mathbb{R}^n$
- ▶ Raw data: text, graphs, missing values, etc

Python pandas for data wrangling

Reading class data

```
import pandas as pd
X = pd.read_excel("data/class.xlsx")
X["First_Name"]
```

- ▶ Array columns correspond to features
- ▶ Columns can be accessed through namesx

Summarising class data

```
X.hist()
import matplotlib.pyplot as plt
plt.show()
```

Pandas and DataFrames

- ▶ Data in pandas is stored in a **DataFrame**
- ▶ DataFrame is **not the same** as a numpy array.

Core libraries

```
import pandas as pd
import numpy as np
```

Series: A sequence of values

```
# From numpy array:
s = pd.Series(np.random.randn(3),
index=["a", "b", "c"])
# From dict:
d = {"a": 1, "b": 0, "c": 2}
s = pd.Series(d)
# accessing elements
s.iloc[2] #element 2
s.iloc[1:2] #elements 1,2
s.array # gets the array object
s.to_numpy() # gets the underlying numpy array
```

DataFrames

Constructing from a numpy array

```
data = np.random.uniform(size = [3,2])  
df = pd.DataFrame(data, index=["John", "Ali", "Sumi"],  
                  columns=["X1", "X2"])
```

Constructing from a dictionary

```
d = { "one": pd.Series([1, 2], index=["a", "b"]),  
      "two": pd.Series([1, 2, 3], index=["a", "b", "c"])  
df = pd.DataFrame(d)
```

Access

```
X["First_Name"] # get a column  
X.loc[2] # get a row  
X.at[2, "First_Name"] # row 2, column 'first name'  
X.loc[2].at["First_Name"] # row 2, element 'first name'  
X.iat[2,0] # row 2, column 0
```

Modelling single variables




Figure: $x \in \mathbb{N}$




Figure: $x \in \mathbb{R}$

Means using python

Example (Calculating the mean of our class data)

```
X.mean() # gives the mean of all the variables through pandas  
X["Height"].mean()  
np.mean(X["Weight"])
```

- ▶ The mean here is **fixed** because we calculate it on the same data.
- ▶ If we were to **collect new data** then the answer would be different.

Example (Calculating the mean of a random variable)

```
import numpy as np  
X = np.random.gamma(170, 1, size=20)  
X.mean()  
np.mean(X)
```

- ▶ The mean is **random**, so we get a different answer everytime.

One variable: expectations and distributions

Definition (The expected value)

Assume $x : \Omega \rightarrow \mathbb{R}$, and $\omega_t \sim P$

- ▶ $x_1, \dots, x_t, \dots, x_T$: random i.i.d. variables with $x_t = x(\omega_t)$
- ▶ Ω : random outcome space
- ▶ P : distribution of outcomes $\omega \in \Omega$
- ▶ $\mathbb{E}_P[x]$: expectation of x under P

$$\mathbb{E}_P[x_t] = \sum_{\omega \in \Omega} x_t(\omega) P(\omega)$$

One variable: expectations and distributions

Definition (The expected value)

Assume $x : \Omega \rightarrow \mathbb{R}$, and $\omega_t \sim P$

- ▶ $x_1, \dots, x_t, \dots, x_T$: random i.i.d. variables with $x_t = x(\omega_t)$
- ▶ Ω : random outcome space
- ▶ P : distribution of outcomes $\omega \in \Omega$
- ▶ $\mathbb{E}_P[x]$: expectation of x under P

$$\mathbb{E}_P[x_t] = \sum_{\omega \in \Omega} x_t(\omega) P(\omega)$$

Definition (The sample mean)

The sample mean of x_1, \dots, x_T is

$$\frac{1}{T} \sum_{t=1}^T x_t$$

Under P , the sample mean is $O(1/\sqrt{T})$ -close to the expected value $\mathbb{E}_P[x_t]$.

Reminder: expectations of random variables

A gambling game

What are the expected winnings if you play this game?

- ▶ [a] With probability 1%, you win 100 CHF
- ▶ [b] With probability 40%, you win 20 CHF.
- ▶ [c] Otherwise, you win nothing

Solution

Reminder: expectations of random variables

A gambling game

What are the expected winnings if you play this game?

- ▶ [a] With probability 1%, you win 100 CHF
- ▶ [b] With probability 40%, you win 20 CHF.
- ▶ [c] Otherwise, you win nothing

Solution

- ▶ Let x be the amount won, then $x(a) = 100, x(b) = 20, x(c) = 0$.

Reminder: expectations of random variables

A gambling game

What are the expected winnings if you play this game?

- ▶ [a] With probability 1%, you win 100 CHF
- ▶ [b] With probability 40%, you win 20 CHF.
- ▶ [c] Otherwise, you win nothing

Solution

- ▶ Let x be the amount won, then $x(a) = 100, x(b) = 20, x(c) = 0$.
- ▶ We need to calculate

$$\mathbb{E}_P(x) = \sum_{\omega \in \{a,b,c\}} x(\omega)P(\omega) = x(a)P(a) + x(b)P(b) + x(c)P(c)$$

Reminder: expectations of random variables

A gambling game

What are the expected winnings if you play this game?

- ▶ [a] With probability 1%, you win 100 CHF
- ▶ [b] With probability 40%, you win 20 CHF.
- ▶ [c] Otherwise, you win nothing

Solution

- ▶ Let x be the amount won, then $x(a) = 100, x(b) = 20, x(c) = 0$.
- ▶ We need to calculate

$$\mathbb{E}_P(x) = \sum_{\omega \in \{a,b,c\}} x(\omega)P(\omega) = x(a)P(a) + x(b)P(b) + x(c)P(c)$$

- ▶ $P(c) = 59\%$, as $P(\Omega) = 1$. Substituting,

Reminder: expectations of random variables

A gambling game

What are the expected winnings if you play this game?

- ▶ [a] With probability 1%, you win 100 CHF
- ▶ [b] With probability 40%, you win 20 CHF.
- ▶ [c] Otherwise, you win nothing

Solution

- ▶ Let x be the amount won, then $x(a) = 100, x(b) = 20, x(c) = 0$.
- ▶ We need to calculate

$$\mathbb{E}_P(x) = \sum_{\omega \in \{a,b,c\}} x(\omega)P(\omega) = x(a)P(a) + x(b)P(b) + x(c)P(c)$$

- ▶ $P(c) = 59\%$, as $P(\Omega) = 1$. Substituting,
$$\mathbb{E}_P(x) = 1 + 8 + 0 = 9.$$

Models

Models as summaries

- ▶ They summarise what we can see in the data
- ▶ The ultimate model of the data **is** the data

Models as predictors

- ▶ They make predictions about things **beyond** the data
- ▶ This requires some assumptions about the **data-generating process**.

Example models

- ▶ A numerical mean
- ▶ A linear classifier
- ▶ A linear regressor
- ▶ A deep neural network
- ▶ A Gaussian process
- ▶ A large language model

The Bernoulli distribution

Definition (Bernoulli distribution)

We say that $x \in \{0, 1\}$ has Bernoulli distribution with parameter θ and write

$$x \sim \text{Bernoulli}(\theta),$$

when

$$\mathbb{P}(x) = \begin{cases} \theta & x = 1 \\ 1 - \theta & x = 0. \end{cases}$$

Example (Applications of the Bernoulli distribution)

- ▶ A biased coin flip.
- ▶ Classification errors.

Exercise: The expected value

If x is Bernoulli with parameter θ , then what is the expected value of

- ▶ The variable $f(x) = x$?
- ▶ The variable $g(x) = x^2$?

Two-variable models




Figure: $x \in \mathbb{R}^2$




Figure: $x \in \mathbb{R} \rightarrow y \in \mathbb{N}$




Figure: $x \in \mathbb{N} \rightarrow y \in \mathbb{R}$




Figure: $x \in \mathbb{R} \rightarrow y \in \mathbb{R}$

Predicting y from x , discrete case.

Consider two variables, x, y . We can either care about

- ▶ $\mathbb{E}[y|x]$ the expectation of y for all x .
- ▶ $\mathbb{P}[y|x]$ the distribution of y for all x .

Probability table for $P(x, y)$

| $P(x, y)$ | $y = 0$ | $y = 1$ |
|-----------|---------|---------|
| $x = 0$ | 54% | 6% |
| $x = 1$ | 16% | 24% |

- ▶ How can we graph this?
- ▶ What is $P(x)$?

Conditional probability table for $P(y|x)$

| $P(y x)$ | $y = 0$ | $y = 1$ |
|------------|---------|---------|
| $x = 0$ | 90% | 10% |
| $x = 1$ | 40% | 60% |

- ▶ What is $\mathbb{E}[y | x]$?

Distributions of two variables

In this setting, both x and y have a Bernoulli distribution. If we use a model whereby x is sampled first, and then y , then we can define two Bernoulli distributions. The first, for x is unconditional, while the second, for y , depends on the value of x :

$$\begin{aligned}x &\sim \text{Bernoulli}(\theta) \\ y \mid x &\sim \text{Bernoulli}(\phi_x).\end{aligned}$$

In our example, $\phi_0 = 0.1$ and $\phi_1 = 0.6$.

Homework

Probability table for $P(x, y)$

| $P(x, y)$ | $y = -1$ | $y = 0$ | $y = 1$ |
|-----------|----------|---------|---------|
| $x = 0$ | 10% | 20% | 10% |
| $x = 1$ | 30% | 20% | 10% |

Given the above table, calculate

- ▶ $P(x)$
- ▶ The conditional probability table for $P(y|x)$.
- ▶ $\mathbb{E}[y|x]$ for all values of x .

Two variables: conditional expectation

The height of different genders

The conditional expected height

$$\mathbb{E}[h \mid g = 1] = \sum_{\omega \in \Omega} h(\omega) P[\omega \mid g(\omega) = 1]$$

The empirical conditional expectation

$$\mathbb{E}[h \mid g = 1] \approx \frac{\sum_{t: g(\omega_t)=1} h(\omega_t)}{|\{t : g(\omega_t) = 1\}|}$$

Python implementation

Two variables: conditional expectation

The height of different genders

The conditional expected height

$$\mathbb{E}[h \mid g = 1] = \sum_{\omega \in \Omega} h(\omega) P[\omega \mid g(\omega) = 1]$$

The empirical conditional expectation

$$\mathbb{E}[h \mid g = 1] \approx \frac{\sum_{t: g(\omega_t)=1} h(\omega_t)}{|\{t : g(\omega_t) = 1\}|}$$

Python implementation

```
h[g==1] / sum(g==1)
## alternative
import numpy as np
np.mean(h[g==1])
```


The problems of Machine Learning (1 week)

Introduction

Estimation

Answering a scientific problem

Pandas and dataframes

Single variable models

Two variable models

Statistics, validation and model selection

Course summary

Course Contents

Reading for this week

Reading

Populations, samples, and distributions




Figure: The world population



Figure: A sample

Statistical assumptions

Independent, Identically Distributed data

- ▶ $\omega_t \sim P$: individuals $\omega_t \in \Omega$ are drawn from some **distribution** P
- ▶ $\mathbf{x}_t \triangleq \mathbf{x}(\omega_t)$ are some **features** of the t -th individual
- ▶ Here we are interested in properties of the **unknown** distribution P .

Representative sample from a fixed population

- ▶ Finite population $\Omega = \{\omega_1, \omega_2, \dots, \omega_N\}$
- ▶ A subset $S \subset \Omega$ of size $T < N$ is selected with a **uniform distribution**, i.e. so that

$$P(S) = T/N, \quad \forall S \subset \Omega.$$

- ▶ Here we are interested in statistics of the **unknown** population Ω .
- ▶ We assume an underlying distribution P for convenience.
- ▶ We can try both cases essentially the same.

Learning from data

Unsupervised learning

- ▶ Given data x_1, \dots, x_T .
- ▶ Learn about the data-generating process.
- ▶ Example: Estimation, compression, text/image generation

Supervised learning

- ▶ Given data $(x_1, y_1), \dots, (x_T, y_T)$
- ▶ Learn about the relationship between x_t and y_t .
- ▶ Example: Classification, Regression

Online learning

- ▶ Sequence prediction: At each step t , predict x_{t+1} from x_1, \dots, x_t .
- ▶ Conditional prediction: At each step t , predict y_{t+1} from $x_1, y_1, \dots, x_t, y_t, x_{t+1}$

Reinforcement learning

Learn to act in an **unknown** world through interaction and rewards

Validating models

Training data

- ▶ Calculations, optimisation
- ▶ Data exploration

Validation data

- ▶ Fine-tuning
- ▶ Model selection

Test data

- ▶ Performance comparison

Simulation

- ▶ Interactive performance comparison
- ▶ White box testing

Real-world testing

- ▶ Actual performance measurement

Model selection

- ▶ Train/Test/Validate
- ▶ Cross-validation
- ▶ Simulation

The problems of Machine Learning (1 week)

Introduction

Estimation

Answering a scientific problem

Pandas and dataframes

Single variable models

Two variable models

Statistics, validation and model selection

Course summary

Course Contents

Reading for this week

Reading

Course Contents

Models

- ▶ k-Nearest Neighbours.
- ▶ Linear models and perceptrons.
- ▶ Multi-layer perceptrons (aka deep neural networks).
- ▶ Bayesian Networks

Algorithms

- ▶ (Stochastic) Gradient Descent.
- ▶ Bayesian inference.

Reproducibility

- ▶ Modelling assumptions
- ▶ Interactions and feedback

Fairness

- ▶ Implicit biases in training data
- ▶ Fair decision rules and meritocracy

The problems of Machine Learning (1 week)

Introduction

Estimation

Answering a scientific problem

Pandas and dataframes

Single variable models

Two variable models

Statistics, validation and model selection

Course summary

Course Contents

Reading for this week

Reading

Reading for this week

ISLP Chapter 1