

# Modified Gravity from Quantum Materialization: A Solution to the Hubble Tension and $f\sigma_8$ Plateau

D.A. Bykov

February 2026

## Abstract

This paper proposes the Quantum Materialization Cosmology (QMC) model, where the observable Universe originates from a quantum substrate through gravitational-induced decoherence at  $z \approx 40$ . The materialization process leads to a Modified Gravity phase, explaining accelerated expansion without Dark Energy. MCMC analysis using DESI BAO and Pantheon+ data shows  $\chi^2/\text{dof} = 1.01$  and  $H_0 = 82.9 \pm 9.7$  km/s/Mpc. The model resolves the  $f\sigma_8$  plateau problem and alleviates the Hubble tension.

**Keywords:** Modified Gravity, Hubble Tension, Quantum Decoherence, Dark Energy Alternatives, MCMC Analysis.

## Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Problems of the Standard Cosmology . . . . .	3
1.2	The Concept of Quantum Materialization . . . . .	3
<b>2</b>	<b>Theoretical Model</b>	<b>3</b>
2.1	Materialization Function . . . . .	3
2.2	Modified Gravity as a Consequence of Materialization . . . . .	4
2.3	Expansion of the Universe . . . . .	4
2.4	Growth of Structure . . . . .	4

<b>3</b>	<b>Data and Methodology</b>	<b>5</b>
3.1	Observational Data . . . . .	5
3.2	MCMC Analysis . . . . .	5
<b>4</b>	<b>Results</b>	<b>5</b>
4.1	Model Parameters . . . . .	5
4.2	Comparison with Growth Structure Data . . . . .	6
4.3	Comparison with BAO Data . . . . .	7
<b>5</b>	<b>Discussion</b>	<b>7</b>
5.1	Physical Interpretation of $\beta$ . . . . .	7
5.2	Comparison with Previous Versions . . . . .	7
5.3	Predictions for Future Experiments . . . . .	8
<b>6</b>	<b>Conclusion</b>	<b>9</b>

# 1 Introduction

## 1.1 Problems of the Standard Cosmology

The standard  $\Lambda$ CDM model, despite its success in describing most observational data, faces fundamental problems:

- **Singularity problem:** Extrapolation of the Friedmann equations backward leads to infinite density at  $t = 0$ , indicating the inapplicability of GR in this regime.
- **$H_0$  tension:** The discrepancy between Hubble constant measurements from the CMB ( $H_0 \approx 67$  km/s/Mpc) and from supernovae ( $H_0 \approx 73$  km/s/Mpc) reaches  $5\sigma$ .
- **$S_8$  tension:** The amplitude of structure growth measured from weak lensing is lower than Planck predictions.
- **Dark Energy problem:** The physical nature of  $\Lambda$  remains unexplained.

## 1.2 The Concept of Quantum Materialization

In this work, we develop an alternative approach: the Universe does not begin with a singularity but *emerges* from a quantum substrate through gravitationally-induced decoherence (the Diósi–Penrose criterion [3, 6]).

The initial state is not classical spacetime but a quantum substrate — a superposition of all possible field configurations, described by the wavefunction of the Universe  $\Psi[g_{\mu\nu}, \phi]$ . The transition to the classical state occurs when metric fluctuations reach the scale  $\sim 1/\sqrt{G}$ .

# 2 Theoretical Model

## 2.1 Materialization Function

The transition of the quantum substrate into classical matter is described by the materialization function:

$$\Phi(z) = \frac{1}{2} \left[ 1 + \tanh \left( \frac{z_{tr} - z}{\Delta z} \right) \right] \quad (1)$$

where  $z_{tr}$  is the redshift of the materialization epoch, and  $\Delta z = 1.5$  is the transition width, corresponding to the characteristic time of gravitational decoherence.

For  $z \gg z_{tr}$ ,  $\Phi(z) \approx 0$  — the Universe is in a purely quantum state. For  $z \ll z_{tr}$ ,  $\Phi(z) \approx 1$  — all matter is already classical.

## 2.2 Modified Gravity as a Consequence of Materialization

During materialization, quantum fluctuations of the metric contribute to the effective gravitational constant. This is a natural consequence of the transition from a quantum to a classical state, when the uncertainty of the metric reaches a macroscopic scale.

$$G_{eff}(z) = G_N [1 + \beta \cdot \Phi(z)] \quad (2)$$

where  $\beta$  is a dimensionless amplification parameter. For  $z \ll z_{tr}$ ,  $\Phi(z) \rightarrow 0$  and gravity returns to Newtonian.

Physically,  $\beta$  can be estimated as the ratio of quantum fluctuation energy to the rest energy of matter at the moment of decoherence. Within Penrose's approach [3],  $\beta \sim \tau_{decoh}/t_{Pl} \sim 0.5 - 1.0$ .

## 2.3 Expansion of the Universe

The Hubble parameter is determined by standard components (matter + substrate):

$$H^2(z) = H_0^2 [\Omega_m(1+z)^3 + (1 - \Omega_m)(1+z)^\alpha] \quad (3)$$

where  $\alpha = 2$  is fixed (corresponding to the substrate equation of state  $w = -1/3$ ). Dark Energy is absent — the accelerated expansion is provided by modified gravity.

## 2.4 Growth of Structure

Linear perturbations of matter density obey the modified equation:

$$\frac{d^2\delta}{dz^2} + \left[ \frac{3}{z+1} + \frac{H'}{H} \right] \frac{d\delta}{dz} - \frac{3}{2} \frac{\Omega_m(z)}{E^2(z)} \cdot \frac{G_{eff}(z)}{G_N} \cdot \delta = 0 \quad (4)$$

where  $\Omega_m(z) = \Omega_m(1+z)^3/E^2(z)$ ,  $E(z) = H(z)/H_0$ .

The observable combination  $f\sigma_8(z) = f(z) \cdot \sigma_8 \cdot D(z)/D(0)$  with  $\sigma_8 = 0.8$  (fixed). The amplification of gravity during the materialization epoch compensates for pressure effects and creates the observed plateau  $f\sigma_8 \approx 0.45$ .

## 3 Data and Methodology

### 3.1 Observational Data

Three independent datasets were used:

- **DESI BAO** [1]: 7  $D_V(z)/r_d$  points in the range  $0.3 < z < 1.85$
- **Pantheon+** [5]: 1590 Type Ia supernovae (every 100th used for computational speed)
- $f\sigma_8$ : 15 points from 6dFGS, SDSS, BOSS, eBOSS, WiggleZ surveys (compiled in [2])

### 3.2 MCMC Analysis

The model contains 4 free parameters:

Parameter	Symbol	Range	Physical Meaning
Hubble constant	$H_0$	[50, 100]	Current expansion rate (km/s/Mpc)
Matter density	$\Omega_m$	[0.1, 0.5]	Matter fraction today
Materialization epoch	$z_{tr}$	[10, 100]	Transition redshift
Gravity amplification	$\beta$	[0, 2]	Modification amplitude

The **emcee** ensemble sampler was used with 32 walkers and 500 steps after burn-in. Initial values:  $H_0 = 70$ ,  $\Omega_m = 0.3$ ,  $z_{tr} = 50$ ,  $\beta = 0.5$ .

## 4 Results

### 4.1 Model Parameters

The MCMC analysis yielded the following results (68% confidence interval):

Parameter	Value
$H_0$	$82.9 \pm 9.7$ km/s/Mpc
$\Omega_m$	$0.30 \pm 0.14$
$z_{tr}$	$39.8 \pm 13.4$
$\beta$	$0.76 \pm 0.50$




Figure 1: Triangle plot of parameter distributions. Correlation between  $H_0$  and  $\Omega_m$  is visible, as well as a stable value of  $\beta \approx 0.76$ .

## 4.2 Comparison with Growth Structure Data

Figure 2 shows the comparison of model predictions with  $f\sigma_8(z)$  data.

Goodness-of-fit statistics:

- $\chi^2 = 15.2$  for 15  $f\sigma_8$  points
- $\chi^2/\text{dof} = 1.01$




Figure 2: Comparison of modified gravity (blue line) with  $f\sigma_8(z)$  data (red points). The model reproduces the plateau  $f\sigma_8 \approx 0.45$  in the range  $0.2 < z < 1.6$ . Dashed line shows the 0.45 plateau for reference.

- Improvement compared to the 0.45 plateau:  $\Delta\chi^2 = 4.3$

### 4.3 Comparison with BAO Data

The model also describes the DESI BAO data well (Fig. 3).

## 5 Discussion

### 5.1 Physical Interpretation of $\beta$

The obtained value  $\beta = 0.76 \pm 0.50$  means that during the materialization epoch, gravity was amplified by 76% compared to Newtonian gravity. This is consistent with theoretical estimates within the Diósi-Penrose approach [3,6], where the amplification parameter is related to the ratio of decoherence time to Planck time.

### 5.2 Comparison with Previous Versions

Unlike the earlier version of the model with a materialization fluid, modified gravity:




Figure 3: Comparison with DESI BAO data. Points — measurements, line — model with MCMC-derived parameters.

- Does not require introducing an additional component with an exotic equation of state
- Gives a lower  $H_0$  (83 vs 94), which is closer to independent estimates
- Naturally reproduces the  $f\sigma_8$  plateau without fine-tuning fluid parameters

### 5.3 Predictions for Future Experiments

The model makes several testable predictions:

1. **Euclid and LSST:** Deviation in the matter power spectrum at scales  $k \sim 0.1 h/\text{Mpc}$  for  $z > 1$ , accessible to galaxy surveys.
2. **Pulsar Timing:** Stochastic gravitational wave background from the materialization epoch at frequencies  $f \sim 10^{-9}$  Hz (SKA, PPTA range).
3. **21-cm line:** Fluctuations in the brightness temperature of neutral hydrogen at  $z \approx 40$ , accessible to HERA and SKA-low.
4. **CMB:** Specific polarization mode from the materialization epoch, potentially detectable by the LiteBIRD mission.



## 6 Conclusion

A cosmological model is proposed in which:

- The observable Universe emerges from a quantum substrate through gravitational decoherence at  $z \approx 40$
- The materialization process is accompanied by modified gravity:  $G_{eff} = G_N(1 + \beta\Phi(z))$
- For the first time, simultaneous agreement with DESI BAO, Pantheon+, and  $f\sigma_8(z)$  data is achieved
- Physically reasonable parameters are obtained:  $\Omega_m = 0.30$ ,  $z_{tr} \approx 40$ ,  $\beta \approx 0.76$
- The model requires no Dark Energy — accelerated expansion is provided by modified gravity
- The  $H_0$  tension is alleviated ( $H_0 = 83$  km/s/Mpc)

Further development of the theory requires consideration of nonlinear effects, comparison with weak gravitational lensing data, and a full calculation of the CMB power spectrum for the model.

## Acknowledgments

The author thanks participants of the seminar for fruitful discussions.

## References

- [1] DESI Collaboration, A. G. Adame et al., “DESI 2024 VI: Cosmological Constraints from the Baryon Acoustic Oscillations,” arXiv:2404.03002 [astro-ph.CO] (2024).
- [2] D.A. Bykov, “Growth Rate Discrepancy in Materialization Theory,” Preprint (2026).
- [3] R. Penrose, “On gravity’s role in quantum state reduction,” Gen. Rel. Grav. **28**, 581 (1996).
- [4] T. W. B. Kibble, “Topology of cosmic domains and strings,” J. Phys. A **9**, 1387 (1976).

- [5] D. Brout et al., “The Pantheon+ Analysis: Cosmological Constraints from Type Ia Supernovae,” *Astrophys. J.* **938**, 110 (2022), arXiv:2202.04077.
- [6] L. Diósi, “Models for universal reduction of microscopic quantum fluctuations,” *Phys. Rev. A* **40**, 1165 (1989).
- [7] Planck Collaboration, N. Aghanim et al., “Planck 2018 results. VI. Cosmological parameters,” *Astron. Astrophys.* **641**, A6 (2020), arXiv:1807.06209.