Q2. Find approximate cashflow for Hoorlet on one month GBOR, using VasicEx Model

We start, as instructed, with the yield are power socies expansion.

D We know that zero-corpor bonds satisfy:

(1)
$$\frac{J^2}{Jt} + \frac{1}{2}w^2J^2_{zz} + (u-\lambda w)\frac{J^2}{J^2} - F^2_{zz} = 0$$
 where w, μ, λ are functions of (t, ν)

We look to a sollion of equation above of the form (Taylor seves about t=T)

Finding decirations:
$$\frac{dz}{dt} = -a - 2b(T-t)$$
; $\frac{dz}{dt} = a'(T-t) + b'(T-t)^2$

$$\frac{d^2z}{dt^2} = a''(T-t) + b''(T-t)^2$$

Now, replacing in (+)

$$-a-2b(T-t)+1u^{2}(a''(T-t)+b''(T-t)^{2})+(4*\lambda w)(a'(T-t)+b'(T-t)^{2})-r(1+a(T-t)+b(T-t))=0$$

ve-avanging (grouping on (T-6) and (T-6)2)

$$-a - (T-t) \left[2b - \frac{1}{2} w^{2} a'' - a' (u - \lambda w) + ra \right] + (T-t)^{2} \left[\frac{1}{2} w^{2} b'' + (u - \lambda w) b' - rb \right] = V$$

Now equating coefficients:

(9)
$$-q=V \Rightarrow |q=-V| \Rightarrow |q=-V| \Rightarrow |q'=0|$$

(b)
$$2b - \frac{1}{2}w^{2}a' - a'(u - \lambda w) + Va = 0$$

replacing
$$\Rightarrow 2b - 0 - (-1)(u - \lambda w) + t - (-t) = 0$$

$$b = \frac{V^2 + \lambda w}{2}$$

bow replacing in agration (2):

So ZN |- V(T-t) + 1 (T-t2) (12-u+xw)+.... (3) this is a Taylor saies approximation for a zero-corpon bond solution Now, as we are sughating I wonth LiBOR, we want to find the short term 1/2. So, the relation between Zeno-Corpon and rate: $2(r,t;T) = e^{-r(T-t)}$ finding r: ln(Z) = -r(T-t) $Y = -\ln(z) \tag{4}$ So, we need to find (4) in eq. (3): Applying In 11 in (3): ln (2) ~ ln(1-r(T-t)+1(T-t)(r-a+hw) +.... near the short and, is small, so: In (1+x) & X near X = 0 lm(2) ~- +(T-t) + 1 (T-t)2(12u+hw) +... ananging + ln(2) ~- r(T-E) + 1(T-E) 2 - 1(T-E)2 (u- hu) +... lan(2) ~- Y(T-6) - 1 (T-6)2(u-)w) +... $\frac{-2n(2)}{(T-t)} \sim V + \frac{1}{2} (T-t) (u-\lambda_w) +$ (all as toT) or (T-6)-00 by (4), this is an approximation for the short vale (Imenth 4BOR) Then, the Hoalet cashflow: MAX [K-L, O] = MAX[Hoalet - KiBOR, O] In this case: Kyloovlot = Ky HAX[Y,-(Y-1-1/12-(η-μν),0]

MAX[Y,-(Y-1/2-1/12-(η-μν),0] YUBOR = Y-1 (T-t) (U-) W but: T-t= 1 (Amonth) by Vasicer: dr= (n-yr)de+ cdw + la-xw=n-yr


- 3 We with to obtain a model of the form: dr = ululd+ Yr dw FROM DATA
 - @ First, let's examine the volatility structure: Ur dw

Forso:
$$(dr)^2 = (u(r)dt + y)r dw)^2 = u(u)(dt)^2 + y)^2 \frac{2B}{dt} (dw)^2 + 2u(u)yr \frac{B}{dt} dw$$

$$(dv)^2 = y^2 \frac{2B}{dt} (1)$$

- Now, let's divide our data im Jr.
- Then, we use bucketing technique. We build tuckets of Jr covering a range in rvalues.
- Then we calculate (Jr)2 toveach bucket.
- We then averge the values of (Si)2 for each broket.
- -then, thom (1) we have: E[(Jr)2]=y220dt at thom data time step Je is one day -Applying log: $log(E[(3r)^2]) = 2log V + 2\beta log V$

 - So whatity inverses as spot vale inverses



(b) let's examine the drift structure. Is difficult to estimate drift from data, the drift term is smaller than the volatility term, and this subject to larger relative errors.

Better approach is to find the drift truction via empirical and analytical determination of the steady-state probability trunction for r

it v statisties: dv = uculd+ yr dw, then the puobability donsity function part to Y satisfies the Former Plance equation:

$$\frac{JP}{Jt} = \frac{1}{2} v^2 J^2 (v^2 P) - \frac{1}{2} (uu) P$$
 (2)

the steady state Poo(r) satisfies the time-independent vasion of (2): 1 y2/2 (r28) - d (ulu) po) = 0 Now, integraling once (3) $\Rightarrow \frac{1}{2}v^2d^2(rp_{\infty}) = \frac{1}{2}(v(l)p_{\infty})$ 1 V2 (1 Poo) = u(r) Poo P. 2 2 2 (r Poo) = u(r) Now, Rodrotrule => 1 1 y2 [2Br. Poo + rPoo] = u(r) $V_{B}^{2} + \frac{P_{0}^{1}}{P_{0}} \cdot \frac{1}{2} V_{F}^{2} = u(r)$ but d (log Po) = 1. Poo (Chain rule) So Linally: $u(r) = y^{2}_{13}r^{2\beta-1} + \frac{1}{2}v^{2}r^{2\beta} + \frac{1}{2}(\log P_{\infty})$ (4) Now we explare the empirical data, build an histogram, and fit a density Lunction Po : We tind a and F titting on data. Now, replacing in (4) $\Rightarrow \alpha(r) = \nu^2 \beta r^{2\beta-1} + \frac{1}{2} \nu^2 \frac{zB}{2r} \left[\log \left(\frac{1}{qr\sqrt{z}r} \cdot \exp \left(\frac{1}{2a^2} \left(\log (r/r) \right)^2 \right) \right]$ $= \frac{1}{2r} \left[\log \left(\frac{1}{arR} \right) + \log \left(\frac{1}{2ar} \left(\log (V_L) \right)^2 \right) \right] = \frac{1}{ar} \left[\log \left(\frac{1}{arR} \right) - \frac{1}{2a^2} \left(\log \frac{V}{r} \right)^2 \right] = \frac{1}{ar} \left[\log \left(\frac{1}{r} \right) + \log \left(\frac{1}{arR} \right) - \frac{1}{2a^2} \left(\log \frac{V}{r} \right)^2 \right]$ Evaluating $\frac{1}{\sqrt{dr}} = \left[V - \frac{1}{\sqrt{2}} + 0 - \frac{1}{\sqrt{2}} \left[2 \log \left(\frac{r}{F} \right) - \left[\frac{1}{r} + 0 \right] - 1 \right] = \left[-\frac{1}{r} - \frac{1}{\sqrt{2}} \log \left(\frac{r}{F} \right) \right] = + \frac{1}{r} \left[-1 - \frac{1}{\sqrt{2}} \log \left(\frac{r}{F} \right) \right]$ NOW, replacing in (5) $u(r) = y^2 p r^2 + \frac{1}{2} y^2 r^2 \left[\frac{1}{r} \left(-1 - \frac{1}{q^2} \log \left(\frac{r}{r} \right) \right) \right]$ u(r) = 1223-1 (13-1-202 log(r/F)) (6) So now from (6) we have a mean reverting drift a(1).

Now we need to examine I, martetprice of risk. There is no information in the poess. We examine the shart end of the come.

So, we know that zero-coupon satisfy:
$$\frac{dz}{dt} + \int w^2 \frac{d^2z}{dr^2} + (u - \lambda w) \frac{dz}{dr} - rz = 0$$
 (7)

bing Taylor inT=E => Z~ 1+a(v) (T-E)+b(+) (T-E) 2+....

In question (2) we do all the procedure, and show that: (replacing Taylor in (71)) $\frac{2 \times (-r(t-\epsilon))}{2} + \frac{1}{2}(t-\epsilon)^2 (r^2 + 1) + \dots$

We also show that, to find the spot rate we need: V = -ln(2)where l is the spot rate we need: V = -ln(2)

vow, for small (T-E) and assuming &(1+x) xx near x=0, we also show that =

$$\frac{-\ln(2)}{(T-\epsilon)} \sim V + \frac{1}{2} (T-\epsilon) (u - \lambda w) + \cdots (8)$$

Now, from the data, we can examine the time-sense data to estimate $(a-\lambda w)$ empirically. Since we have u and w alwordy, we can find λ .

Here we assume that u, w and \(\) only depend on \(Y\). Empirically, we can see that \(\) is time vary, nog,

Spot vote model:
$$dr = (\eta - \mu r) dt + (dr + B) dw$$
 (Parametes η, μ, α, β , constant)

For the following bond priving equation:
$$\frac{dz}{dt} + \frac{1}{2}w^2 \frac{d^2z}{dt^2} + (y-\lambda w)\frac{dz}{dt} - tz = 0$$
 (1)

In this case:
$$(u-\lambda w) = (\gamma - \mu r)$$

 $w = (\lambda r + \beta)^{1/2}$

Replacing on (1):
$$\frac{dz}{dt} + \frac{1}{2} \left(dr + \beta \right) \frac{d^2z}{dr^2} + \left(\eta - \mu r \right) \frac{dz}{dr} - 1z = 0$$
 (2) With Lincol Londinion

We call take decidation 1: 1

$$\frac{d^{2}}{dt} = \left(\frac{dA}{dt} - rdB\right) \cdot e^{A-rB} = \left(\frac{dA}{dt} - rdB\right) \cdot Z$$

$$\frac{d^{2}}{dr} = -Be^{A-rB} = -BZ$$

$$\frac{d^{2}}{dr^{2}} = B^{2}e^{A-rB} = BZ$$

$$\left(\frac{\partial A}{\partial t} - r\frac{\partial B}{\partial t}\right) \cdot \frac{1}{2} + \frac{1}{2} \left(\partial r + \beta^2\right) \cdot B^2 + \left(\eta - \mu r\right) \cdot \left(-B \cdot \frac{1}{2}\right) - r \cdot \frac{1}{2} = 0$$

Dividing by Z and le-awanging by grouping "r" terms:

$$\left(\frac{JA}{Jt} + \frac{J^3}{2} \cdot B^2 - \eta^B\right) + r\left(-\frac{JB}{Jt} + \frac{\alpha B^2}{2} + \mu B - 1\right) = 0$$
 (3)

Now, with v \$0, expressions between premises must be zero to fullfill the equation (3). So we

$$\frac{\partial A}{\partial t} + \frac{B \cdot B^2}{2} - \eta B = 0 \Rightarrow \frac{\partial A}{\partial t} = \eta B - \frac{1}{2} \beta B^2$$
 (4)

$$-\frac{JB}{Jt} + \frac{\Delta B^2}{2} + \gamma B - 1 = 0 \qquad \Rightarrow \qquad \frac{JB}{Jt} = \frac{1}{2} \Delta B^2 + \gamma B - 1 \quad (5)$$

Now, we know that at maturity zero-corpor bond is worth 1: Z(K,T,T=1 Hen eACH)-rB(F) = 1 ACTIT- + BCTIT) = 0 (6) then (ACTIT) = 0 = BCTIT) (40rv+0) So we now need to solve: $\frac{JB}{Jt} = \frac{1}{2} dB^2 + yB - 1$ with condition A(T,T) = B(T,T) = O(T,T)Solving: $\frac{2}{d} \cdot \frac{dB}{dt} = B^2 + \frac{2\mu B}{d} - \frac{2}{d}$ $\Rightarrow \frac{dB}{(B^2 + 2\mu B - \frac{2}{d})} = \frac{\alpha}{2} dt$ Roots: $B_{1,2} = \frac{-2\gamma}{d} + \sqrt{4\gamma^2 - 4.1 \cdot \left(-\frac{2}{d}\right)}$ Solving roots $= \frac{-2y}{2} + \sqrt{\frac{4}{2}(y^2 + 2d)} = \frac{2}{2}(-y + \sqrt{y^2 + 2d})$ $B_{1,2} = -\gamma + \sqrt{\gamma^2 + 2\lambda}$ So now we have: dB = adt $(B-B_1)(B-B_2) = 2dt$ Now, for integration purposes, let's define: a = B, $b = -B_2$ $\begin{cases} a,b = \pm y + \sqrt{y^2 + 2d} \end{cases}$ Hom: $\frac{dB}{(B-a)(B+b)} = \frac{d}{2} dt$ $\Longrightarrow \int \frac{dB'}{(B'-q)(B'+b)} = \frac{d}{2} dt$ where B' is articary variable for integration $\implies \log \left(\frac{B-a}{b+B} \right) = \frac{\alpha}{2} \left(\xi - T \right) \quad (6)$

Now we have:
$$\frac{(1-e^{4(4-1)})(4-1)(4+1)}{d(1+4-(1-4)e^{4(4-1)})} = \frac{(1-e^{4(4-1)})(4^2-1^2)}{d(1+4-(1-4)e^{4(4-1)})}$$

But, we man that:
$$V_1 = \sqrt{\mu^2 + 2d} \Rightarrow 50 \quad V_1^2 - \mu^2 = 2d$$
 (2)

Replacing:
$$(1-e^{4(4-1)})$$
 $= 2(1-e^{4(4-1)})$ $= 2(1-e^{4(4-1)})$ $= 2(1-e^{4(4-1)})$ $= 2(1-e^{4(4-1)})$ $= 2(1-e^{4(4-1)})$

le ananging:
$$B = \frac{2(e^{V_1(1-V_1)})}{-y_1-y_1+y_2e^{V_1(1-V_1)}} + \frac{2(e^{V_1(1-V_1)})}{-2y_1e^{V_1(1-V_1)}}$$

$$B = 2(e^{y(t-1)})$$

$$(y+y_1)(e^{y_1(t-1)}) - 2y_1e^{y_1(t-1)}$$

$$B = \frac{2(1 - e^{-\psi_{i}(t+1)})}{(\mu + \psi_{i})(1 - e^{-\psi_{i}(t+1)}) - 2\psi_{i}} / (-1)$$

$$B = \frac{2(e^{-\psi_{i}(t-T)}-1)}{(y+\psi_{i})(e^{-\psi_{i}(t-T)}-1)+2\psi_{i}}$$

Then:
$$B = 2(e^{V_1(T-t)})$$

$$(y_1+V_1)(e^{V_1(T-t)}-1)+2V_1$$

How defining:
$$V_{i} = \sqrt{\mu^{2} + 2d}$$
 and exaliating (6)
$$\log \left[\frac{B - (y + v_{i})}{a}\right] - \log \left[\frac{O + y - v_{i}}{y + v_{i}}\right] = \frac{\alpha}{2} (+T) / \frac{2v_{i}}{a}$$

$$= \frac{\alpha}{2} (+T) / \frac{2v_{i}}{a}$$

$$= \frac{2v_{i}}{a}$$

$$\log \left[\frac{\alpha B + y - v_{i}}{a B + y + v_{i}}\right] - \log \left[\frac{y - v_{i}}{y + v_{i}}\right] = v_{i} (+T)$$

$$\log \left[\frac{(\alpha B + y - v_{i})}{(\alpha B + y + v_{i})} \cdot \frac{(y + v_{i})}{(y - v_{i})}\right] = v_{i} (+T) / e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y + v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_{i}}{\alpha B + y + v_{i}} = \frac{(y - v_{i})}{(y - v_{i})} \cdot e^{(t)}$$

$$\frac{\alpha B + y - v_$$

JI+W.

```
Q5
   Two factor in lest vale model:
                                                                                                                    dr= u(r,t)dt+ w(r,t)dW,(t)
                                                                                                                           d\ell = p(r_1 t) dt + q(r_1 t) dW_2(t)
                                                                                                                       #[dw,dw2]= pdt
Bond with moderly T. V (r, l, t; T)
q T=V(Y, l,t;T)-Δ,V, (Y, l,t;T)-Δ2V2(Y, l,t;T2) Drowtfolio hodged by the
Now dT = dV(v,l,t;T) - \Delta, dV, (v,l,t;T_1) - \Delta dV_2(v,l,t;T_2) (1)
Eq(2) is by Itô, dt and was temos from dt 20
Now. dv^2 = u(v,t)^2 dt^2 + w(v,t)^2 dw_1^2 + 2u(v,t)w(v,t) dt dw_1

dv^2 = w(v,t)^2 dt \quad (3)
e(dt^3 \hbar) \approx 0
lau dl2= p(r, e)2de2+q(r, e)2dw2+2p(r, e)q(r, e)dedw2
          dl2= govisde (4)
dvdl = a(x+)p(x+)de2+ a(x+)q(x+)dtdwz + w(x+)dw, p(x+)dt + w(x+)q(x+)dw, dwz
             dudl= w(r,t) q(r,t) dw, dwz
             But . #[dudin] = Pdt
           s: drdl=w(rit)q(rit) pdt (5)
low, replacing in (2)
dV = \frac{dV}{dt} \frac{dt}{dr} + \frac{dV}{dr} \frac{dr}{dt} + \frac{dV}{dl} \frac{dl}{dt} + \frac{1}{2} \frac{d^2V}{dr} \frac{w(v,t)^2 dt}{dt} + \frac{1}{2} \frac{d^2V}{dl} \frac{w(v,t)^2 dt}{dt} + \frac{1}{2} \frac{d^2V}{dl} \frac{w(v,t)^2 dt}{dt} + \frac{1}{2} \frac{d^2V}{dl} \frac{w(v,t)^2 dt}{dt} + \frac{1}{2} \frac{d^2V}{dt} \frac{w(v,t)^2 dt}{dt} + \frac{1}{
```

So:
$$dV = \left(\frac{dV}{dt} + \frac{1}{2}\frac{d^2V}{dv^2}w(v,t)^2 + \frac{1}{2}\frac{d^2V}{dt^2}q(v,t)^2 + \frac{J^2V}{dv}w(v,t)q(v,t)\rho\right)dt + \frac{dV}{dt}dr + \frac{dV}{dt}dt$$

$$dV = \int (V)dt + \frac{dV}{dt}dt + \frac{dV}{dt}dt + \frac{dV}{dt}dt$$

$$dV = \int (V)dt + \frac{dV}{dt}dt + \frac{dV}{dt}dt$$

NOW, Apply 196) for V, and Vz, then replacing in (1), we have:

$$dT = \int (V)dt + \frac{dV}{dr} \frac{dV}{dr} + \frac{dV}{dr} \frac{dV}{dr} - \Delta_1 \left(\int (V_1)dt + \frac{dV}{dr} \frac{dr}{dr} \frac{dV}{dr} \right) - \Delta_2 \left(\int (V_2)dt + \frac{dV}{dr} \frac{dr}{dr} + \frac{dV}{dr} \frac{dr}{dr} \right)$$

$$dV_2$$

bow, grouping Previous equation dil

$$dT = \left(\int (V) - \Delta_{1} \int (V_{1}) - \Delta_{2} \int (V_{2}) dt + \left(\frac{\partial V}{\partial r} - \Delta_{1} \frac{\partial V_{1}}{\partial r}\right) dr + \left(\frac{\partial V}{\partial l} - \Delta_{1} \frac{\partial V_{1}}{\partial l}\right) dl$$

$$B$$

By No-ARBITRAGE PRINCIPLE, a visk FACE ASSET MUST FAVOR HE HISK FACE PATE.

So, that we eliminate how (randomness) by doing B=0 and C=0:

$$\frac{\partial V}{\partial r} - \Delta_1 \frac{\partial V_1}{\partial r} - \Delta_2 \frac{\partial V_2}{\partial r} = 0$$

$$\frac{\partial V}{\partial l} - \Delta_1 \frac{\partial V_1}{\partial l} - \Delta_2 \frac{\partial V_2}{\partial r} = 0$$

$$\frac{\partial V}{\partial l} - \Delta_1 \frac{\partial V_1}{\partial l} - \Delta_2 \frac{\partial V_2}{\partial l} = 0$$
(8)

then, the put-tolio must garm the hise trace PATE, So Alt = LITCH to The V-DIV, -DIV, -DIV

$$\int (V) - \Delta_1 \int (V_1) - \Delta_2 \int (V_2) = Y \left(V - \Delta_1 V_1 - \Delta_2 V_2 \right)$$

Reducinging:
$$\int (V) - rV = \Delta_1 \left(\int (V_1) - V_1 r \right) + \Delta_2 \left(\int (V_2) - V_2 r \right)$$

we have:
$$\int'(V) = \Delta_i \int'(V_i) + \Delta_2 \int'(V_2)$$

So finally:
$$\int (V) - \Delta_1 \int (V_1) - \Delta_2 \int (V_2) = 0$$
 (9)

Now, we have Beguations (7,8 and 9), and two unknowns D, and Dz, deliming an inlansistent system (over-possised)

Now, the system can be written as:

$$\begin{pmatrix}
S'(v) & S'(v_1) & S'(v_2) \\
Jv_{JL} & Jv_{JL} & Jv_{ZJL} \\
Jv_{JL} & Jv_{JL} & Jv_{ZJL}
\end{pmatrix}
\begin{pmatrix}
I \\
-\Delta_1 \\
-\Delta_2
\end{pmatrix} = \begin{pmatrix}
O \\
O \\
O
\end{pmatrix}$$

So we now set det(M)=0, this means that the first now will be a linear compination of the other two:

$$\int_{V}^{1}(v) = \alpha_{r} \frac{JV}{Jr} + \alpha_{l} \frac{JV}{Jl}$$

Now, defining of and all interms of market price of risk Ar (V, I, t) and Al (V, I, t) associated with vould: «1= >19-P

We have:

splacing S(V):

$$\frac{\partial V}{\partial t} + \frac{1}{2} \frac{\partial^{2} V}{\partial r^{2}} + \frac{1}{2} \frac{\partial^{2} V}{\partial z^{2}} + \frac{1}{2} \frac{\partial^{2} V}{\partial$$

$$\frac{dV}{dt} + \frac{1}{2} \frac{d^2 v}{dr^2} w^2 + \frac{1}{2} \frac{d^2 q}{dr^2} + \frac{d^2 v}{dr} wq p - W + (u - \lambda r w) \frac{dV}{dr} + (p - \lambda q) \frac{dV}{dl} = 0$$
 (10)

So, given:
$$u-\lambda_{r}w=0=p-\lambda_{r}q$$

and: $w=q=\sqrt{a+br+c}l$ with a,b,c (and the stands).

$$\frac{dV}{dt} + \frac{1}{2} \frac{J^2 V}{dt^2} (a + br + cl) + \frac{1}{2} \frac{d^2 V}{dl^2} (a + br + cl) + \frac{1}{2} \frac{J^2 V}{dl} (a + br + cl) + \frac{1}{2} \frac{J^2 V}{dl} (a + br + cl) + \frac{1}{2} \frac{J^2 V}{dl} = 0$$

Solution of the town:
$$V = \exp(A(t;T) - tB(t;T) - lC(t;T))$$
 (Redemption Value: $V(t;lT;T) - l$

Redemption Valle:
$$V(t) = \exp(A(t;T) - tB(t;T) - lC(t;T))$$

Redemption Value:
$$V = \exp(A(t;T) - tB(t;T) - lC(t;T))$$
 (12)

 $V(t;l;T;T) = 1$
 $V(t;l;T;T) = 0$
 $V(t;$

$$\frac{\partial V}{\partial t} = \left(\frac{\partial A}{\partial t} - r \frac{\partial B}{\partial t} - l \frac{\partial C}{\partial t}\right) e^{A - rB - lC} = \left(\frac{\partial A}{\partial t} - r \frac{\partial B}{\partial t} - l \frac{\partial C}{\partial t}\right) \cdot V$$

$$\frac{d\hat{V}}{dv^2} = B^2 e^{A-VB-QC} = B^2 V$$

$$\frac{dV}{dt} = BCe = BCV$$

$$\left(\frac{dA}{dt} - r\frac{dB}{dt} - l\frac{dC}{dt}\right) \cdot V + \frac{1}{2} \left(B^2V\right) \left(a + br + cl\right) + \frac{1}{2} \left(CV\right) \left(a + br + cl\right) + BCV\left(a + br + cl\right) - VV = 0$$

how dividing by V and graping rand terms:

$$V\left(-\frac{JB}{Jt} + \frac{B^{2}b}{2} + \frac{C^{2}b}{2} + BCb - 1\right) + \sqrt{-\frac{JC}{Jt} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + BCc} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + BCc + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + BCc - \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{B^{2}c}{2} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{C^{2}c}{2} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{C^{2}c}{2} + \frac{C^{2}c}{2} + \frac{C^{2}c}{2} + \frac{B^{2}c}{2} + \frac{C^{2}c}{2} + \frac{C^{2}c$$

Now, with 140, lto, aspressions between prenderes mist be zero to fulfil the equation (B).

$$\frac{-JB}{Jt} + \frac{B^{2}b}{2} + \frac{C^{2}b}{2} + BCb - 1 = 0 \implies \frac{JB}{Jt} = \frac{B^{2}b}{2} + \frac{C^{2}b}{2} + BCb - 1$$

$$\frac{JC}{Jt} + \frac{B^{2}c}{2} + \frac{C^{2}b}{2} + \frac{BCb}{2} - 1$$

$$\frac{JC}{JE} + \frac{B^{2}}{2} + \frac{Z}{2} + BC_{c} = 0$$

$$\frac{JC}{JE} + \frac{B^{2}}{2} + \frac{C_{c}}{2} + BC_{c}$$

$$\frac{JC}{JE} + \frac{B^{2}}{2} + \frac{C_{c}}{2} + BC_{c}$$

$$\frac{JA}{Jt} + \frac{B_q}{2} + \frac{C_q}{2} + \frac{BC_q}{2} = 0$$
 $\Rightarrow \frac{JA}{Jt} = -\frac{B^2q}{2} - \frac{C_q}{2} - \frac{BC_q}{2}$

With boundary and its size A

With boundary conditions: A(T;T) = B(T;T) = C(T; +) = 0 (moreoing condition: V(Y, I, T;T)=1)