


## 16.7. Surface integrals: vector fields

★ Def Given an orientable surface  $S$  parametrized by  $\vec{r}(u,v)$  on a domain  $D$ , the surface integral (or flux) of a vector field  $\vec{F}$  over  $S$  is


$$\iint_S \vec{F} \cdot d\vec{S} := \iint_D \vec{F}(\vec{r}(u,v)) \cdot (\vec{r}_u \times \vec{r}_v) dA$$

Note (1) If  $\vec{F}$  is a velocity field of a fluid, then the flux


$\iint_S \vec{F} \cdot d\vec{S}$  is the rate of flow across  $S$ .



(2) The orientation of  $S$  is determined by the direction of normal vectors



oriented upward



oriented downward

★ (3) The surface integral of a vector field depends on the orientation of the surface

$$\iint_S \vec{F} \cdot d\vec{S} = - \iint_{-S} \vec{F} \cdot d\vec{S}$$

where  $-S$  is the surface  $S$  with the opposite orientation.


(4) If  $\vec{n}$  denotes the unit normal vector of  $S$ , then

$$\iint_S \vec{F} \cdot d\vec{S} = \iint_S \underbrace{\vec{F} \cdot \vec{n}}_{\text{Scalar}} dS$$

★ This is useful if  $S$  is a sphere or a flat surface.

Ex Let  $S$  be the part of the plane  $x+y+z=1$  that lies in the first octant. Find the upward flux of the vector field  $\vec{F}(x,y,z) = (z, x^2, x+y)$  across  $S$ .

Sol



$$x+y+z=1 \rightsquigarrow z=1-x-y.$$

$S$  is parametrized by  $\vec{r}(x,y) = (x, y, 1-x-y)$

The domain  $D$  is given by  $0 \leq x \leq 1, 0 \leq y \leq 1-x$ .

$$\vec{r}_x = (1, 0, -1), \quad \vec{r}_y = (0, 1, -1)$$

$$\Rightarrow \vec{r}_x \times \vec{r}_y = (1, 1, 1) : \text{oriented upward}$$


$$\iint_S \vec{F} \cdot d\vec{S} = \iint_D \vec{F}(\vec{r}(x,y)) \cdot (\vec{r}_x \times \vec{r}_y) dA$$

$$\vec{F}(\vec{r}(x,y)) \cdot (\vec{r}_x \times \vec{r}_y) = (1-x-y, x^2, x+y) \cdot (1, 1, 1) = x^2 + 1$$

$$\begin{aligned} \Rightarrow \iint_S \vec{F} \cdot d\vec{S} &= \int_0^1 \int_0^{1-x} x^2 + 1 \, dy \, dx = \int_0^1 (1-x)(x^2+1) \, dx \\ &= \int_0^1 -x^3 + x^2 - x + 1 \, dx = \left( -\frac{x^4}{4} + \frac{x^3}{3} - \frac{x^2}{2} + x \right) \Big|_{x=0}^{x=1} \\ &= \boxed{\frac{7}{12}} \end{aligned}$$

Ex Let  $S$  be the part of the paraboloid  $z = 2 - x^2 - y^2$  that lies in the cone  $z^2 = x^2 + y^2$ . Find the downward flux of the vector field  $\vec{F}(x, y, z) = (y, -x, z)$  across  $S$ .

Sol



$S$  is parametrized by  $\vec{r}(x,y) = (x, y, 2-x^2-y^2)$

The domain  $D$  is given by  $x^2 + y^2 \leq 1$ .

$$\vec{r}_x = (1, 0, -2x), \quad \vec{r}_y = (0, 1, -2y)$$

$$\Rightarrow \vec{r}_x \times \vec{r}_y = (2x, 2y, 1) : \text{oriented upward}$$

$$\iint_S \vec{F} \cdot d\vec{S} = - \iint_D \vec{F}(\vec{r}(x,y)) \cdot (\vec{r}_x \times \vec{r}_y) dA$$

↑  
opposite orientation

$$\vec{F}(\vec{r}(x,y)) \cdot (\vec{r}_x \times \vec{r}_y) = (y, -x, 2-x^2-y^2) \cdot (2x, 2y, 1) = 2-x^2-y^2.$$

In polar coordinates,  $D$  is given by  $0 \leq \theta \leq 2\pi$ ,  $0 \leq r \leq 1$ .


$$\begin{aligned} \iint_S \vec{F} \cdot d\vec{S} &= - \iint_D 2 - x^2 - y^2 \, dA = - \int_0^{2\pi} \int_0^1 (2 - r^2) r \, dr \, d\theta \quad \text{Jacobian} \\ &= - \int_0^{2\pi} \left. r^2 - \frac{r^4}{4} \right|_{r=0}^{r=1} d\theta = - \int_0^{2\pi} \frac{3}{4} d\theta = \boxed{-\frac{3\pi}{2}} \end{aligned}$$

★ Ex Consider the inverse square field

$$\vec{F}(x,y,z) = \left( \frac{x}{(x^2+y^2+z^2)^{3/2}}, \frac{y}{(x^2+y^2+z^2)^{3/2}}, \frac{z}{(x^2+y^2+z^2)^{3/2}} \right)$$

Find  $\iint_S \vec{F} \cdot d\vec{S}$  where  $S$  is a sphere centered at the origin, oriented outward.

Sol



Let  $R$  be the radius of  $S$   
 $\Rightarrow S$  is given by  $x^2 + y^2 + z^2 = R^2$   
 $\leadsto$  a level surface of  
 $f(x,y,z) = x^2 + y^2 + z^2$   
 $\Rightarrow$  A normal vector of  $S$  is  
 $\nabla f = (2x, 2y, 2z)$

The unit normal vector of  $S$  is given by

$$\vec{n} = \frac{\nabla f}{|\nabla f|} = \frac{(2x, 2y, 2z)}{\sqrt{4x^2 + 4y^2 + 4z^2}} = \frac{(2x, 2y, 2z)}{2R} = \left( \frac{x}{R}, \frac{y}{R}, \frac{z}{R} \right)$$

$\vec{n}$  is oriented outward (pointing away from the origin)

$$\iint_S \vec{F} \cdot d\vec{S} = \iint_S \vec{F} \cdot \vec{n} \, dS$$

$$\begin{aligned} \vec{F} \cdot \vec{n} &= \frac{x^2}{R(x^2+y^2+z^2)^{3/2}} + \frac{y^2}{R(x^2+y^2+z^2)^{3/2}} + \frac{z^2}{R(x^2+y^2+z^2)^{3/2}} \\ &= \frac{x^2+y^2+z^2}{R(x^2+y^2+z^2)^{3/2}} = \frac{R^2}{R \cdot R^3} = \frac{1}{R^2} \end{aligned}$$

$\uparrow$   
 $x^2+y^2+z^2 = R^2$  on  $S$

$$\Rightarrow \iint_S \vec{F} \cdot d\vec{S} = \iint_S \frac{1}{R^2} \, dS = \frac{1}{R^2} \text{Area}(S) = \frac{1}{R^2} \cdot \underbrace{4\pi R^2}_{\text{Area of sphere}} = \boxed{4\pi}$$