


## General idea of a perturbation theory

Stationary State Perturbation Theory  $\int$  ref. Davydov: ¶ 47-50,  
Sakurai: Secs. 5.1-5.2.

The key idea of a perturbation theory is to start with a well-defined and well-known system, add a small change (i.e. perturbation), and then develop a method for specifying the presumably small excursions in system parameters.

In this general sense, the WKB approximation is a perturbation theory -- we start from free particle eigenfns & eigenvalues (or from  $\psi^s$  &  $E^s$  in a const external potential), and document how things change when we add "small" (remember  $|k'/k^2| \ll 1$ ) departures from a const potential.

A classical example<sup>¶</sup> of a perturbation is a string of length  $l$  & uniform mass density, pegged at both ends and set into vibration. The wave amplitudes  $\psi_n$  and eigenfrequencies  $\omega_n$  are easy to get. Now a small point mass  $m$  (small means  $m \ll$  string mass) is attached to the string -- this changes  $\psi_n \rightarrow \psi'_n$  and  $\omega_n \rightarrow \omega'_n$  by small amounts. The problem is to calculate the new  $\psi'_n$  &  $\omega'_n$  from the old (unperturbed)  $\psi_n$  &  $\omega_n$ .



1) We shall now do the QM version of the above string problem, viz.

Suppose a stationary (i.e. time-independent), bound-state (i.e. discrete energies) QM system with known eigenfns  $\psi_n^{(0)}$  & eigenenergies  $E_n^{(0)}$  is described by:  $\mathcal{H}_0(p, x) \psi_n^{(0)}(x) = E_n^{(0)} \psi_n^{(0)}(x)$ . A "small" time-independent potential  $V(x)$  is added to  $\mathcal{H}_0$ , i.e.  $\mathcal{H}_0 \rightarrow \mathcal{H} = \mathcal{H}_0 + V$ . The new QM problem is:  $\mathcal{H} \psi_n = E_n \psi_n$ . How are the perturbed  $(\psi_n, E_n)$  related to the known  $(\psi_n^{(0)}, E_n^{(0)})$ ?

(1)

<sup>¶</sup> See Sec. 44 of A. Fetter & J. Walecka "Theoretical Mechanics..." (McGraw-Hill, 1980)

## Basic perturbation equation.

SS(2)

### REMARKS

1. For the unperturbed system, the  $\{\psi_n^{(0)}\}$  are a complete orthonormal set. The energies  $E_n^{(0)}$  are assumed non-degenerate (we'll treat degeneracy later).
  2.  $V(x)$  "small" means  $|\langle \psi_n^{(0)} | V | \psi_n^{(0)} \rangle| \ll |E_n^{(0)}|$  within  $n^{\text{th}}$  state, or when  $m \neq n$ :  $|\langle \psi_m^{(0)} | V | \psi_n^{(0)} \rangle| \ll |E_m^{(0)} - E_n^{(0)}|$ . So  $V$  cannot change the spectrum much.
- 

2) With  $V$  added in, our perturbed problem is:

$$\mathcal{H}(p, x) \psi_k(x) = E_k \psi_k(x), \quad \mathcal{H}(p, x) = \mathcal{H}_0(p, x) + V(x).$$

PROBLEM: Find  $\psi_k$  and  $E_k$  in terms of  $\psi_n^{(0)}$ ,  $E_n^{(0)}$  &  $V$ . (2)

Since  $\{\psi_n^{(0)}\}$  are a complete set, we can expand the unknown  $\psi_k$  as...

$$\rightarrow \psi_k(x) = \sum_n a_{nk} \psi_n^{(0)}(x), \quad \text{w/ } a_{nk} = \langle \psi_n^{(0)} | \psi_k \rangle \quad \text{at this point, the } a_{nk} \text{ are unknown.} \quad (3)$$

Plug this version of  $\psi_k$  into  $\mathcal{H} \psi_k = E_k \psi_k, \dots$

$$\sum_n a_{nk} (\mathcal{H}_0 + V) \psi_n^{(0)} = \sum_n a_{nk} E_k \psi_n^{(0)} \quad \leftarrow \text{use } \mathcal{H}_0 \psi_n^{(0)} = E_n^{(0)} \psi_n^{(0)}, \text{ and rearrange terms...}$$

or

$$\sum_n a_{nk} (E_k - E_n^{(0)}) \psi_n^{(0)} = \sum_n a_{nk} V \psi_n^{(0)}. \quad (4)$$

Operate thru Eq. (4) by  $\langle \psi_m^{(0)} |$ , and use  $\langle \psi_m^{(0)} | \psi_n^{(0)} \rangle = \delta_{mn}$ . Then...

$(E_k - E_m^{(0)}) a_{mk} = \sum_n V_{mn} a_{nk}$

, w/  $V_{mn} = \langle \psi_m^{(0)} | V | \psi_n^{(0)} \rangle. \quad (5)$

This is the Fundamental Equation of QM SS perturbation theory. It allows us -- at least in principle -- to solve for the set of coefficients  $\{a_{nk}\}$  (which specify  $\psi_k = \sum_n a_{nk} \psi_n^{(0)}$ ) from an  $\infty$  set of linear coupled equations. The energies  $E_k$  are gotten from a secular eqn, as follows...

Existence of solns  $\{\psi_k, E_k\}$  from  $\{\psi_n^{(0)}, E_n^{(0)}, V_{mn}\}$ .

SS13

let  $\rightarrow \underline{a}_k = \begin{pmatrix} \vdots \\ a_{nk} \\ \vdots \end{pmatrix}$ ,  $\underline{a}_{nk} = \langle \psi_n^{(0)} | \psi_k \rangle$  [a rep<sup>n</sup> of  $\psi_k$  on basis  $\{\psi_n^{(0)}\}$ ]; (6a)

$\rightarrow \underline{V} = \begin{pmatrix} \vdots \\ \dots V_{mn} \dots \\ \vdots \end{pmatrix}$ ,  $\underline{V}_{mn} = \langle \psi_m^{(0)} | V | \psi_n^{(0)} \rangle$  [rep<sup>n</sup> of  $V$  on  $\{\psi_n^{(0)}\}$ ]; (6b)

$\rightarrow \underline{\Delta}_k = \begin{pmatrix} \vdots \\ \dots (E_k - E_m^{(0)}) \delta_{mn} \dots \\ \vdots \end{pmatrix}$ , a diagonal energy matrix. (6c)

Then Eq. (5) reads, in these terms...

$\left[ \begin{array}{l} \underline{\Delta}_k \underline{a}_k = \underline{V} \underline{a}_k, \text{ or } (\underline{\Delta}_k - \underline{V}) \underline{a}_k = 0; \\ \text{and } \det(\underline{\Delta}_k - \underline{V}) = \det[(E_k - E_m^{(0)}) \delta_{mn} - V_{mn}] = 0. \end{array} \right.$  Secular equation. (7)

The secular eqn is imposed to avoid the trivial solution  $\underline{a}_k \equiv 0$ . It will evidently give a specific  $E_k$  in terms of all the  $E_m^{(0)}$  &  $V_{mn}$ . Then, with the  $\{E_k\}$ , we can find the (normalized) eigenvectors  $\{\underline{a}_k\}$  as usual.

This statement of the perturbation problem in Eq. (5) shows that solutions  $\{\psi_k, E_k\}$  exist, and they can be gotten from the  $\{\psi_n^{(0)}, E_n^{(0)}\}$  and  $V$  alone. But Eq. (7) is not a practical solution, since it involves  $\infty \times \infty$  matrices like  $\{V_{mn}\}$  and  $\infty$ -component eigenvectors  $\underline{a}_k$ .

3) A practical method of dealing with Eq. (5) capitalizes on the "smallness" of  $V$  and the notion we can correct for it by a series of ever smaller terms. Let:

$V \rightarrow \lambda V$   $\left\{ \begin{array}{l} \text{parameter } \lambda = 0 \Rightarrow \text{no perturbation; solns are } \psi_n^{(0)} \text{ \& } E_n^{(0)}; \\ \lambda = 1 \Rightarrow \text{full perturbation; solns are } \psi_k \text{ \& } E_k. \end{array} \right.$  (8)

## Expansion of Fund<sup>l</sup> Eq. (5) in terms of order parameter $\lambda$

SS14

In what follows, we will always take  $\lim_{\lambda \rightarrow 1}$  in the final result, to get the full solution of the  $(\mathcal{H}_0 + V)$  problem.  $\lambda$  is just a convenient parameter to keep track of what "order-of-approximation" we are doing; this will soon become clear.

If  $V$  is "small",  $E_k$  must be close to  $E_k^{(0)}$ , so we write...

$$\rightarrow E_k = E_k^{(0)} + \lambda E_k^{(1)} + \lambda^2 E_k^{(2)} + \dots = \sum_{\mu=0}^{\infty} \lambda^{\mu} E_k^{(\mu)} \quad \int \lim_{\lambda \rightarrow 1} \text{ is understood.} \quad (9)$$

$E_k^{(\mu)}$  is called the  $\mu^{\text{th}}$  order correction to  $E_k^{(0)}$  [for  $\mu \geq 1$ ].

Hope is: if  $V$  is "small", then  $|E_k^{(\mu+1)}| \ll |E_k^{(\mu)}|$ , so series converges.

In the same spirit, we expand the unknown  $a_{nk}$ 's in Eq. (5), viz...

$$\rightarrow a_{nk} = a_{nk}^{(0)} + \lambda a_{nk}^{(1)} + \lambda^2 a_{nk}^{(2)} + \dots = \sum_{\nu=0}^{\infty} \lambda^{\nu} a_{nk}^{(\nu)} \quad \int \lim_{\lambda \rightarrow 1} \text{ is understood.} \quad (10)$$

Put the  $\lambda$ -expansions of Eqs. (9) & (10) into the Fundamental Equation (5);

$$\rightarrow (E_k - E_m^{(0)}) a_{mk} = \sum_n \lambda V_{mn} a_{nk}, \text{ becomes:}$$

$$\left( \sum_{\mu=0}^{\infty} E_k^{(\mu)} \lambda^{\mu} \right) \left( \sum_{\nu=0}^{\infty} a_{mk}^{(\nu)} \lambda^{\nu} \right) - \sum_{\nu=0}^{\infty} E_m^{(0)} a_{mk}^{(\nu)} \lambda^{\nu} = \sum_n \lambda V_{mn} \left( \sum_{\nu=0}^{\infty} a_{nk}^{(\nu)} \lambda^{\nu} \right). \quad (11)$$

Use general formula for product of power series

$$\left( \sum_{\mu=0}^{\infty} A_{\mu} \lambda^{\mu} \right) \left( \sum_{\nu=0}^{\infty} B_{\nu} \lambda^{\nu} \right) = \sum_{\mu=0}^{\infty} C_{\mu} \lambda^{\mu}, \quad C_{\mu} = \sum_{\sigma=0}^{\mu} A_{\mu-\sigma} B_{\sigma}$$

so/ 1<sup>st</sup> term LHS in Eq. (11) =

$$\sum_{\mu=0}^{\infty} \left( \sum_{\sigma=0}^{\mu} E_k^{(\mu-\sigma)} a_{mk}^{(\sigma)} \right) \lambda^{\mu},$$

and Eq. (11) can be written...

$$\left[ \sum_{\mu=0}^{\infty} \left[ \sum_{\sigma=0}^{\mu} E_k^{(\mu-\sigma)} a_{mk}^{(\sigma)} - E_m^{(0)} a_{mk}^{(\mu)} \right] \lambda^{\mu} = \sum_{\nu=0}^{\infty} \left[ \sum_n V_{mn} a_{nk}^{(\nu)} \right] \lambda^{\nu+1} \right] \quad (12)$$

This is now the Master Equation. It can be simplified, by the following steps:

## Master Eqn for Perturbation Theory as an Iteration

SS(5)

Simplifying Eq. (12)...

1. On the LHS, split off the  $\mu=0$  term, and set  $\sum_{\mu=1}^{\infty} = \sum_{\nu=0}^{\infty}$ , with  $\mu = \nu+1$ . Then...

$$\left\{ \begin{array}{l} \text{LHS of} \\ \text{Eq. (12)} \end{array} \right\} = (E_k^{(0)} - E_m^{(0)}) a_{mk}^{(0)} + \sum_{\nu=0}^{\infty} \left[ \sum_{\sigma=0}^{\nu+1} E_k^{(\nu+1-\sigma)} a_{mk}^{(\sigma)} - E_m^{(0)} a_{mk}^{(\nu+1)} \right] \lambda^{\nu+1}; \quad (13)$$

$= E_k^{(0)} a_{mk}^{(\nu+1)} + \sum_{\sigma=0}^{\nu} E_k^{(\nu+1-\sigma)} a_{mk}^{(\sigma)}.$

Take entire RHS of (12), and put over to LHS. Then (12) becomes...

$$\rightarrow (E_k^{(0)} - E_m^{(0)}) a_{mk}^{(0)} + \sum_{\nu=0}^{\infty} \left[ (E_k^{(0)} - E_m^{(0)}) a_{mk}^{(\nu+1)} + \sum_{\sigma=0}^{\nu} E_k^{(\nu+1-\sigma)} a_{mk}^{(\sigma)} - \right.$$

$$\left. \sum_n V_{mn} a_{nk}^{(\nu)} \right] \lambda^{\nu+1} = 0. \quad (14)$$

2. Consider Eq. (14) as a power series in  $\lambda$  for  $\lambda \neq 0$  ( $0 < \lambda \leq 1$  is OK). Since  $\lambda$  is an independently variable parameter, the only way the power series can  $= 0$  is if every one of its coefficients vanish, i.e.

$$\text{I. } (E_k^{(0)} - E_m^{(0)}) a_{mk}^{(0)} = 0; \quad (15a)$$

$$\text{II. } (E_k^{(0)} - E_m^{(0)}) a_{mk}^{(\nu+1)} + E_k^{(\nu+1)} a_{mk}^{(0)} = \sum_n V_{mn} a_{nk}^{(\nu)} - \sum_{\sigma=1}^{\nu} E_k^{(\nu+1-\sigma)} a_{mk}^{(\sigma)}. \quad (15b)$$

This extremely powerful simplification justifies the whole procedure of using power series in  $\lambda$ . Note that II is a recursion relation for  $a_{mk}^{(\nu+1)} = \text{fcn } a_{mk}^{(\mu \leq \nu)}$ .

3. In Eq. (15a),  $a_{mk}^{(0)} \equiv 0$  for  $m \neq k$  (the  $E_n^{(0)}$  are not degenerate). Then, for  $m=k$  we need  $a_{kk}^{(0)} = 1$ , so that  $\psi_k = \psi_k^{(0)}$  when  $V$  vanishes. So:  $a_{mk}^{(0)} = \delta_{mk}$ . Use of this in Eq. (15b), plus rewrite of last term RHS (let  $\mu = \sigma-1$ ) gives...

$$(E_k^{(0)} - E_m^{(0)}) a_{mk}^{(\nu+1)} + E_k^{(\nu+1)} \delta_{mk} = \sum_n V_{mn} a_{nk}^{(\nu)} - \sum_{\mu=0}^{\nu-1} E_k^{(\nu-\mu)} a_{mk}^{(\mu+1)}; \quad \nu=0,1,2,\dots \quad (16)$$

term contributes for  $\nu \geq 1$  only.

This is the new version of our Master Eq. for perturbation. The order parameter  $\nu=0,1,2,\dots$  corresponds to working on  $\mathcal{O}(\lambda)$ ,  $\mathcal{O}(\lambda^2)$ ,  $\mathcal{O}(\lambda^3)$  terms, i.e. to working on 1st, 2nd, 3rd order perturbative corrections.

4) Now we iterate the Master Eqn to get the  $a_{mk}^{(1)}$  from the already known  $a_{mk}^{(0)} = \delta_{mk}$ , the  $a_{mk}^{(2)}$  from the  $a_{mk}^{(1)}$  &  $a_{mk}^{(0)}$ , etc. Also the  $E_k^{(1)}$  from the  $E_k^{(0)}$ , etc. Like this:

In Eq. (16), Choose  $v=0$  ( $\Rightarrow$  working to  $\theta(\lambda)$ , 1<sup>st</sup> order perturbation theory).

$$\xrightarrow{\text{So}} (E_k^{(0)} - E_m^{(0)}) a_{mk}^{(1)} + E_k^{(1)} \delta_{mk} = \sum_n V_{mn} a_{nk}^{(0)} \xleftarrow{\delta_{nk}} = V_{mk}. \quad (17)$$

$$\underline{1. \ m \neq k, (17) \Rightarrow a_{mk}^{(1)} = V_{mk} / (E_k^{(0)} - E_m^{(0)})}. \quad (17a)$$

$$\text{So } \psi_k = \psi_k^{(0)} + \underbrace{\sum_m a_{mk}^{(1)} \psi_m^{(0)}}_{\text{"Small" correction only if}} + \dots \quad \int |a_{mk}^{(1)}| \ll 1, \quad |V_{mk}| \ll |E_k^{(0)} - E_m^{(0)}|$$

(as advertised at top of p. SS 2).

$$\underline{2. \ m=k, (17) \Rightarrow (0) + E_k^{(1)} = V_{kk} + (0), \text{ i.e. } E_k^{(1)} = V_{kk}. \quad (17b)$$

$$\text{So } E_k = E_k^{(0)} + \underbrace{V_{kk}}_{\text{"Small" only if}} + \dots \quad \int |V_{kk}| \ll |E_k^{(0)}|, \text{ within state } k.$$

(as advertised on top of p. SS 2).

NOTE: the  $m=k$  eqn for  $v=0$  gives no information on  $a_{kk}^{(1)}$ .

ASIDE: What value do we give  $a_{kk}^{(1)}$ ? Assertion: we can set  $a_{kk}^{(1)} = 0$ .

(1) We can defend the assertion on grounds that  $\psi_k$  should be normed:  $\langle \psi_k | \psi_k \rangle = 1$ .

The argument goes as follows. Write the wavefn  $\psi_k$  as ...

$$\left[ \begin{aligned} \psi_k &= \sum_n a_{nk} \psi_n^{(0)} = \sum_n \left( \sum_{\sigma=0}^{\infty} \lambda^{\sigma} a_{nk}^{(\sigma)} \right) \psi_n^{(0)} = \sum_{\sigma=0}^{\infty} \psi_k^{(\sigma)}, \\ \text{w/ } \psi_k^{(\sigma)} &= \lambda^{\sigma} \sum_n a_{nk}^{(\sigma)} \psi_n^{(0)} = \lambda^{\sigma} \left[ a_{kk}^{(\sigma)} \psi_k^{(0)} + \sum_n' a_{nk}^{(\sigma)} \psi_n^{(0)} \right]. \end{aligned} \right. \quad (18)$$

(prime means  $\sum_{n \neq k}$  (Sum w/o  $n=k$  term))

$\psi_k^{(\sigma)}$  is the  $\sigma^{\text{th}}$  order correction to  $\psi_k^{(0)}$ ; note  $\psi_k^{(\sigma)}|_{\sigma=0} = \psi_k^{(0)}$ , as should be (here use  $a_{nk}^{(0)} = \delta_{nk}$ ). Now we calculate the norm,  $\langle \psi_k | \psi_k \rangle$ , after splitting off  $\psi_k^{(0)}$ , i.e. put  $\psi_k = \psi_k^{(0)} + \sum_{\sigma=1}^{\infty} \psi_k^{(\sigma)}$  and calculate ...

# Normalization of perturbed wavefn $\psi_k$ .

SS(7)

$$\begin{aligned} \rightarrow \langle \psi_k | \psi_k \rangle &= \langle \psi_k^{(0)} + \sum_{\sigma=1}^{\infty} \psi_k^{(\sigma)} | \psi_k^{(0)} + \sum_{\mu=1}^{\infty} \psi_k^{(\mu)} \rangle \\ &= 1 + 2 \operatorname{Re} \sum_{\sigma=1}^{\infty} \underbrace{\langle \psi_k^{(0)} | \psi_k^{(\sigma)} \rangle}_{\textcircled{1}} + \sum_{\sigma, \mu=1}^{\infty} \underbrace{\langle \psi_k^{(\sigma)} | \psi_k^{(\mu)} \rangle}_{\textcircled{2}} \end{aligned} \quad (19)$$

(2) Use  $\psi_k^{(\sigma)}$  from Eq. (18) to evaluate the projections  $\textcircled{1}$  &  $\textcircled{2}$ . Thus...

$$\textcircled{1} = \lambda^{\sigma} \langle \psi_k^{(0)} | a_{kk}^{(\sigma)} \psi_k^{(0)} + \sum_n' a_{nk}^{(\sigma)} \psi_n^{(0)} \rangle = \lambda^{\sigma} a_{kk}^{(\sigma)} ; \quad (20a)$$

$\uparrow$  only contrib  $\uparrow$

$$\begin{aligned} \textcircled{2} &= \lambda^{\sigma+\mu} \langle a_{kk}^{(\sigma)} \psi_k^{(0)} + \sum_n' a_{nk}^{(\sigma)} \psi_n^{(0)} | a_{kk}^{(\mu)} \psi_k^{(0)} + \sum_m' a_{mk}^{(\mu)} \psi_m^{(0)} \rangle \\ &= \lambda^{\sigma+\mu} \left\{ a_{kk}^{(\sigma)*} a_{kk}^{(\mu)} + \sum_{n,m} a_{nk}^{(\sigma)*} a_{mk}^{(\mu)} \underbrace{\langle \psi_n^{(0)} | \psi_m^{(0)} \rangle}_{\delta_{nm}, \text{ now sum over } m} \right\}. \end{aligned} \quad (20b)$$

Put results of Eqs. (20) into Eq. (19) to get the  $\psi_k$  norm as...

$$\langle \psi_k | \psi_k \rangle = 1 + 2 \operatorname{Re} \sum_{\sigma=1}^{\infty} \lambda^{\sigma} a_{kk}^{(\sigma)} + \sum_{\sigma, \mu=1}^{\infty} \lambda^{\sigma+\mu} \left\{ a_{kk}^{(\sigma)*} a_{kk}^{(\mu)} + \sum_n' a_{nk}^{(\sigma)*} a_{nk}^{(\mu)} \right\}. \quad (21)$$

This expression is exact. Clearly  $\langle \psi_k | \psi_k \rangle \neq 1$  in general (despite  $\langle \psi_k^{(0)} | \psi_k^{(0)} \rangle = 1$ ).

(3) Now suppose we are working in the  $\sigma = 1^{\text{st}}$  order of perturbation theory, as we did in Eqs. (17) on p. SS6. Eq. (21) prescribes (to  $\mathcal{O}(\lambda)$  and no higher)...

$$\rightarrow \langle \psi_k | \psi_k \rangle = 1 + 2\lambda \operatorname{Re} a_{kk}^{(1)} + \cancel{\mathcal{O}(\lambda^2)},$$

so reqt. that  $\langle \psi_k | \psi_k \rangle = 1 \Rightarrow \underline{\operatorname{Re} a_{kk}^{(1)} = 0}$ , or  $a_{kk}^{(1)} = i b_k^{(1)}$   $\int a_{kk}^{(1)}$  is pure imaginary

$$\text{only } \psi_k \approx \psi_k^{(0)} + \psi_k^{(1)} = [1 + \lambda a_{kk}^{(1)}] \psi_k^{(0)} + \underbrace{\left( \lambda \sum_n' a_{nk}^{(1)} \psi_n^{(0)} \right)}_{\psi_k^{(1)} = \frac{\lambda V_{nk}}{E_k^{(0)} - E_n^{(0)}}}$$

this is  $\mathcal{O}(\lambda^2) \rightarrow 0$ .

$$\text{or } \psi_k \approx [1 + i\lambda b_k^{(1)}] \psi_k^{(0)} \rightarrow [e^{i\lambda b_k^{(1)}}] \psi_k^{(0)}, \text{ to } \mathcal{O}(\lambda). \quad (22)$$

So the evanescent  $a_{kk}^{(1)}$  enters  $1^{\text{st}}$  order theory at most as a phase factor, which is arbitrary. We can set  $b_k^{(1)} \equiv 0$  w/o harm. So, as advertised,  $a_{kk}^{(1)} = 0$ .


(4) Now we can claim with confidence that in 1<sup>st</sup> order theory...

$$\left[ \begin{aligned} a_{mk}^{(1)} &= \begin{cases} V_{mk} / (E_k^{(0)} - E_m^{(0)}), & \text{for } m \neq k; \\ \underline{0}, & \text{for } m = k; \end{cases} \\ \text{So } \psi_k &\simeq \psi_k^{(0)} + \sum_n' \left( \frac{V_{nk}}{E_k^{(0)} - E_n^{(0)}} \right) \psi_n^{(0)} \end{aligned} \right] \quad \left. \begin{array}{l} \text{is 1<sup>st</sup> order perturbed wavefn} \\ \text{[prime on } \sum_n' \Rightarrow n \neq k \text{ in sum]} \end{array} \right\} \quad (23)$$

This maneuver removes the possibility that  $a_{kk}^{(1)} = \left( \frac{V_{kk}}{E_k^{(0)} - E_n^{(0)}} \right) \Big|_{n=k}$  diverges for  $V_{kk} \neq 0$ .

Also, it ensures that  $\langle \psi_k | \psi_k \rangle = 1 + \mathcal{O}(\lambda^2)$ , i.e. that  $\psi_k$  is normed to 1 to within the order of approx  $[\mathcal{O}(\lambda)]$ .

Note that in 1st order theory, the mixed state  $\psi_k$  has a contribution from  $\psi_n^{(0)}$  only if the "Coupling"  $V_{nk} \neq 0$ .



(5) It is clear from the Master Eq. (16) that in general the  $a_{kk}^{(\sigma)}$ ,  $\sigma \geq 1$ , are free constants (phase factors) because for  $m = k$  the eqn reads...

$$\rightarrow (E_k^{(0)} - E_k^{(0)}) a_{kk}^{(v+1)} + E_k^{(v+1)} = \sum_n' V_{kn} a_{nk}^{(v)} + \left[ V_{kk} a_{kk}^{(v)} - \sum_{\sigma=1}^v E_k^{(v+1-\sigma)} a_{kk}^{(\sigma)} \right]. \quad (24)$$

Since the first term LHS drops out, we get no restrictions on  $a_{kk}^{(v+1)}$  from any of the previous iterations for  $a_{nk}^{(\sigma)} \neq E_k^{(\sigma)}$ ,  $\sigma \leq v$ . We thus have the freedom:

Choose:  $a_{kk}^{(\sigma)} \equiv 0$ , for all  $\sigma \geq 1$

$$\text{So } \underline{1.} \quad \langle \psi_k | \psi_k \rangle = 1 + \sum_n' \left( \sum_{\sigma, \mu=1}^{\infty} a_{nk}^{(\sigma)*} a_{nk}^{(\mu)} \lambda^{\sigma+\mu} \right), \text{ from Eq. (21);}$$

$$\underline{2.} \quad \langle \psi_k^{(\sigma)} | \psi_k \rangle = \lambda^{\sigma} \sum_n' \left( \sum_{\mu=1}^{\infty} a_{nk}^{(\sigma)*} a_{nk}^{(\mu)} \lambda^{\mu} \right) \quad \left\{ \begin{array}{l} \text{the } ( ) = \mathcal{O}(\lambda^2) \text{ at least, so} \\ \psi_k^{(\sigma)} \text{ is } \perp \psi_k, \text{ negl. } \mathcal{O}(\lambda^{\sigma+2}). \end{array} \right.$$

$$\underline{3.} \quad E_k^{(v+1)} = \sum_n' V_{kn} a_{nk}^{(v)}, \text{ from Eq. (24).} \leftarrow \text{NICE SIMPLIFICATION!}$$

END of ASIDE