

An electron neutrino  $\nu_e$  created in subatomic interactions can later be observed as a muon neutrino  $\nu_\mu$ , and vice versa, in a process known as *neutrino oscillation*. Such oscillations occur **not** as decays, but as a result of a mismatch between the neutrino *flavor eigenstates* ( $\nu_e$  and  $\nu_\mu$ , with definite lepton numbers) and the *mass eigenstates*  $\nu_1$  and  $\nu_2$  (with definite masses  $m_1$  and  $m_2$ , respectively). The eigenstates in these complete, orthonormal bases<sup>1</sup> are coupled via an arbitrary real unitary matrix:

$$U = \begin{pmatrix} \langle \nu_e | \nu_1 \rangle & \langle \nu_e | \nu_2 \rangle \\ \langle \nu_\mu | \nu_1 \rangle & \langle \nu_\mu | \nu_2 \rangle \end{pmatrix} \equiv \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

where the unknown *mixing angle*  $\theta$  is real.

<sup>1</sup>We ignore here the tau neutrino  $\nu_\tau$  and any possible *sterile* neutrinos.

- (a) Briefly discuss any physical constraints that require  $U$  to be unitary.
- (b) Calculate the probability as a function of time  $t$  that a  $\nu_e$  at time  $t = 0$  with momentum  $p$  will transform into a  $\nu_\mu$ , expressing your answer in terms of  $\theta$  and the energies  $E_1 = \sqrt{p^2 c^2 + m_1^2 c^4}$  and  $E_2 = \sqrt{p^2 c^2 + m_2^2 c^4}$  of the two mass eigenstates.
- (c) Consider now the *ultrarelativistic limit*, i.e.,  $pc \gg mc^2$  for both mass eigenstates. Re-express your answer from part (a) in terms of  $p$ ,  $\theta$ ,  $\Delta m^2 \equiv m_2^2 - m_1^2$ , and distance traveled  $L \simeq ct$ .
- (d) Make an accurate plot of probability as a function of  $p$  for given values of  $L$ ,  $\theta$ , and  $\Delta m^2$ , and describe an experiment or set of experiments that could be used to determine both  $\theta$  and  $\Delta m^2$ .