Largest use of CO_2 (~50%) is not a chemical use but a refrigerant

CO₂(s) f.p. = -78.5°C
"dry ice"

CO_{2(s)}
$$\xrightarrow{>-78.5^{\circ}C}$$
 CO_{2(g)} sublimation directly from solid to gas!

How do you get liquid CO₂?

You have to keep a pressure on the solid when it is melting.

 CO_2 is a liquid $\rightarrow CO_{2(1)}$ at 5.2 atm and -5.6°C

- CO₂₍₁₎ is used to extract caffeine from coffee. It leaves no harmful residues.
- CO₂₍₁₎ is used in fire extinguishers. The more dense CO₂ will displace air around the burning material and keep O₂ from fueling the flames.
- ~25% of CO₂ produced is used in carbonation of beverages

"Chemistry of Soda"
$$CO_{2(aq)} + H_2O \Longrightarrow H_2CO_{3(aq)}$$

$$H_2CO_{3(aq)} + H_2O \Longrightarrow HCO_{3(aq)} + H_3O^+$$

TABLE 2.6 Some Polyatomic Ions

lon	Name (Alternate Name in Parentheses)
NH ₄ +	Ammonium ion
H ₃ O+	Hydronium ion ^a
OH-	Hydroxide ion
CN-	Cyanide ion
NO_2^-	Nitrite ion
NO ₃ -	Nitrate ion
CIO-	Hypochlorite ion
CIO ₂ -	Chlorite ion
CIO ₃ -	Chlorate ion
CIO ₄ -	Perchlorate ion
MnO_4^-	Permanganate ion
C ₂ H ₃ O ₂ -	Acetate ion
CO ₃ ²⁻	Carbonate ion
HCO ₃ -	Hydrogen carbonate ion (bicarbonate ion)b
SO ₃ 2-	Sulfite ion
SO ₄ 2-	Sulfate ion
HSO ₄ -	Hydrogen sulfate ion (bisulfate ion)
CrO ₄ 2-	Chromate ion
Cr ₂ O ₇ ²⁻	Dichromate ion
PO ₄ 3-	Phosphate ion (orthophosphate ion)
HPO ₄ 2-	Monohydrogen phosphate ion
H ₂ PO ₄	Dihydrogen phosphate ion

^a You will only encounter this ion in aqueous solutions.

^b Although "hydrogen carbonate ion" is formally correct, "bicarbonate ion" is what you will see and hear the most. We'll use "bicarbonate" too.

CO₂ Reaction with Base:

$$\begin{array}{l} CO_{2(aq)} + OH^{\text{-}}_{(aq)} \longrightarrow HCO_{3}^{\text{-}}_{(aq)} \\ HCO_{3}^{\text{-}}_{(aq)} + OH^{\text{-}}_{(aq)} \longrightarrow CO_{3}^{\text{-}}_{(aq)} + H_{2}O_{(1)} \\ \underline{Other\ carbon\ compounds} \end{array}$$

Cyanides CN⁻
 [:C≡N:]⁻¹ strong base
 NaCN sodium cyanide

HCN hydrogen cyanide (weak acid)

• Reaction of CN with water:

$$CN_{(aq)} + H_2O = HCN_{(aq)} + OH_{(aq)}$$
 equilibrium

• Reaction of CN with a strong acid:

strong base
$$CN^{-}_{(aq)} + HCl_{(aq)} \rightarrow HCN_{(aq)} + Cl^{-}_{(aq)}$$
 strong acid Reaction is complete!

Properties of Cyanides

- HCN is extremely poisonous
- it was used in gas chambers. CN binds to the heme in your mitochondria that transport O₂. Not the same heme as CO which is your blood heme.
- NaCN is also lethal
 Hemoglobin in blood is based on Fe
 Heme in mitrochondria is Cu based

Nitrogen

- 78% of air is N₂
- 25-30 million tons N₂/ year

Industrial Preparation of N₂:

$$Air \xrightarrow{liquify} N_2 \text{ (collected first)}$$

$$Ar$$

$$O_2$$

Uses of N₂:

- Enhanced oil recovery to force oil from subterranean deposits (~30% of N₂ made)
- Coolant (low b.p. 77k (-196°C)) freezing perishables (meat, seafood)
- Unreactive gas in chemical industry used as a blanketing atmosphere

Nitrogen Cycle helps to maintain balance of N_2 in the atmosphere.


- Plants remove N₂ to make <u>NH</u>₃
- Plants decay back to N₂

Nitrogen Fixation:

Name given to the reactions that microorganisms use to make NH_3 from N_2 .

Ammonia:

- Sharp odor
- Irritates lungs. Can cause death if inhaled in large quantities




• Used as a fertilizer by injecting directly into the soil

Haber – Bosch Process:

catalysts
$$N_2(g) + 3H_2(g) = 2NH_3(g)$$
 $\Delta H = -92.38KJ$ requires a catalyst and high pressure $AH = -92.38KJ$ is the heat of reaction $AH = -92.38KJ$ Negative $AH = -92.38KJ$ means exothermic.

The extreme conditions are required because of the large <u>kinetic barrier</u> to the reaction.



Properties of NH₃:

- b.p. -33.4°C
- f.p. -77.7°C
- very soluble in H_2O due to \underline{H} -bonding ability. It is a <u>weak base</u> in H_2O .

$$NH_{3(aq)} + H_2O \rightleftharpoons NH_{4(aq)}^+ + OH_{(aq)}^-$$

(an equilibrium exists in H_2O)

- Reacts completely with strong acids $NH_{3(aq)} + HCl_{(aq)} \rightarrow NH_4Cl_{(aq)}$
- Dissolves Group IA, IIA metals $Na + NH_{3(1)} \rightarrow Na^+ + NH_{3(1)} + e^-$ The e^- is "solvated" by $NH_3!!$

- NH₂ is a powerful <u>base</u> and is found in combination with metals such as Na⁺ and K⁻
- Amides react with H₂O to give strongly <u>basic</u> solutions.

$$NaNH_2 + H_2O \rightarrow NH_{3(aq)} + NaOH_{(aq)}$$
 (very exothermic reactions!)

Ammonium

 NH_4^+

- NH₄⁺ is slightly <u>acidic</u>
- Reacts with bases:

$$NH_4^+Cl_{(aq)}^- + NaOH_{(aq)} \longrightarrow NH_{3(aq)} + NaCl_{(aq)} + H_2O_{(1)}$$

Nitrides


 N^{3}

- N³- combined with metals
- (i.e. Mg₃N₂, Li₃N) are ionic
- N³⁻ combined with non-metals (i.e. P₃N₅, BN) are covalent

Nitric Acid

 HNO_3

The most important oxo acid of nitrogen



Resonance forms of Lewis structure

Ostwald Process to make Nitric Acid:

- Very important reaction
- Discovered by the German scientist Ostwald in 1902
- It's discovery is thought to have prolonged WWI because Germany had been cut off from importing nitrate salts from Chile by the Allies. Nitrates are used in explosives.

Oswald Process is 3 steps:

(1)
$$4NH_{3(g)} + 5O_{2(g)} \xrightarrow{Pt/Rh} 4NO_{(g)} + 6H_2O_{(g)}$$

 $900 \, ^{\circ}C$ $\Delta H = -1170 \, \text{KJ}$
 $\underline{\text{Very exothermic!}}$
(Ammonia to nitric oxide)

- (2) $2NO_{(g)} + O_{2(g)} \rightarrow 2NO_{2(g)}$ This is down-stream from the NH₃+ O₂ reaction (1). (nitric oxide to nitrogen dioxide)
- (3) $3NO_{(g)} + H_2O \rightarrow 2HNO_{3(aq)} + NO_{(g)}$ (nitrogen dioxide to nitric acid and nitric oxide)

Overall:

$$NH_{3(g)} + 2O_{2(g)} \rightarrow HNO_{3(aq)} + H_2O$$
Uses of Nitric Acid HNO_3

- 1) To make NH_4NO_3 for <u>fertilizers</u> $NH_3 + HNO_3 \rightarrow NH_4NO_3$
- 2) To make NH₄NO₃ for <u>explosives</u>

Alfred Nobel, a Swedish chemist, discovered how to safely handle <u>nitroglycerine</u> with the NH₄NO₃ and amassed a fortune, some of which he used to fund the Nobel Prizes.

NH₄NO₃ is unstable.

Decomposition of NH₄NO₃:

Under mild conditions: $NH_4NO_3(s) \xrightarrow{200-260 \text{ °C}} N_2O(s) + 2H_2O(g)$

With strong heating:

2NH₄NO₃(s)
$$\rightarrow$$
 300 °C 2N₂(g) + O₂(g) + 4H₂O(g) (2 moles solid \rightarrow 2 + 1 + 4 = 7 moles of gas) Rapid expansion occurs!

The violent explosion of a ship being loaded with fertilizer (NH₄NO₃) in Texas City, Texas took the lives of ~600 persons in 1947.

Other Nitrogen Explosives

$$\begin{array}{c} \underline{\text{Nitroglycerine}} \\ C_3H_5N_3O_9 \\ (\text{liquid}) \\ \end{array} \begin{array}{c} H_2C \longrightarrow \text{ONO}_2 \\ \\ HC \longrightarrow \text{ONO}_2 \\ \\ H_2C \longrightarrow \text{ONO}_2 \end{array}$$

$$4C_3H_5N_3O_9 \rightarrow 6N_2 + 12CO + 10H_2O + 7O_2$$

Nitroglycerine
(4 molecules of liquid \rightarrow 35 moles of gas!)

The expanding gases cause a violent detonation but no smoke!

<u>Dynamite</u> – this is a mixture of Nitroglycerine/NH₄NO₃/wood pulp/CaCO₃

(this is a filter used to neutralize any acids that may form during storage)

<u>TNT</u> – trinitrotoluene (solid)

Oxygen

50% of all atoms on earth are oxygen. 61% of all atoms in earth's crust are O. (crust is 16-40 km thick)

 O_2

- Diatomic
- Colorless, odorless, tasteless

Allotropes:

(different molecular forms of the same element)

- \bullet O₂
- O₃

Isotopes: Three for O:

16O 17O 18O

99.8% 0.2%

Dry air is $\sim 21\%$ of $O_2 \rightarrow$ this has not changed for millions of years due to the <u>oxygen cycle</u> in nature which maintains the balance.

Oxygen $\underline{consumed} \equiv oxygen \underline{produced}$

- respiration

- photosynthesis by cholorophyll-

- decay

containing organisms

- combustion of fuels

Photosynthesis

$$nCO_2 + nH_2O \xrightarrow{\text{sunlight}} (CH_2O)_n + nO_2$$
 carbohydrates (especially glucose)

Note: >50% of all O₂ from photosynthesis comes from photoplankton in oceans

- The cycle continues when decay, respiration and combustion take O₂ back to CO₂ and H₂O.
- Q What would happen if the oxygen cycle did not maintain O_2 concentration in air at ~21%?

<u>A</u>

Everything would burn out of control - forest fires, house fires etc.,

Why?

Because the rates of reactions increase with higher concentrations of reagents.

<u>Industrial Production of O2:</u>

Air
$$\stackrel{\text{liquify}}{\longrightarrow}$$
 Liquid Air $\stackrel{}{\longrightarrow}$ N₂ (1)
Ar (2)
O₂ (3)

 N_2 , Ar boil off first; wait to collect pure O_2

Laboratory (small scale) Synthesis of O₂:

2 KClO_{3(s)}
$$\xrightarrow{\text{MnO}_2 \text{ (cat)}}$$
 3O_{2(g)} + 2KCl_(s)

Uses of O₂:


(top 5th chemical in the U.S. ~ 19 million tons)

- 1) steel industry blast furnaces
- 2) chemical industry plastics
- 3) sewage treatment aerobic bacteria
- 4) health industry respirators
- 5) rocket industry fuel

Ozone

 O_3

Pungent odor Lewis structure/VSEPR



• Importance of O₃ for life:

It absorbs ho in the UV range which screens us from this harmful radiation

$$O_3 + UV \text{ light} \rightarrow O_2 + O$$

- Oxidizing Ability of O₃
 Very strong oxidant in basic and acidic media.
 Second only to fluorine in its oxidizing ability
- Ozone is a dangerous pollutant in smog. It attacks trees, fabrics, rubber, plastics, & <u>lungs!</u>
- at 0.000005% O₃ in air (0.5 <u>parts per million</u>) young children and elderly people are at risk
- at 0.000001% O₃ (1 ppm) O₃ is dangerous to everyone

Oxides

All elements except Noble gases form oxides

Three catagories: (THIS IS ALL REVIEW)

- Basic ionic oxides (form with metals)
- Acidic covalent oxides (form with non-metals, metalloids, some metals)
- Amphoteric can be ionic or covalent (form with metals)

Basic Oxides

- Form OH in H₂O
- Groups I, IA (except Be), In, Tl, some transition metals

Examples:

$$NaO_{(s)} + H_2O \rightarrow 2NaOH_{(aq)}$$

$$O^{2-} \qquad \qquad O^{3-} \qquad \qquad O \rightarrow O$$

 $MgO_{(s)} + H_2O \rightarrow Mg(OH)_{2(s)}$ (insoluble hydroxide)

Acidic Oxides

- Form acids in water
- All non-metals except noble gases. SO₃, SO₂, NO, NO₂, SiO₂, Sb₂O₃, etc., and some transition elements

Examples:

$$SO_3 + H_2O \rightarrow H_2SO_{4(aq)}$$

 $CrO_3 + H_2O \rightarrow H_2CrO_{4(aq)}$

Amphoteric Oxides

- Can be either acidic or basic
- Al, Ga, Sn, Pb and most transition metals
- They can neutralize acid or base

Example:
$$Al_2O_3$$
 (amphoteric)
(C) $Al_2O_{3(s)} + 2OH_{(aq)}^- + 7H_2O \rightarrow 2[Al(H_2O)_2(OH)_4]_{(aq)}^-$

(B)
$$Al_2O_{3(s)} + 6H^+_{(aq)} + 9H_2O \rightarrow 2[Al(H_2O)_6]^{3+}_{(aq)}$$

In reaction (A), Al₂O₃ is an acid In reaction (B), Al₂O₃ is a base How do you predict if a transition metal oxide will be acidic, basic or amphoteric?

Two trends

Trend 1

The <u>higher</u> the <u>oxidation state</u> of the metal, the more <u>covalent</u> (acidic) it will be.

Trend 2

The <u>lower</u> the <u>oxidation state</u> of the metal, the more <u>ionic</u> (basic) it will be.

Consider: Cr⁺²O, Cr₂⁺³O₃, Cr⁺⁶O₃ Most ionic is CrO (lowest ox. state) Most covalent is CrO₃ (highest ox. state)

∴ CrO would be basic CrO₃ would be acidic Cr₂O₃ would be amphoteric

Practice these: Mn₂O₃, MnO, [MnO₄]⁻ OsO₄, OsO₂, OsO

Peroxides $| (O_2^{2-})$ ox. state is O^{-1}

- Hydrogen peroxide: H₂O₂
- colorless liquid
- strong oxidizing agent
- used as a bleach, disinfectant $H_2O_2 \xrightarrow{\Delta} H_2O + \frac{1}{2}O_2$ Exothermic!
- Alkali Metal Peroxides: M₂O₂

 Na_2O_2 K_2O_2

							7			6			5			4			w			2		_	-		
						(223)	Fr	87	132.9	Cs	55	85.47	Rb	37	39,10	×	19	22.99	Na	=	6.941	L	w	1.008	Ħ	:-	-
						(226)	Ra	88	137.3	Ва	56	87.62	Sr	38	40.08	Ca	20	24.31	Mg	12	9.012	Be	4	2			
						(227)	Ac	89	138.9	La	57	16.88	X	39	44.96	Sc	21	1									
						(261)	Rf	2	178.5	Hf	4	91.22	Zr	6	47.88]	22		文	Source de la	,						
232.0	Th	8	140.1	Ce	58	(262)	Hn	105	180.9	Ta	t	92.91	Zb.	41	50.94	<	23		SA CO	2	>				アガス	1	
231.0	Pa	91	140.9	Pr	59	(263)	Unh	100	183.9	¥	74	95.94	Mo	42	52.00	Cr	24		202		ごう				IODI		
238.0	U	93	144.2	Nd	8	(262)	Zs	107	186.2	Re	75	(98)	Te	43	54.94	Mn	25	1		7:	7				CIA	3	
237.0	Np	93	(145)	Pm	61	(265)	Hs	108	190.2	Os	76	101.1	Ru	4	55.85	Fe	26		CLAD	AND SCIENT FORM	X				PERIODIC TABLE OF ELEMENTS		
(244)	Pu	94	150.4	Sm	62	(266)	M	109	192.2	Ir	77	102.9	Rh	45	58,93	Co	27	THE PERSON NAMED IN	TION		oxida				OH EI	1	
(243)	Am	25	152.0	Eu	63				195.1	P	78	106.4	Pd	46	58.69	Z	28		0	2	1				LEMI	7 2 2	
(247)	Cm	8	157.3	Gd	2				197.0	Au	E	107.9	Ag	47	63.55	Cu	29	No maconomic		8					NIN		
(247)	Bk	97	. 158.9	47	65				200.6	Hg	80	112.4	Cd	48	65.39	Zn	30		L	P					5		1
(249)	Cf	98	162.5	Ď	8				204.4	11	00	114.8	In	49	69.72	Ga	31	26.98_	N	: 53	18:04	B	5	3	(2	171
(254)	Es	99	164.9	Ho	67				207.2	Pb	82	118.7	Sn	50	72.59	Ge	32	23	101	Ta la	12.01	9	10	4	1	12	-
(253)	Fm	100	167.3	Er	68				209.0	Bi	83	121.8	Sb	51	74.99	As	38	30.97	7	12	74.01	Z	7	5		5	2
(256)	Md	101	168.9	Tm	69				(209)	Po	22	127.6	Per	52	78.96	Se	¥	27.06	C.	16	16.00	0	00	6		13	5:5
(254)	Zo	25	173.0	4.4	5				1010	マネア	85	NZ6.9)	53	79.90	Br	25	35.45	The state of the s	17	1	1	9	7		1	0
(256)	Lr	103	175.0	Lu	71				(222)	1 Rn	86		Xe	72	83.80	T	36	39.95	Ar	18	20.18	Ne	10	4.003	He	2	00

						02_2	7	l ,		6			5			A			w			2			-		
						(223)	Fr	87	132.9	Cs	55	85.47	Rb	37	39.10	X	19	22,99	Na	Π	6.941	Li	w	1.008	H	-	
						(226)	Ra	88	137.3	Ва	56	87.62	Sr	38	40.08	Ca	20	24.31	Mg	12	9.012	Ве	4	2			
						(227)	Ac	89	138.9	Z.	57	88.91	×	39	44.96	Sc	21	1	C	6	3	į.				5	I
						(261)	Rf	2	178.5	Hf	7	91.22	Zr	46	47.88	T	22		CINGRAD	-	子ロアく					1	MUNC
232.0	Th	8	140.1	Ce	58	(262)	Hn	105	180.9	Ta	7	92.91	Zb	41	50.94	<	23	The same of the sa	CAT						PER	5	20
231.0	Pa	91	140.9	Pr	59	(263)	Unh	8	183.9	¥	74	95.94	Mo	42	52.00	Cr	24		092	*	のなうる	1			IODI		さ
238.0	U	22	144.2	Nd	8	(262)	Zs	107	186.2	Re	75	(98)	Te	43	54.94	Mn	25				9				CTA		7
237.0	Z Z	93	(145)	Pm	19	(265)	Hs	108	190.2	Os	76	1.101	Ru	4	55.85	Fe	26	Decree of the last	Jun	3	77	7			PERIODIC TABLE OF ELEMENTS		C
(244)	Pu	20	150.4	Sm	62	(266)	Mt	109	192.2	Ir	77	102.9	Rh	45	58.93	Co	27	COLUMN TANKS	3	さっちつい	101 01	5			OF EI	1	2
(243)	Am	95	152.0	Eu	63				195.1	Pt	78	106.4	Pd	46	58.69	Z	28	St. Common	,	2	P. 1 . 1 . 1 . 1	3			EMI	Andrew Park	77
(247)	Cm	8	157.3	Gd	2				197.0	Au	78	107.9	Ag	47	63.55	Cu	29	10000		1 0	1	3			STN		V
(247)	Bk	97	158.9	T _b	83				200.6	Hg	88	112.4	Cd	48	65.39	Zn	3	4									
(249)	Cf	98	162.5	Dy	8	», <u> </u>			204.4	11	81	114.8	In	49	Per-097	ゆ		26.98	100	3	18.01	Ħ	S	u			
(254)	Es	99	164.9	Ho	67				ATT.	- 11	Rank Contract of	TOP OF	183	1		_		2.4						10			
(253)	Fm	100	167,3	Er	68				209.0	Bi	83	121.8	Sb	51	74.92	As	33	30.97	P	15	14,01	Z	7	Ci			
(256)	Md	101	168.9	Tm	69				(209)	Po	20	127.6	Te	52	78.96	Se	¥	32.06	S	16	16.00	0	8	6			
(254)	No	102	173.0	4,4	75			100000000000000000000000000000000000000	(210)	At	85	126.9)==(53	79.90	Br	35	35.45	C	17	19.00	H	9	7			
(256)	Lr	103	175.0	Lu	71			THE STREET	(222)	Rn	88	131.3	Xe	2	83.80	不	8	39.95	Ar	100	20.18	Ne	10	4.003	He	١.	∞

								7			6			S			A			w	0.007.00		2	_		-		
						- Contractor - Con	(223)	石	180	132.9	S.S.	55	25/17	RB	K	39,10	4	99	22.99	Za	=	6.01	4	W.	1.008	H-	-	_
						(Chinese Land	(226)	Ra	000	27.3	Ba	156	87.62	ST	108	40.08	Ca	20	124.31	No.	12	9.012	Be	4	2	/.		- Contract
							(227)	Ac	89	138.9	2	57	88.91	×	39	44.96	Sc	21	1		0.570					3		2
							(261)	Rf	104	178.5	Hf	7	91.22	Zr	8	47.88	T:	22		会	STA)				>	~	1
0 000	Th	8	140.1	Ce	58		(262)	Hn	105	180.9	Ta	E	92.91	Z _b	41	50.94	<	23	00	N N	_	-223				AEG AEG	1	1
2110	Pa	91	140.9	Pr	59		(263)	Unh	106	183.9	×	74	95.94	Mo	42	52.00	Cr	24	1	Amundanout	SIC NO					TODI) (
0 856	U	92	144.2	Zd	8		(262)	Zs	107	186.2	Re	75	(98)	Tc	43	54.94	Mn	25	1	tan						PERIODIC TABLE OF ELEMENTS	1	X
2770	Zp	93	(145)	Pm	61		(265)	Hs	108	190.2	Os	76	101.1	Ru	44	55.85	Fe	26	-	Thursi tion -	- 5	りんごとう				BLE (-
(344)	Pu	28	150.4	Sm	62	,	(266)	M	9	192.2	Ir	T	102.9	Rh	45	58.93	Co	27				>				OF EI	(Charleson of the Charleson of the Charl	2
CAC	Am	25	152.0	Eu	63					195.1	R	78	106.4	Pd	\$	58.69	Z	28	ACTION DESCRIPTION	ALL Y		9				EME		
(242)	Cm	8	157.3	Gd	2					197.0	Au	b b	107.9	Ag	47	63.55	Cu	29	The second second	F						STN		
(747)	Bk	97	158.9	4	83					200.6	Hg	8	112.4	Cd	48	65.39	Zn	30										
(249)	Cf	98	162.5	Dy	86					204	A	8/	14.8	5	6	69.72	Ga	31	26.98	A	13	10.81	В	5	u			
354	Es	99	164.9	Ho	67				1	207.2	Pb	82	118.7	Sn	50	72.59	Ge	32	28.09	S:	14	12.01	0	6	4			
(253)	Fm	100	167.3	Er	68					209.0	Bi	83	121.8	Sb	51	74.92	As	33	30.97	ď	15	14.01	Z	7	5			
(256)	pW	101	168.9	Tm	69					(209)	Po	200	127.6	Te	52	78.96	Se	2	32.06	S	16	16.00	0	8	6			
(254)	No	102	173.0	4Y	70					(210)	J.	85	126.9	1	53	79.90	Br	35	35.45	0	. 17	19.00	דו	9	7			
(256)	Lr	103	175.0	Lu	71					(222)	Rn	86	131.3	Xe	X	83.80	7	8	39.95	Ar	180	20.18	Ne	10	4.003	He	2	00