

성욕저하시

testosterone inj. tx

가. 테스토스테론주사 반앰플씩 사용하고 있어요
나. 테스토스테론 수치는 개별차가 심해서, 부정확합니다. Free testosterone(CX232) 추가 검사해보세요
다. 예나스테론은 저는 3개월마다 삼분의1씩 주는데, 그 전 다른 원장님은 아마 한달마다 반앰플씩 주신다고 했던것
도같네요. 저도 예나스테론 1/3 A씩 한달에 한두번 씹니다 더 쓰면 턱수염도 날 수 있어서 소량씩 줍니다
라. 대표적 부작용들 이 이렇습니다.[5 35% 발현]

- 1) HDL 감소
- 2) Acne
- 3) Hirsutism
- 4) Clitorimegaly
- 5) Voice deepening(산공방 어느 선생님)

TESTOSTERONE THERAPY FOR TX OF DISORDERS OF DESIRE

- Screening
 - Baseline testosterone levels(free and total), baseline lipid profile, baseline liver enzyme level, mammography, PAP smear
- Initiate Therapy
 - Combination product(Estratest or Estratest hs)
 - Methyltestosterone(Android), 1.25 to 2.5mg daily
 - Micronized oral testosterone, 5mg twice daily
 - Testosterone proprionate 2 percent in petroleum applied daily to every other day
 - Testosterone injectables/pellets
- Reevaluation at three to four months
 - Repeat testosterone levels, lipid profile, liver enzyme levels
 - Monitor symptoms, side effects
- Continued therapy
 - Taper to lowest effective dosage
 - Monitor lipid levels, liver enzyme levels once or twice yearly
 - Routine PAP smear and mammography schedules.