

Zaire

No code

Investigating the Applegate effect for the post-common-envelope binary V471 Tau

Semester : 2020B

Science Cat. : Stars and stellar population

Abstract

This proposal will tackle the key question about whether an Applegate mechanism can be operating in the V471 Tau system or not. This effect is theoretically predicted to operate in several binary systems, but yet no definitive proof of its action have been found. We will address this question by using spectropolarimetric time-series to reconstruct the magnetic field of the K dwarf companion and to obtain the surface differential rotation of the star. With these results, we will be able to discuss if an Applegate effect can be the responsible mechanism for the eclipse timing variations observed in the system.

Telescopes

Telescope	Observing mode	Instruments
CFHT	QSO Regular	ESPaDOnS

Applicants

Name	Affiliation	Email	Country	Potential observer
Bonnie Zaire	Institut de Recherche en Astrophysique et Planétologie (IRAP)	bonniezaire@gmail.com	France	Pi
Jean-Francois Donati	CNRS / Univ de Toulouse (IRAP / OMP)	jean-francois.donati@irap.omp.eu	France	
Baptiste Klein	IRAP	baptiste.klein@irap.omp.eu	France	

Contact Author

Title

Name

Bonnie Zaire

Email

bonniezaire@gmail.com

Phone(first)

+330769838544

Phone(second)

Fax

Institute

Institut de Recherche en Astrophysique et Planétologie (IRAP)

Department

Address

14 Avenue Edouard Belin

Zipcode

31400

City

Toulouse

State

Country

France

Requested Time (exposure + overheads) in hours (in nights for Classical)

15.6

Overall scheduling requirements

2020/11/29 to 2020/12/04

We request a total of 15.6h, spanning in 5 consecutive days. The observations will take place in three different nights with 5.2h of observation each. An interval of one day is asked from one observation to the other. This setup requires observations from mid-October to late December to achieve the requested hours with an airmass <1.5.

Observing runs

Run	Instrument	Seeing	Config	Details
A	ESPaDOnS	< 1.00" (ESPaDOnS)	Observing mode: Polarimetry Read-out mode : Normal	Stokes V polarimetry We request a total of 15.6h, spanning in 5 consecutive days. The observations will take place in three different nights with 5.2h of observation each. An interval of one day is asked from one observation to the other. This setup requires observations from mid-October to late December to achieve the requested hours with an airmass <1.5.

Targets

Field	RA	Dec	Epoch		Runs	Moon	Seeing Lower	Seeing Upper	S/N	Magnitude	Diameter	Comments
V* V471 Tau	03:50:24.97	+17:14:47.4	J2000		A	full	0.1	1	110	9.373		wavelength of the SN: 596 nm

Science justification

Overall context: V471 Tau is an eclipsing binary system whose components are a K dwarf main-sequence star and a hot white dwarf star (Nelson B., Young A., 1970, PASP, 82, 699). In the last 50 years, V471 Tau has been exhaustively observed in an attempt to understand the evolution of binary systems. The current scenario for V471 Tau indicates that the system is a pre-cataclysmic variable that, in early stages, has undergone a common-envelope phase (Paczynski B., 1976, IAU Symposium Vol. 73, p. 75).

As in most pos-common-envelope binary systems, eclipse timing variations are observed when considering a linear ephemeris to predict the time of mid-eclipse (Young A., Lanning H. H., 1975, PASP, 87, 461). The existence of a third body perturbing the orbit of the system was disclaimed with an image of the binary system obtained with the instrument SPHERE (Hardy A., et al., 2015, ApJ, 800, L24). A plausible explanation that still stands is the existence of a variable magnetic field that modulates the orbital period of the system. This mechanism, known as **Applegate effect**, drives period variations as a result of magnetically induced changes in the quadrupole momentum of the K dwarf component (Applegate J. H., 1992, ApJ, 385, 621). Its feasibility would require a cyclic behaviour in the magnetic field and a significant variation in the differential rotation.




Figure 1: Polar view of magnetic field topology for 2004 (top panels) and 2005 (bottom panels). Columns depict different components of the magnetic field, namely radial, azimuthal and meridional field components. Magnetic field strengths are saturated at 600G, with red shades representing positive values and blue shades negative values.

Recent analysis of data collected with ESPaDoNS in 2004 and 2005 unveiled a magnetic variability at the K dwarf surface (Zaire B.R., Donati J.F., Klein B., 2020, in preparation). Fig. 1 shows the reconstructions of the surface field of the K dwarf star for both years. The results indicate that a dynamo mechanism operate within the convective zone of the K dwarf, which is a necessary ingredient for action of the Applegate mechanism. Moreover, this first analysis shows a significant fluctuation in the differential rotation that

suggest inverse correlation with the magnetic field strength. Although this results favor the action of an Applegate effect, there is no observational proof that the magnetic field has the cyclic behaviour required to justify the eclipse timing variations.

Specific objective of the proposal: We propose to apply the Zeeman-Doppler imaging technique to spectropolarimetric data of the K dwarf star to reconstruct its magnetic topology. This work will be a follow up of two observational campaigns performed in 2004 and 2005. The 15 years gap between the observation proposed in this work and the last observation performed for the system corresponds to half of the cycle obtained with eclipse timing variations. It represents an opportune time to constrain the magnetic variability and to get hints about the existence of a **cyclic magnetic field**. Furthermore, we can measure the **differential rotation** at the star surface and extend previous investigations about the angular momentum distribution inside the K dwarf companion. Finally, we aim to discuss whether an Applegate mechanism operating in the V471 Tau system could explain the series of observations (2004, 2005, and, now, 2020).

Technical justification

Observation strategy: The observational setup proposed here follows the one adopted in previous observational campaigns for the same system (performed in 2004*B* and 2005*B*). We aim to use the ESPaDoNS instrument in the circular polarization mode to observe the V471 Tau binary system ($\alpha = 03 : 50 : 24.97$, $\gamma = +17 : 14 : 47.43$). We ask for observations every other day spanning in 5 consecutive days. Given the short rotational period of ~ 12.5 h, this scheme of observation maximizes the coverage of the data with respect to the rotational phase of the K dwarf companion. We highlight that the phase coverage is a key factor to achieve our scientific goals. Another possible setup is to adopt a two days interval between each observation, however this would represent an upper limit because placing the observed nights farther apart may compromise the necessary coherence between the data set.

Requested signal-to-noise ratio: We request a **SNR of 110 in each individual subexposure**, which results in a SNR of ~ 260 over each polarimetric sequence. Guided by previous data acquisitions, we can then expect the SNR in the LSD profiles to reach ~ 4000 after summing up all spectral lines of V471 Tau.

Requested time: Using the ESPaDoNS Exposure Time Calculator, we find that at a wavelength of 596 nm the exposure time needed to reach a SNR of 110 is 180 s for an object of magnitude 9.373 and T_{eff} 5066 K, under a seeing of 0.8" and airmass of 1.5. Therefore we suggest full observation sequences totaling 800 s each, including an overhead time for each subexposure of 20 s. We aim at obtaining 66 Stokes V polarimetric sequence (made of 4 individual subexposures) in this campaign. With the estimation described above, this can be achieved with **5.2 h of observation in three different nights**. We thus request a total of **15.6 h = 2 nights** of ESPaDoNS time. Using the ‘Object Observability’ tool of ESO (<http://www.eso.org/sci/bin/skycalcw/observability>), one sees that with a airmass ≤ 1.5 the requested amount of night hours can be achieved in 2020B **from mid October to late December**.

No PhD Students involved

Linked proposal submitted to this TAC: No

Linked proposal submitted to other TACs: No

Any other expenditure

Relevant previous Allocations: Yes

No additional remarks

Observing run info :

Run: A backup strategy: Compute etime, order:38, wavelength:596, snr:110, saturation time: 8939, magv:9.373, teff:5066, airmass:1.5, seeing:0.8, omode:polarimetric, rmode:normal, etime:180