

THE UNIVERSITY OF NEW SOUTH WALES

DEPARTMENT OF STATISTICS

Additional exercises for MATH5905, Statistical Inference

Part four: Multinomial distribution. Order statistics. Robustness

1. a) X_1, X_2, X_3 has a multinomial $(8; 0.2, 0.3, 0.5)$ distribution. Find $P(X_1 = 2, X_2 = 2, X_3 = 4)$, $E(X_2)$, $Var(X_2)$ and $Cov(X_1, X_3)$.
b) X_1, X_2, X_3 has a multinomial $(6; 0.5, 0.2, 0.3)$ distribution. Find $P(X_1 = 3, X_2 = 1, X_3 = 2)$ and $P(X_1 + X_2 = 2)$.

2. Find the probability density function of the second order statistic $X_{(2)}$ in a random sample of size four from a population with the density function

$$f(x) = \begin{cases} e^{1-x}, & 1 < x < \infty, \\ 0 & \text{elsewhere} \end{cases}$$

3. Find the probability density function of $X_{(4)}$ in a random sample of size five from a population with the density function

$$f(x) = \begin{cases} \frac{1}{x^2}, & 1 \leq x < \infty, \\ 0 & \text{elsewhere} \end{cases}$$

4. The opening prices per share of two similar stocks Y_1 and Y_2 are independent random variables, each with density function

$$f(y) = \begin{cases} \frac{1}{2}e^{-\frac{y-4}{2}}, & y \geq 4, \\ 0 & \text{elsewhere} \end{cases}$$

On a given morning Mr. A is going to buy shares of whichever stock is less expensive. Find the probability density function and the expected value for the price per share that Mr. A will have to pay.

5. Find the expected value of the largest order statistic in a random sample of size 3 from:
a) the exponential distribution with density $f(x) = e^{-x}, x \geq 0$.
b) the standard normal distribution

6. Electric components of a certain type have lifetime Y with probability density given by

$$f(y) = \begin{cases} \frac{1}{100}e^{-\frac{y}{100}}, & y > 0, \\ 0 & \text{elsewhere} \end{cases}$$

- a) Suppose that two such components operate independently and in series in a certain system (that is, the system fails when either component fails). Find the density function for X , the lifetime of the system.
b) Now suppose that the components operate in parallel (that is, the system does not fail until both components fail). Find the density function for X , the lifetime of the system.
7. A continuous random variable X has a standard exponential distribution

$$f(x) = \begin{cases} e^{-x}, & x > 0, \\ 0 & \text{elsewhere} \end{cases}$$

For a random sample of size 3, let $X_{(1)}, X_{(2)}, X_{(3)}$ denote the ordered sample.

- a) Write down the joint distribution of $X_{(1)}$ and $X_{(3)}$.
 - b) Obtain the distributions of $X_{(1)}$ and $X_{(3)}$.
 - c) Evaluate $EX_{(1)}$ and $EX_{(3)}$.
 - d) Find the sampling distribution of the range $R = X_{(3)} - X_{(1)}$.
8. A random sample of size 3 is taken from a population with density

$$f(x) = \begin{cases} 2x, & 0 \leq x < 1, \\ 0 & \text{elsewhere} \end{cases}$$

Find the sampling distribution of the range R .

9. For a random sample of size 2 from a standard normal distribution, find the distribution of the range.
10. The Cauchy-Schwartz Inequality tells us that for random variables X, Y with $E(X^2) < \infty$ and $E(Y^2) < \infty$, $\{E(|XY|)\}^2 \leq E(X^2)E(Y^2)$ holds. Using this Inequality, show that when estimating the location parameter θ in $f(x - \theta)$ by using an M-estimator (defined by its ψ function):
- a) we get a loss in asymptotic efficiency in comparison to the MLE estimator (i.e. the Inequality

$$\sigma^2(F, \psi) = \frac{\int \psi^2(x)f(x)dx}{(\int \psi'(x)f(x)dx)^2} \geq \frac{1}{\int [\frac{f'(x)}{f(x)}]^2 f(x)dx}$$

holds).

- b) Equality holds only when the M-estimator coincides with the MLE. (**But: despite the above observation we still may decide to use the M-estimator instead of the MLE because of the better robustness properties of the former in comparison to the latter**).
11. Assume that we are estimating the location parameter θ_0 for the family $f_\theta(x) = f(x - \theta)$ in a robustness context. The “true” value θ_0 to be estimated in robustness context is usually taken as the solution of the Equation $E_{\theta_0}\psi(X - \theta_0) = 0$. Derive the asymptotic normality statement from the lecture notes for the M-estimator a defined as the solution to the equation $\sum_{i=1}^n \psi(X_i - \hat{\theta}_M) = 0$:

$$\sqrt{n}(\hat{\theta}_M - \theta_0) \rightarrow^d N(0, \frac{\int \psi^2(x)f(x)dx}{(\int \psi'(x)f(x)dx)^2}).$$

Answers

- 1 a) 0.0945, 2.4, 1.68, -0.8 b) 0.135 0.059535
- 2) $12\exp(3(1 - x_{(2)}))(1 - \exp(1 - x_{(2)})), 1 \leq x_{(2)} < \infty$
- 3) $20(x_{(4)} - 1)^3/x_{(4)}^6, 1 \leq x_{(4)} < \infty$
- 4) $\exp(-y_{(1)} - 4), y_{(1)} \geq 4$. The expected value is 5.
- 5) a) 11/6, b) $\frac{3}{2\sqrt{\pi}}$.
- 6) a) $\frac{1}{50}\exp(-\frac{x}{50})$; b) $\frac{1}{50}(1 - \exp(-\frac{x}{100}))\exp(-\frac{x}{100})$
- 7) a) $6(e^{-x_{(1)}} - e^{-x_{(3)}})e^{-(x_{(1)}+x_{(3)})}, 0 < x_{(1)} < x_{(3)} < \infty$.
b) $3\exp(-3x_{(1)}), x_{(1)} > 0; 3(1 - \exp(-x_{(3)}))^2\exp(-x_{(3)}), x_{(3)} > 0$.
- c) 1/3, 11/6; d) $2\exp(-u)(1 - \exp(-u)), u > 0$.
- 8) $12u(1 - u)^2, 0 < u < 1$.
- 9) $\frac{1}{\sqrt{\pi}}\exp(-\frac{u^2}{4}), u > 0$.