

Query Complexity of Mastermind Variants

Aaron Berger, Christopher Chute, Matthew Stone

August 7, 2015

Mastermind

- i. Codemaker vs. Codebreaker
- ii. Queries: Guess a vector from $\{1, 2, \dots, 6\}^4$.
- iii. Response
 - i. Black hits
 - ii. White hits



Knuth Paper – 1976

At most five turns needed to guarantee a victory

Minimax

- For each possible guess

 - For each possible response to that guess

 - Check how many possible solutions remain

 - Let *score* be min. number solutions eliminated

- Make guess with maximum score

Optimality of the Minimax Algorithm

Lemma (Pigeonhole Principle)

For a query with r possible responses, there exists a response that would leave at least $1/r$ of possible solutions remaining.

Analyze the worst-case performance of any algorithm.

- i. Worst case response to first guess \Rightarrow At least 256 solutions remain.
- ii. Second guess: At least $\lceil 256/14 \rceil = 19$ solutions remain.
- iii. Third guess: At least $\lceil 19/14 \rceil = 2$ solutions remain
- iv. Fourth guess: At least two possible solutions left \Rightarrow cannot guarantee to guess the solution on the fourth turn.

Extensions

- i. Basic Extension: n spots, k colors
- ii. Repeats vs. no repeats
- iii. Non-adaptive vs. adaptive strategies

Trivial Lower Bound

Theorem

(Game: n spots, k colors, repeats, adaptive allowed). For any strategy that guarantees a win in s turns,

$$s \geq O\left(\frac{n \log k}{\log n^2}\right)$$

Proof.

Trivial Lower Bound

Theorem

(*Game: n spots, k colors, repeats, adaptive allowed*). For any strategy that guarantees a win in s turns,

$$s \geq O\left(\frac{n \log k}{\log n^2}\right)$$

Proof.

1. Winning Condition: Distinct sequence of responses for each of k^n vectors. Let s be number of guesses to guarantee a win.

Trivial Lower Bound

Theorem

(Game: n spots, k colors, repeats, adaptive allowed). For any strategy that guarantees a win in s turns,

$$s \geq O\left(\frac{n \log k}{\log n^2}\right)$$

Proof.

1. Winning Condition: Distinct sequence of responses for each of k^n vectors. Let s be number of guesses to guarantee a win.
2. $\binom{n+2}{2}$ possible responses (cf. stars and bars), giving $\binom{n+2}{2}^s$ possible response sequences.

Trivial Lower Bound

Theorem

(Game: n spots, k colors, repeats, adaptive allowed). For any strategy that guarantees a win in s turns,

$$s \geq O\left(\frac{n \log k}{\log n^2}\right)$$

Proof.

1. Winning Condition: Distinct sequence of responses for each of k^n vectors. Let s be number of guesses to guarantee a win.
2. $\binom{n+2}{2}$ possible responses (cf. stars and bars), giving $\binom{n+2}{2}^s$ possible response sequences.
3. Therefore, s must satisfy

$$\binom{n+2}{2}^s \geq k^n$$

(Somewhat) Trivial Upper Bound

1. Represent guesses and solutions as matrices ($Q_{ij} = 1$ iff the i -th spot is the j -th color)

Example

The guess $(1, 1, 3)$ would become:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

(Somewhat) Trivial Upper Bound

1. Represent guesses and solutions as matrices ($Q_{ij} = 1$ iff the i -th spot is the j -th color)

Example

The guess $(1, 1, 3)$ would become:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

2. $Q_{ij} \in \mathbb{R}^{nk}$

(Somewhat) Trivial Upper Bound

1. Represent guesses and solutions as matrices ($Q_{ij} = 1$ iff the i -th spot is the j -th color)

Example

The guess $(1, 1, 3)$ would become:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

2. $Q_{ij} \in \mathbb{R}^{nk}$
3. # of black hits of Q with hidden matrix X is $Q \cdot X$.

(Somewhat) Trivial Upper Bound

1. Represent guesses and solutions as matrices ($Q_{ij} = 1$ iff the i -th spot is the j -th color)

Example

The guess $(1, 1, 3)$ would become:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

2. $Q_{ij} \in \mathbb{R}^{nk}$
3. # of black hits of Q with hidden matrix X is $Q \cdot X$.
4. Guess a basis of span of all such matrices ($\leq nk$ guesses)

(Somewhat) Trivial Upper Bound

1. Represent guesses and solutions as matrices ($Q_{ij} = 1$ iff the i -th spot is the j -th color)

Example

The guess $(1, 1, 3)$ would become:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

2. $Q_{ij} \in \mathbb{R}^{nk}$
3. # of black hits of Q with hidden matrix X is $Q \cdot X$.
4. Guess a basis of span of all such matrices ($\leq nk$ guesses)
5. Dot products with basis \Rightarrow Uniquely determine X

(Somewhat) Trivial Upper Bound

1. Represent guesses and solutions as matrices ($Q_{ij} = 1$ iff the i -th spot is the j -th color)

Example

The guess $(1, 1, 3)$ would become:

$$\begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Theorem

There exists a set of at most nk guesses such that any hidden vector is uniquely determined by the responses to those guesses.

Entropy Method

Surprise Function: For an event x , we want

Definition (Surprise Function)

Entropy Method

Surprise Function: For an event x , we want

Definition (Surprise Function)

1. $S(x) = 0$ when $\mathbb{P}[x] = 1$

Entropy Method

Surprise Function: For an event x , we want

Definition (Surprise Function)

1. $S(x) = 0$ when $\mathbb{P}[x] = 1$
2. $S(x) = 1$ when $\mathbb{P}[x] = 1/2$

Entropy Method

Surprise Function: For an event x , we want

Definition (Surprise Function)

1. $S(x) = 0$ when $\mathbb{P}[x] = 1$
2. $S(x) = 1$ when $\mathbb{P}[x] = 1/2$
3. Decreasing function of $\mathbb{P}[x]$

Entropy Method

Surprise Function: For an event x , we want

Definition (Surprise Function)

1. $S(x) = 0$ when $\mathbb{P}[x] = 1$
2. $S(x) = 1$ when $\mathbb{P}[x] = 1/2$
3. Decreasing function of $\mathbb{P}[x]$
4. $S(x \wedge y) = S(x) + S(y|x)$ ($= S(x) + S(y)$ if independent)

Entropy Method

Surprise Function: For an event x , we want

Definition (Surprise Function)

1. $S(x) = 0$ when $\mathbb{P}[x] = 1$
2. $S(x) = 1$ when $\mathbb{P}[x] = 1/2$
3. Decreasing function of $\mathbb{P}[x]$
4. $S(x \wedge y) = S(x) + S(y|x)$ ($= S(x) + S(y)$ if independent)

Unique Surprise Function

$$S(x) = -\log_2(\mathbb{P}[x]).$$

Entropy Method

Entropy is the expected surprise of a random variable.

Entropy Method

Entropy is the expected surprise of a random variable.

Definition (Entropy)

Let X be a random variable with domain D .

$$H(X) = \sum_{x \in D} \mathbb{P}[X = x] \cdot (-\log_2 (\mathbb{P}[X = x]))$$

Entropy Method

Lemma

A single turn in a non-adaptive strategy has < 4 bits of entropy.

Entropy Method

Lemma

A single turn in a non-adaptive strategy has < 4 bits of entropy.

Lemma

The number of bits of entropy in any winning strategy is exactly

$$\log_2 \left(\frac{k!}{(k-n)!} \right).$$

Entropy Method

Lemma

A single turn in a non-adaptive strategy has < 4 bits of entropy.

Lemma

The number of bits of entropy in any winning strategy is exactly

$$\log_2 \left(\frac{k!}{(k-n)!} \right).$$

Theorem

The number of queries submitted by any winning non-adaptive strategy is at least

$$\frac{1}{4} \log_2 \left(\frac{k!}{(k-n)!} \right).$$

Probabilistic Method

Choose a random set of queries $Q = \{q_1, q_2, \dots, q_s\}$.

Calculate $\mathbb{P}[Q \text{ is a winning set of guesses}]$

Probabilistic Method

Choose a random set of queries $Q = \{q_1, q_2, \dots, q_s\}$.

Calculate $\mathbb{P}[Q \text{ is a winning set of guesses}]$

Theorem

When the random set of guesses is of magnitude $s = 4k \log k$

$$\mathbb{P}[Q \text{ is a winning set of guesses}] > 0$$

Probabilistic Method

Choose a random set of queries $Q = \{q_1, q_2, \dots, q_s\}$.

Calculate $\mathbb{P}[Q \text{ is a winning set of guesses}]$

Theorem

When the random set of guesses is of magnitude $s = 4k \log k$

$$\mathbb{P}[Q \text{ is a winning set of guesses}] > 0$$



\exists a winning set of $4k \log k$ guesses

Thanks!

1. Special thanks to Danny Montealegre, Nathan Kaplan.
2. Thanks to SUMRY for this research opportunity.
3. Thanks to MathFest for the opportunity to present.