

Homework-03

Callie Baker

2026-05-28

Table of contents

Set up	2
Problem 1. Giant kelp fronds	2
a. An appropriate test	2
b. Create a visualization	2
c. Check your assumptions and run your test	3
Assumption checks	3
Running the test	5
d. Results communication	6
e. Test implications	6
f. Double check your work	6
Problem 2. Personal data	7
a. Updating your visualizations	7
b. Captions	10
Problem 3. Affective visualization	10
a. Describe your affective visualization	10
b. Create a paper sketch of your idea	10
c. Make a draft of your visualization	11
d. Write an artist statement	12
e. Prep your materials to share in class	13
Problem 4. Statistical critique	13
a. Revist and summarize	13
b. Visual clarity	14
c. Aesthetic clarity	14
d. Recommendations	14

Set up

```
# read in packages
library(tidyverse) # general use
library(here) # file organization
library(janitor) # cleaning data frames

# read in data
kelp <- read_csv(here("data", "temp-kelp.csv"))
my_data <- read_csv(here("data", "envs193ds-personal-data.csv"))
```

Problem 1. Giant kelp fronds


a. An appropriate test

To determine the strength of the relationship between temperature and giant kelp frond elongation rate, I can test for the correlation between the two continuous variables using Pearson's correlation (r) or Spearman rank correlation (ρ). Pearson's correlation is used when variables are continuous and normally distributed; Pearson's r represents the linear relationship between two variables and is used for parametric studies. In contrast, Spearman rank correlation is the non parametric version which describes monotonic relationship between variables.

b. Create a visualization

```
# scatterplot for temperature vs kelp growth rate
# base layer: kelp data
ggplot(data = kelp,
       # x-axis: temperature
       # y-axis: elongation rate
       mapping = aes(x = temp_c,
                     y = kelp_elong)) +
# first layer: points to represent individual kelp observations
# changed point size, stroke, shape, and color from defaults
geom_point(size = 2,
           stroke = 1,
           shape = 21,
           fill = "darkgreen") +
# relabeled x- and y-axes
```

```
# added title
labs(x = "Ocean Temperature (°C)",
     y = "Kelp Elongation Rate (cm/day)",
     title = "Kelp Grows Faster at Cooler Ocean Temperatures") +
# changed theme from default
theme_minimal()
```



Note: There appears to be a negatively linear relationship between kelp growth and ocean temperature. In other words, kelp growth rate decreases at increased ocean water temperatures.

c. Check your assumptions and run your test


Assumption checks

Check 1: The data includes independent observations, and both variables are continuous.

Check 2: Linearity was assessed in Question 1b. There appears to be a negative linear relationship between kelp growth and ocean temperature seen through the created scatterplot.


Check 3: Normality was assessed using two QQ plots, one for each variable.

```
# base layer: ggplot
# data frame: kelp
ggplot(data = kelp,
       # y-axis: temperature
       aes(sample = temp_c)) +
# first layer: reference line colored red
geom_qq_line(color = "red") +
# second layer: QQ plot
geom_qq()
```



Note: The data for ocean temperature appears to be normally distributed.

```
# base layer: ggplot
# data frame: kelp
ggplot(data = kelp,
       # y-axis: kelp elongation
       aes(sample = kelp_elong)) +
# first layer: reference line colored red
geom_qq_line(color = "red") +
# second layer: QQ plot
geom_qq()
```



Note: The data for kelp elongation appears to be normally distributed.

Summary of assumption checks: The data includes independent observations, and both variables are continuous. Linearity was assessed by looking at the scatterplot created in Problem 1b and was met because data points appear to follow a straight line. Normality was assessed using two Q-Q plots; both variables appear normally distributed as the data for each variable aligns closely with their red reference line. Finally, no significant outliers were detected after visual assessment from the scatterplot. Therefore, all assumptions for Pearson's correlation were met.

Running the test

```
# ran Pearson's test for correlation
# variables: ocean temperature and kelp elongation rate
cor.test(kelp$temp_c, kelp$kelp_elong,
         method = "pearson") # Pearson's r
```

Pearson's product-moment correlation

```
data: kelp$temp_c and kelp$kelp_elong
t = -5.189, df = 30, p-value = 1.366e-05
```

```
alternative hypothesis: true correlation is not equal to 0
95 percent confidence interval:
 -0.8359665 -0.4460157
sample estimates:
      cor
-0.6877489
```

d. Results communication

To evaluate the strength of the relationship between temperature and giant kelp frond elongation rate, I used a Pearson's correlation test. I found a strong negative linear relationship between ocean temperature (°C) and giant kelp elongation rate (cm/day) (Pearson's $r = -0.69$, $t(30) = -5.19$, $p < 0.001$, $\alpha = 0.05$). Therefore, the null hypothesis should be rejected because there is a strong relationship between ocean water temperature and giant kelp frond elongation rate.

e. Test implications

The results outlined above indicate that giant kelp frond growth rate decreases with increased ocean water temperatures. Therefore, warming ocean temperatures from climate change could threaten giant kelp productivity and species that rely on healthy giant kelp ecosystems.

f. Double check your work

```
# ran Spearman rank test for correlation
# variables: ocean temperature and kelp elongation rate
cor.test(kelp$temp_c, kelp$kelp_elong,
         method = "spearman", # Spearman rank
         exact = FALSE)
```

Spearman's rank correlation rho

```
data: kelp$temp_c and kelp$kelp_elong
S = 9216.1, p-value = 1.29e-05
alternative hypothesis: true rho is not equal to 0
sample estimates:
      rho
-0.6891684
```

Both tests led to the same decision which is to reject the null hypothesis because there is a strong negative correlation between ocean temperature and giant kelp frond growth rate. Using the Spearman rank test, I found a strong negative monotonic relationship between ocean temperature (°C) and giant kelp elongation rate (cm/day) (Spearman (ρ) = -0.69, $S = 9216.1$, $p < 0.001$, $r = 0.05$). Both tests produced correlation coefficients of -0.69 and p-values of less than 0.001, suggesting that warmer temperatures tend to produce lower kelp growth rates.

Problem 2. Personal data

a. Updating your visualizations

Figure 1:

```
# boxplot with jitter to visualize workout duration vs workout type

# cleaned column names
my_data <- my_data |>
  clean_names()

# base layer: ggplot with x- and y-axes
# data frame: my_data
ggplot(data = my_data,
       # x-axis: workout type (categorical predictor variable)
       # y-axis: workout duration (response variable)
       mapping = aes(x = workout_type,
                     y = duration_min,
                     # colored points by workout type
                     color = workout_type)) +
# first layer: boxplot to show distribution
  geom_boxplot() +
# second layer: jitter to show underlying data
  geom_jitter(height = 0, # no jitter in vertical direction
             width = 0.2, # narrowed jitter in horizontal direction
             alpha = 0.5, # made points more transparent
             shape = 1) + # changed points to be open circles
# changed colors from default
  scale_color_manual(values = c("Lift" = "darkgreen",
                                "Run" = "navyblue",
                                "Surf" = "darkorange",
                                "Yoga" = "darkred")) +
# relabeled x- and y-axes
```

```
# added title summarizing takeaway
# added subtitle with recent date
labs(title = "Runs show the most variable workout duration",
      x = "Workout type",
      y = "Workout duration (minutes)",
      subtitle = "Most recent observation: 2026-05-26") +
# changed ggplot theme from default
theme_minimal() +
# removed legend
theme(legend.position = "none")
```




Figure 2:

```
# boxplot with jitter: workout type vs sleep the previous night (hours)


# base layer: ggplot with x- and y- axes
# data frame: my_data
ggplot(data = my_data,
       mapping = aes(x = workout_type,
                     y = sleep_hours,
                     color = workout_type)) +
# first layer: boxplot to show distribution
geom_boxplot() +
```

```

# second layer: jitter to show underlying data
geom_jitter(height = 0, # no jitter in vertical direction
            width = 0.2, # narrowed jitter in horizontal direction
            alpha = 0.5, # made points more transparent
            shape = 1) + # changed points to be open circles
# changed colors from default
scale_color_manual(values = c("Lift" = "darkgreen",
                              "Run" = "navyblue",
                              "Yoga" = "darkred",
                              "Surf" = "darkorange")) +

# relabeled x- and y-axes
# added title summarizing main message
# added subtitle with recent date
labs(title = "More sleep is associated with going on runs",
     x = "Workout type",
     y = "Sleep the previous night (hours)",
     subtitle = "Most recent observation: 2026-05-26") +
# changed ggplot theme from default
theme_minimal() +
# removed legend
theme(legend.position = "none")

```



b. Captions

Figure 1. Runs show the most variable workout duration. Points represent individual workout sessions ($n = 30$, most recent observation: 2026-05-26). Colors represent workout type (green: lift, blue: run, orange: surf, red: yoga). Boxes show the median and interquartile range of workout duration (minutes) for each workout type (lift, run, surf, yoga). Note that surf sessions ($n = 2$) are limited in sample size.

Figure 2. More sleep is associated with going on runs. Points represent individual workout sessions ($n = 30$, most recent observation: 2026-05-26). Colors represent workout type (green: lift, blue: run, orange: surf, red: yoga). Boxes show the median and interquartile range of sleep the previous night (hours) for each workout type (lift, run, surf, yoga). Note that surf sessions ($n = 2$) are limited in sample size.


Problem 3. Affective visualization

a. Describe your affective visualization

In analyzing my personal data, I found that run duration is more variable than lifting, surfing, and yoga, and that I tend to sleep more on nights before runs than before other workout types. With these observations in mind, I plan to create an affective visualization exploring the relationship between sleep duration and workout type. My visualization will be a hand drawn landscape where hills of varying heights represent hours of sleep the previous night. Each section of the landscape will be colored and labeled by workout type, and stick figures doing that workout will be placed on top.


b. Create a paper sketch of your idea

Note: This sketch was hand drawn on my iPad (I don't have paper...)



c. Make a draft of your visualization

Note: This draft was also hand drawn on my iPad. I am still deciding whether I want to draw my visualization out on paper for my final copy.



d. Write an artist statement

Content: This piece visualizes the relationship between sleep duration and workout type from my personal workout data. Each colored landscape section represents a workout type (lift, run, surf, yoga), with the height of the rolling hills representing hours of sleep the previous night; taller peaks indicate more sleep. Stick figures placed on the hills are in the position of their workout type and are situated at the height corresponding to their sleep value.

Influences: Jill Peltó's paintings, which overlay data directly into natural landscapes, were a primary influence on this piece. The layered contour lines were generally inspired by topographic maps, which I hope add depth and a sense of flow to my visualization.

Form: This piece is hand drawn using Notability on my iPad. I am still determining if my final visualization will be created online with my iPad or on paper with watercolors. If I decide to do watercolors on paper, I will use Jill Peltó's paintings as reference because I love her style.

Process: The first step was sketching the horizontal landscape divided into sections by workout type. Then, I adjusted the hill heights to more accurately reflect the sleep distributions I observed in my data (see Figure 2 in Problem 2a). Stick figures were added on top of each hill to represent workout sessions; although I still have to decide if I want the number of figures to accurately reflect the exact number of observations. Contour lines were drawn between the hills to reinforce the differences in sleep duration (hill height).

e. Prep your materials to share in class

[View](#) my slides.

Problem 4. Statistical critique

a. Revist and summarize

The statistical tests the authors use are three-factor ANOVAs with equal replication to test the effects of temperature, food quantity, and food quality on abalone growth (shell length and body mass), reproductive output (gonad bulk index), disease burden (WS-RLP score), and mortality. Each response variable is tested separately, and the ANOVA also tests all two-way and three-way interactions among the three predictors. Additionally, the authors apply multivariate analyses (cluster analysis and multidimensional scaling based on Euclidean distances) to examine how the 27 treatment combinations group together across all response variables simultaneously.

TABLE 1. *F* values and probabilities from three-factor ANOVAs with equal replication, testing the effects of varying temperature, food quantity, and food quality on abalone growth, reproductive output, disease, and mortality.

Source of variation	df	Growth							Reproduction (GBI)		
		Length			Mass						
		MS	F	P	MS	F	P	MS	F	P	
<i>Haliotis rufescens</i>											
Temperature	2	0.05	181.87	0.00	1.80	276.15	0.00	0.22	3.87	0.03	
Quantity	2	0.08	272.42	0.00	1.40	215.86	0.00	1.86	32.51	0.00	
Quality	2	0.02	79.22	0.00	0.51	79.08	0.00	0.50	8.74	0.00	
Temperature : Quantity	4	0.00	4.16	0.01	0.02	3.71	0.01	0.05	0.90	0.47	
Temperature : Quality	4	0.00	4.76	0.00	0.03	5.30	0.00	0.03	0.55	0.70	
Quantity : Quality	4	0.00	10.22	0.00	0.06	9.55	0.00	0.29	5.14	0.00	
Temperature : Quantity : Quality	8	0.00	0.34	0.94	0.01	1.06	0.41	0.03	0.59	0.78	
Residuals	54	0.00			0.01			0.06			
<i>H. fulgens</i>											
Temperature	2	0.00	11.34	0.00	0.01	2.41	0.10	0.84	8.05	0.00	
Quantity	2	0.12	690.91	0.00	1.57	474.76	0.00	7.32	70.11	0.00	
Quality	2	0.02	133.68	0.00	0.29	88.96	0.00	0.36	3.45	0.04	
Temperature : Quantity	4	0.00	9.10	0.00	0.05	14.36	0.00	0.52	4.99	0.00	
Temperature : Quality	4	0.00	1.46	0.23	0.00	0.85	0.50	0.13	1.24	0.30	
Quantity : Quality	4	0.00	22.46	0.00	0.03	10.10	0.00	0.05	0.47	0.76	
Temperature : Quantity : Quality	8	0.00	0.82	0.59	0.00	1.08	0.39	0.08	0.78	0.62	
Residuals	54	0.00			0.00			0.10			

Notes: Growth was evaluated through standardized changes in animal mass and shell length [(initial – final value)/initial value]. Reproductive output was evaluated from gonad bulk indices (GBI). Disease was evaluated by the burden of withering syndrome, Rickettsiales-like prokaryote (WS-RLP) and scores of morphological changes associated with withering syndrome (digestive gland atrophy [DGA] and foot muscle degeneration [FM-DG]), and accumulated mortalities. All variables were derived from cell means since that was the experimental unit and not individual abalone (see *Methods* for details). A posteriori tests were run on those treatments with significant effects (see Appendix).

TABLE 1. Extended.

Disease and morbidity											
WS-RLP			DGA			FM-Deg			Mortality		
MS	F	P	MS	F	P	MS	F	P	MS	F	P
4.74	19.64	0.00	2.64	25.32	0.00	1.80	9.48	0.00	0.38	41.11	0.00
0.34	1.40	0.25	1.78	17.07	0.00	2.08	10.90	0.00	0.04	3.97	0.02
0.07	0.28	0.76	0.40	3.84	0.03	0.98	5.13	0.01	0.00	0.08	0.92
0.19	0.79	0.54	0.32	3.05	0.02	0.15	0.81	0.53	0.03	3.49	0.01
0.10	0.41	0.80	0.15	1.43	0.24	0.05	0.25	0.91	0.00	0.20	0.94
0.31	1.29	0.28	0.02	0.23	0.92	0.24	1.24	0.30	0.00	0.16	0.96
0.35	1.45	0.20	0.12	1.14	0.35	0.08	0.43	0.90	0.00	0.34	0.94
0.24			0.10			0.19			0.01		
0.02	0.86	0.43	0.00	1.00	0.37	0.08	1.19	0.31	0.00	2.00	0.15
0.00	0.21	0.81	0.00	1.00	0.37	0.05	0.81	0.45	0.00	0.50	0.61
0.02	0.86	0.43	0.00	1.00	0.37	0.26	3.94	0.03	0.00	0.50	0.61
0.02	1.07	0.38	0.00	1.00	0.42	0.03	0.50	0.74	0.00	0.50	0.74
0.03	1.71	0.16	0.00	1.00	0.42	0.07	1.09	0.37	0.00	0.50	0.74
0.02	1.07	0.38	0.00	1.00	0.42	0.04	0.63	0.65	0.00	1.25	0.30
0.01	0.64	0.74	0.00	1.00	0.45	0.05	0.69	0.70	0.00	1.25	0.29
0.02			0.00			0.07			0.00		

b. Visual clarity

Table 1 presents the ANOVA results organized by species and divided into response variable columns for growth, reproduction, disease, and mortality. Although Table 1 allows for clear comparison of how each predictor affects each outcome across species, it includes so much information that it was extended into two tables in the article. Furthermore, the table only reports MS, F, and p-values with no indication of the direction or magnitude of effects, so viewers cannot assess whether warm temperature increased or decreased growth.

c. Aesthetic clarity

Table 1 has a high data to ink ratio because the authors present dense statistical information in a compact grid without colors or borders. Nonetheless, I find the table difficult to follow due to the large amount of information it presents. Furthermore, significant p-values are not visually distinguished with asterisks or text modifications, making it harder for viewers to see important results.

d. Recommendations

Recommended modifications: To improve Table 1, I recommend adding asterisks for p-values less than 0.05 (the significance level) to improve readability of important takeaways. I

would also recommend adding an effect size column such as eta-squared to indicate the strength of the relationship between variables.

Additional figure: I recommend a bar chart displaying mean growth (specifying either mass or length) on the y-axis and temperature treatment (cool, ambient, warm) on the x-axis; bars would be grouped and colored by species (example: red for *H. rufescens*, green for *H. fulgens*). This figure would allow for direct species comparison across temperature levels and would convey the direction and magnitude of the effect of temperature on abalone growth.
