

Homework-03

Callie Baker

2026-05-28

Table of contents

Set up	2
Problem 1. Giant kelp fronds	2
a. An appropriate test	2
b. Create a visualization	2
c. Check your assumptions and run your test	3
Assumption checks	3
Running the test	5
d. Results communication	6
e. Test implications	6
f. Double check your work	6
Problem 2. Personal data	7
a. Updating your visualizations	7
b. Captions	9
Problem 3. Affective visualization	10
a. Describe your affective visualization	10
b. Create a paper sketch of your idea	10
c. Make a draft of your visualization	10
d. Write an artist statement	10
e. Prep your materials to share in class	10
Problem 4. Statistical critique	10
a. Revist and summarize	10
b. Visual clarity	10
c. Aesthetic clarity	10
d. Recommendations	10

Set up

```
# read in packages
library(tidyverse) # general use
library(here) # file organization
library(janitor) # cleaning data frames
library(readxl) # reading excel files
library(scales) # modifying axis labels
library(ggeffects) # getting model predictions
library(ggplot2)
library(MuMin) # model selection
library(dplyr)

# reading in data
kelp <- read_csv(here("data", "temp-kelp.csv"))
my_data <- read_csv(here("data", "envs193ds-personal-data.csv"))
```

Problem 1. Giant kelp fronds


a. An appropriate test

To determine the strength of the relationship between temperature and giant kelp frond elongation rate, I can test for the correlation between the two continuous variables using Pearson's correlation (r) or Spearman rank (ρ). Pearson's correlation is used when variables are continuous and normally distributed; Pearson's r represents the linear relationship between two variables and is used for parametric studies. In contrast, Spearman rank correlation is the non parametric version which describes monotonic relationship between variables.

b. Create a visualization

```
# scatterplot for temperature vs kelp growth rate
# base layer: kelp data
ggplot(data = kelp,
       # x-axis: temperature
       # y-axis: elongation rate
       mapping = aes(x = temp_c,
                     y = kelp_elong)) +
# first layer: points to represent individual kelp observations
```

```
# changed point size, stroke, shape, and color from defaults
geom_point(size = 2,
           stroke = 1,
           shape = 21,
           fill = "darkgreen") +
# relabeled x- and y-axes
# added title
labs(x = "Ocean Temperature (°C)",
     y = "Kelp Elongation Rate (cm/day)",
     title = "Kelp Grows Faster at Cooler Ocean Temperatures") +
# changed theme from default
theme_minimal()
```



Note: There appears to be a negatively linear relationship between kelp growth and ocean temperature. In other words, kelp growth rate decreases at increased ocean water temperatures.

c. Check your assumptions and run your test


Assumption checks

Check 1: The data includes independent observations, and both variables are continuous.

Check 2: Linearity was assessed in Question 1b. There appears to be a negative linear relationship between kelp growth and ocean temperature seen through the created scatterplot.

Check 3: Normality was assessed using two QQ plots, one for each variable.


```
# base layer: ggplot
# data frame: kelp
ggplot(data = kelp,
       # y-axis: temperature
       aes(sample = temp_c)) +
# first layer: reference line colored red
geom_qq_line(color = "red") +
# second layer: QQ plot
geom_qq()
```



Note: The data for ocean temperature appears to be normally distributed.

```
# base layer: ggplot
# data frame: kelp
ggplot(data = kelp,
       # y-axis: kelp elongation
       aes(sample = kelp_elong)) +
# first layer: reference line colored red
```

```
geom_qq_line(color = "red") +  
# second layer: QQ plot  
geom_qq()
```



Note: The data for kelp elongation appears to be normally distributed.

Summary of assumption checks: The data includes independent observations, and both variables are continuous. Linearity was assessed by looking at the scatterplot created in 1b and was met. Normality was assessed using Q-Q plots; both variables appear normally distributed as the data aligns closely with the red reference line. Finally, no significant outliers were detected from the scatterplot. Therefore, all assumptions for Pearson's correlation were met.

Running the test

```
# ran Pearson's test for correlation  
# variables: ocean temperature and kelp elongation rate  
cor.test(kelp$temp_c, kelp$kelp_elong,  
         method = "pearson") # Pearson's r
```

Pearson's product-moment correlation

```

data: kelp$temp_c and kelp$kelp_elong
t = -5.189, df = 30, p-value = 1.366e-05
alternative hypothesis: true correlation is not equal to 0
95 percent confidence interval:
 -0.8359665 -0.4460157
sample estimates:
      cor
-0.6877489

```

d. Results communication

To evaluate the strength of the relationship between temperature and giant kelp frond elongation rate, I used a Pearson's correlation test. I found a strong negative linear relationship between ocean temperature (°C) and giant kelp elongation rate (cm/day) (Pearson's $r = -0.69$, $t(30) = -5.19$, $p < 0.001$, $r^2 = 0.05$). Therefore, the null hypothesis should be rejected because there is a strong relationship between ocean water temperature and giant kelp frond elongation rate.

e. Test implications

The results outlined above indicate that giant kelp frond growth rate decreases with increased ocean water temperatures. Therefore, warming ocean temperatures from climate change could threaten giant kelp productivity and species that rely on healthy giant kelp ecosystems.

f. Double check your work

```

# ran Spearman rank test for correlation
# variables: ocean temperature and kelp elongation rate
cor.test(kelp$temp_c, kelp$kelp_elong,
         method = "spearman", # Spearman rank
         exact = FALSE)

```

Spearman's rank correlation rho

```

data: kelp$temp_c and kelp$kelp_elong
S = 9216.1, p-value = 1.29e-05
alternative hypothesis: true rho is not equal to 0
sample estimates:

```

```
rho
-0.6891684
```

Both tests led to the same decision which is to reject the null hypothesis because there is a strong negative correlation between ocean temperature and giant kelp frond growth rate. Using the Spearman rank test, I found a strong negative monotonic relationship between ocean temperature (°C) and giant kelp elongation rate (cm/day) (Spearman (ρ) = -0.69, $S = 9216.1$, $p < 0.001$, $\alpha = 0.05$). Both tests produced correlation coefficients of -0.69 and p-values of less than 0.001, suggesting that warmer temperatures tend to produce lower kelp growth rates.

Problem 2. Personal data

a. Updating your visualizations

Figure 1:

```
# created boxplot with jitter to visualize workout duration vs workout type

# cleaned column names
my_data <- my_data |>
  clean_names()

# base layer: ggplot with x- and y-axes
# data frame: my_data
ggplot(data = my_data,
       # x-axis: workout type (categorical predictor variable)
       # y-axis: workout duration (response variable)
       mapping = aes(x = workout_type,
                     y = duration_min,
                     # colored points by workout type
                     color = workout_type)) +
  # first layer: boxplot to show distribution
  geom_boxplot() +
  # second layer: jitter to show underlying data
  geom_jitter(height = 0, # no jitter in vertical direction
             width = 0.2, # narrowed jitter in horizontal direction
             alpha = 0.5, # made points more transparent
             shape = 1) + # changed points to be open circles
  # changed colors from default
  scale_color_manual(values = c("Lift" = "darkgreen",
                                "Run" = "navyblue",
```

```

"Yoga" = "darkred")) +
# relabeled x- and y-axes
# added title summarizing takeaway
labs(title = "Runs show the most variable workout duration",
      x = "Workout type",
      y = "Workout duration (minutes)",
      subtitle = "2026-05-26") +
# changed ggplot theme from default
theme_minimal() +
# removed legend
theme(legend.position = "none")

```




Figure 2:

```

# created scatter plot to visualize workout duration (minutes) vs
# sleep the previous night (hours)

# base layer: ggplot with x- and y-axes
# data frame: my_data
ggplot(data = my_data,
      # x-axis: sleep (hours) the night prior (continuous predictor
      # variable)


```



```

# y-axis: workout duration (response variable)
mapping = aes(x = sleep_hours,
              y = duration_min)) +
# first layer: points for each observation
geom_point(size = 2) + # increased point size
# relabeled x- and y-axes
# added title summarizing main message
labs(title = "More sleep may be associated with longer workouts",
      x = "Sleep the previous night (hours)",
      y = "Workout duration (minutes)") +
# changed ggplot theme from default
theme_minimal()

```



b. Captions

From the boxplot with overlaid jitter (problem 3a), I observed that my run workouts show the greatest variability in duration indicated by the wider spread of points and the taller box. Of the three groups, run workouts have the smallest median duration, while lift workouts have the largest median duration. Yoga workouts shows the most consistency with points clustered closely together at 60 minutes and no vertical box. In the continuous scatter plot showing workout duration vs sleep in hours (problem 3b), there does not appear to be a strong

relationship between variables, seen through the varied points. Collecting more data could shift the medians and spreads in the boxplots, and a larger sample size may reveal a clearer relationship between continuous variables in the scatter plot.

Problem 3. Affective visualization

- a. Describe your affective visualization**
- b. Create a paper sketch of your idea**
- c. Make a draft of your visualization**
- d. Write an artist statement**
- e. Prep your materials to share in class**

Problem 4. Statistical critique

- a. Revist and summarize**
 - b. Visual clarity**
 - c. Aesthetic clarity**
 - d. Recommendations**
-