

ENVS193DS Final

Callie Baker

Table of contents

GitHub repo	2
Set up	2
Problem 1. Research writing	2
a. Transparent statistical methods	2
b. Suggestions for rewriting	3
c. Further analysis	3
Problem 2. Data analysis	3
a. Clean and wrangle	3
b. Response variable	4
c. Purpose of study	4
d. Table of models	4
e. Run the models	4
f. Select the best model	4
g. Check the diagnostics	4
h. Visualize the model predictions	4
i. Write the caption for your figure	4
j. Calculate model predictions	4
k. Interpret your results	4
Problem 3. Affective and exploratory visualizations	4
a. Comparising visualizations	4
b. Designing an analysis	4
c. Check any assumptions and run your analysis	4
Setting up	4
Check assumptions	5
Run the analysis from part b	5
Calculate an effect size	7

d. Create a visualization	9
Visualizing model predictions	9
Describing the model	10
e. Write a caption	11
f. Write about your results	11
g. Sharing your affective visualization	11

GitHub repo

Access my repo [here](#).

Set up

```
# read in packages
library(tidyverse) # general use
library(here) # file organization
library(janitor) # cleaning data frames
library(DHARMA) # for diagnostics
library(ggeffects) # for ggpredict
library(flextable) # for flextable
library(gtsummary)

# read in data
nests <- read_csv(here("data", "nest_data_final.csv"))
# read in personal data to object from csv
my_data <- read_csv(here("data", "envs193ds-personal-data.csv"))
```

Problem 1. Research writing

a. Transparent statistical methods

Part 1: The statistical test used in Part 1 was a generalized linear model (logistic regression) with a binomial distribution. The response variable is a binary outcome to reflect the probability that the American Avocet uses the mircohabitat (1) or doesn't (0). The predictor is a continuous variable: distance to water edge (UNITS).

Part 2: The statistical test used in Part 2 was a chi-square test of independence which is used to determine whether there is a relationship between two categorical variables. The

categorical variables are bird species (American Avocet, Forster's Tern, Black-necked Stilt) and habitat type (managed pond, non-tidal marsh, salt pond, tidal marsh, tidal mudflat). The test compares observed counts to expected counts.

b. Suggestions for rewriting

Part 1: American Avocets were more likely to use microhabitats that were closer to the water's edge, and this relationship was strong enough that is unlikely to be the result of random chance alone.

Part 2: The three shorebird species did not use the five habitat types in equal proportions: each species showed a distinct pattern of habitat preference, and these differences across species are unlikely to reflect chance.

c. Further analysis

Tidal inundation frequency could influence American Avocet microhabitat use because Avocets forage in shallow flooded areas so microhabitats that are too rarely or too frequently inundated may not provided the water depth conditions needed for successful foraging. Tidal inundation frequency is a continuous variable, measured as the proportion of time a microhabitat is flooded per tidal cycle; this variable could be recorded using tidal monitoring stations.

Problem 2. Data analysis

*DO NOT include any code or output from your data exploration

a. Clean and wrangle

```
# cleaned data
nests_clean <- nests |>
  clean_names()
```

- b. Response variable
- c. Purpose of study
- d. Table of models
- e. Run the models
- f. Select the best model
- g. Check the diagnostics
- h. Visualize the model predictions
- i. Write the caption for your figure
- j. Calculate model predictions
- k. Interpret your results

Problem 3. Affective and exploratory visualizations

- a. Comparising visualizations
- b. Designing an analysis
- c. Check any assumptions and run your analysis

Setting up

```
# cleaned column names
my_data <- my_data |>
  clean_names()

# created new column: 1 if workout was a run, 0 if not
my_data <- my_data |>
  mutate(is_run = case_when(
    workout_type == "Run" ~ 1, # run = 1
    .default = 0))             # all other workout types = 0
```

Check assumptions

NEED TO DO indep obs, what else...


Run the analysis from part b

```
# fitted generalized linear model for running
# response: whether or not workout was a run (binary: 1/0)
# predictor: sleep the previous night (hours)
run_mod <- glm(
  # formula: binary run outcome ~ sleep hours
  is_run ~ sleep_hours,
  # data: my_data data frame
  data = my_data,
  # binomial error distribution for binary response
  family = binomial)
```

Checking diagnostics

```
# simulated residuals from GLM for diagnostic plots
plot(simulateResiduals(run_mod))
```

DHARMA residual



Info from left plot (QQ plot of residuals):

- KS test: do my residuals follow a uniform distribution? → generally yes (CHECK)
- Dispersion test: are your residuals more dispersed than expected from a uniform distribution? → no
- Outlier test: are there outliers in your data set that could be influencing model coefficients or predictions? → no

Info from right plot (residuals vs predicted values):

- By default, R generates the reference lines for quartiles: 25, 50, 75
- Not that informative but look at patterns within residuals across the range → DHARMA diagnostics flag issues in 25th quartile due to small sample size, not because the model is wrong.

Looking at the model coefficients

```
# displayed model summary
summary(run_mod)
```

Call:

```
glm(formula = is_run ~ sleep_hours, family = binomial, data = my_data)
```

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-8.7011	3.4964	-2.489	0.0128 *
sleep_hours	1.0873	0.4712	2.307	0.0210 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 36.652 on 29 degrees of freedom
Residual deviance: 28.046 on 28 degrees of freedom
AIC: 32.046

Number of Fisher Scoring iterations: 5

Calculate an effect size

Calculate inverse logit

```
# calculated inverse logit
plogis(-8.7011)
```

```
[1] 0.000166375
```

Note:

- inverse logit is 0.00016
- The probability of going on a run if sleep duration is 0 hours is 0.00016?

```
# calculated odds ratio
gtsummary::tbl_regression(run_mod,
                           exponentiate = TRUE)
```

Note:

Characteristic	OR	95% CI	p-value
sleep_hours	2.97	1.37, 9.40	0.021

Abbreviations: CI = Confidence Interval, OR = Odds Ratio

- results at top of this page
- odds ratio is 2.97
- $2.97 > 1$ so the probability of going on a run increases by a factor of 2.97 with every 1 hour increase in sleep

Summary: For every 1 hour increase in sleep, the the odds of going on a run increase by a factor of 2.97 (95% CI: [1.37, 9.40], $p = 0.021$, $\alpha = 0.05$).

**Looking at the model predictions (where to put this section?)

- using probabilities to determine chance of getting certain outcome
- line represents the probability of 1 (going on a run) actually occurring

```
# what is the probabiltiy of going on a run after sleeping for 10 hours?
ggpredict(run_mod,
  terms = "sleep_hours [10]")
```

Predicted probabilities of is_run

sleep_hours	Predicted	95% CI
10	0.90	0.39, 0.99

```
# what is the probabiltiy of going on a run after sleeping for 4 hours?
ggpredict(run_mod,
  terms = "sleep_hours [4]")
```

Predicted probabilities of is_run

sleep_hours	Predicted	95% CI
4	0.01	0.00, 0.25

Note:

- If I slept for 10 hours, the probability of going on a run the next day is 0.90 (95% CI [0.39, 0.99]).
- If I slept for 4 hours, the probability of going on a run the next day is 0.01 (95% CI [0.00, 0.25]).

d. Create a visualization

Visualizing model predictions


```
# generated model predictions across range of sleep values in data set
mod_preds <- ggpredict(run_mod,
                      terms = "sleep_hours [5:10 by = 1]")

# base layer: ggplot with raw data
# data frame: my_data
ggplot(my_data,
       aes(x = sleep_hours,
           y = is_run)) +
# first layer: raw observations
geom_point(size = 3,
           alpha = 0.4) +
# second layer: confidence interval ribbon from model predictions
geom_ribbon(data = mod_preds,
          aes(x = x,
              y = predicted,
              ymin = conf.low,
              ymax = conf.high),
          alpha = 0.4) +
# third layer: logistic curve from model predictions
geom_line(data = mod_preds,
         aes(x = x,
             y = predicted),
         # custom color
         color = "navyblue") +
# constrained y-axis to 0 and 1
scale_y_continuous(limits = c(0, 1),
                  breaks = c(0, 1)) +
# relabeled axes and added title
labs(title =
     "More sleep is associated with a higher probability of running the next day",
     x = "Sleep the previous night (hours)",
```

```

y      = "Probability of going on a run (1 = yes, 0 = no)" +
# changed theme from default
theme_minimal() +
# bolded title
theme(plot.title = element_text(face = "bold"))

```



Describing the model

```

# displaying model in table
as_flextable(run_mod)

```

	Estimate	Standard Error	z value	Pr(> z)
(Intercept)	-8.701	3.496	-2.489	0.0128 *
sleep_hours	1.087	0.471	2.307	0.0210 *

*Signif. codes: 0 <= '***' < 0.001 < '**' < 0.01 < '*' < 0.05*

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 36.65 on 29 degrees of freedom

Residual deviance: 28.05 on 28 degrees of freedom

e. Write a caption

Figure caption

f. Write about your results

ADD For every 1 hour increase in sleep, the the odds of going on a run increase by a factor of 2.97 (95% CI: [1.37, 9.40], $p = 0.021$, $\alpha = 0.05$).

g. Sharing your affective visualization

Presented visualization in Week 10 Workshop.