

## SPECIFIC AIMS

Genetically-encoded probes have revolutionized our understanding of cellular processes. Fluorescent proteins are a powerful technology that enable tracking of protein concentration and localization over large timescales and with single-molecule resolution. Via simple genetic manipulation, virtually any protein of interest can be labeled and tracked in the cell. However, analogous techniques for study of other biomolecules have yet to be developed.

The importance of RNA in controlling a wide range of cellular functions is only recently being realized.[1] Signaling by messenger and noncoding RNAs in the cell is highly dependent on the location of the transcripts. It is understood that cellular machinery modulates the location of these molecules, but these processes remain difficult to elucidate.[2] This lack of understanding is due in part to the lack of tools available to image this biomolecule. Localization of RNA on a single-molecule level, and multicomponent imaging of RNA transcripts remains difficult. Existing tools utilize aptamers that are unstable in mammalian cells[3], or constructs that are too large for imaging small transcripts. Multicomponent RNA imaging is also difficult due to the design of current tools.

To address this need, *I aim to develop a platform for RNA imaging that will enable facile tracking of multiple transcripts at high resolution.* Riboglow is an RNA imaging platform recently developed in the Palmer lab. It utilizes a fluorescence-quenched pair formed by cobalamin (vitamin B12) and a pendant fluorophore. In solution, this construct shows low fluorescence. When bound to the cobalamin riboswitch aptamer domain, there is an increase in fluorescence. This tool shows promise for RNA imaging because it solves many of the problems faced by traditional RNA probes. The platform is similar to Spinach[4], Broccoli[5], and Mango[6, 7] in that it utilizes a small molecule-binding aptamer for localization, but improves on these tools because it utilizes a natural aptamer that is more stable in the cellular environment. The current gold-standard for imaging is the MS2-FP system[8], which utilizes an aptamer-binding protein fused to a fluorescent protein. Though this technique benefits from the modularity of fluorescent proteins, it requires large constructs to concentrate the fluorescent signal, thus precluding its use with most small RNAs.

Riboglow is already a useful tool for RNA imaging, but several drawbacks are keeping it from widespread utility. The proposed work addresses these drawbacks, and seeks to utilize improved Riboglow tools to study outstanding questions in the field of noncoding RNA. The *objective* of the work is to image small, noncoding RNAs as they function within living cells. My *central hypothesis* is that the Riboglow platform can be improved through chemical modification and RNA selection to provide brighter, multicomponent probes.

### **Aim 1. Synthesize improved Riboglow probes.**

Previously developed Riboglow constructs suffered from poor signal turn-on and low brightness. I aim to derivatize the native cobalamin structure to produce new probe scaffolds. These new molecules will be evaluated for quenching efficiency and will be tested with native cobalamin aptamers.

### **Aim 2. Adapt Riboglow for superresolution imaging.**

The molecules developed in *Aim 1* will be screened against libraries of aptamers to further improve probe properties. Libraries of RNA aptamers will be generated and screening will be carried out in mammalian cells via flow cytometry. Candidate probes will be verified through superresolution imaging of stress granules in living cells.

### **Aim 3. Develop mutually orthogonal Riboglow probes for multicomponent imaging.**

In tandem with brightness optimization, I will develop mutually orthogonal probes to enable labeling of different RNA transcripts in the same cell. The power of SELEX to find selective and tight binders will be used to screen for mutually exclusive aptamer-cobalamin pairs. These pairs will be conjugated to spectrally-resolved fluorophores to enable tracking of multiple RNA simultaneously. These orthogonal probes will be used to image X and Y as they interact in the cell.

### **Aim 4. Application.**

Something here? Or would just applications at the end of aims 2 and 3 be okay?

## Bibliography and References Cited

- [1] Cech, T. R. and Steitz, J. A.: The Noncoding RNA Revolution—Trashing Old Rules to Forge New Ones. Cell. 157(1): 77–94, March 2014.
- [2] Müller-McNicoll, M. and Neugebauer, K. M. How cells get the message: Dynamic assembly and function of mRNA–protein complexes. <https://www.nature.com/articles/nrg3434>, March 2013.
- [3] Etzel, M. and Mörl, M.: Synthetic Riboswitches: From Plug and Pray toward Plug and Play. Biochemistry. 56(9): 1181–1198, March 2017.
- [4] Paige, J. S., Wu, K. Y., and Jaffrey, S. R.: RNA Mimics of Green Fluorescent Protein. Science. 333(6042): 642–646, July 2011.
- [5] Filonov, G. S., Moon, J. D., Svensen, N., and Jaffrey, S. R.: Broccoli: Rapid Selection of an RNA Mimic of Green Fluorescent Protein by Fluorescence-Based Selection and Directed Evolution. J. Am. Chem. Soc. 136(46): 16299–16308, November 2014.
- [6] Autour, A., Jeng, S., Cawte, A., Abdolazadeh, A., Galli, A., Panchapakesan, S. S. S., Rueda, D., Ryckelynck, M., and Unrau, P. J.: Fluorogenic RNA Mango aptamers for imaging small non-coding RNAs in mammalian cells. Nature Communications. 9(1): 656, February 2018.
- [7] Dolgosheina, E. V., Jeng, S. C. Y., Panchapakesan, S. S. S., Cojocaru, R., Chen, P. S. K., Wilson, P. D., Hawkins, N., Wiggins, P. A., and Unrau, P. J.: RNA Mango Aptamer-Fluorophore: A Bright, High-Affinity Complex for RNA Labeling and Tracking. ACS Chem. Biol. 9(10): 2412–2420, October 2014.
- [8] Fusco, D., Accornero, N., Lavoie, B., Shenoy, S. M., Blanchard, J.-M., Singer, R. H., and Bertrand, E.: Single mRNA Molecules Demonstrate Probabilistic Movement in Living Mammalian Cells. Current Biology. 13(2): 161–167, January 2003.