

Adding fluxes to consistent truncations: IIB supergravity on $\text{AdS}_3 \times S^3 \times S^3 \times S^1$

Camille Eloy^{a1}, Michele Galli^{b2}, Emanuel Malek^{b3}

^a *Theoretische Natuurkunde, Vrije Universiteit Brussel, and the International Solvay,
Institutes, Pleinlaan 2, B-1050 Brussels, Belgium*

^b *Institut für Physik, Humboldt-Universität zu Berlin,
IRIS Gebäude, Zum Großen Windkanal 2, 12489 Berlin, Germany*

Abstract

We use $E_{8(8)}$ Exceptional Field Theory to construct the consistent truncation of IIB supergravity on $S^3 \times S^3 \times S^1$ to maximal 3-dimensional $\mathcal{N} = 16$ gauged supergravity containing the $\mathcal{N} = (4, 4)$ AdS_3 vacuum. We explain how to achieve this by adding a 7-form flux to the S^1 reduction of the dyonic $E_{7(7)}$ truncation on $S^3 \times S^3$ previously constructed in the literature. Our truncation Ansatz includes, in addition to the $\mathcal{N} = (4, 4)$ vacuum, a host of moduli breaking some or all of the supersymmetries. We explicitly construct the uplift of a subset of these to construct new supersymmetric and non-supersymmetric AdS_3 vacua of IIB string theory, and show how the moduli space of AdS_3 vacua of six-dimensional gauged supergravity studied in [1] is compactified upon lifting to 10 dimensions. Along the way, we also derive the form of 3-dimensional $\mathcal{N} = 16$ gauged supergravity in terms of the embedding tensor and rule out a 10-/11-dimensional origin of some 3-dimensional gauged supergravities.

¹camille.elay@vub.be

²michele.galli@physik.hu-berlin.de

³emanuel.malek@physik.hu-berlin.de

Contents

1	Introduction	2
2	Review of $E_{8(8)}$ exceptional field theory	3
3	Consistent truncations to 3-dimensional $\mathcal{N} = 16$ gauged supergravity	5
4	Deriving the potential of $\mathcal{N} = 16$ gauged supergravity	6
4.1	Ruling out gaugings	9
5	Adding fluxes to consistent truncations	9
5.1	Adding fluxes to the S^3 truncation of 6-dimensional supergravity	10
5.1.1	The S^3 twist matrix	10
5.1.2	Adding flux	12
5.2	Adding flux to the $S^3 \times S^3 \times S^1$ truncation of IIB supergravity	13
5.2.1	The $S^3 \times S^3$ truncation	13
5.2.2	The $S^3 \times S^3 \times S^1$ truncation	14
5.2.3	Adding flux	14
6	Moduli of $\text{AdS}_3 \times S^3 \times S^3 \times S^1$	16
6.1	The undeformed $\mathcal{N} = (4, 4)$ $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ solution	16
6.2	Moduli in gauged supergravity	17
6.3	Moduli in 10 dimensions	18
6.4	Compactification of the moduli space	22
7	Conclusions	23
A	Representations of $D^1(2, 1; \alpha)$	23

To do:

- Think about title - mention new AdS_3 vacua? Lifting AdS_3 vacua?
- Check the $\text{SL}(n)$ equations
- Move 3-d gauged SUGRA to somewhere else??
- Construct 10-parameter uplift
- Complex geometry of moduli space?
- Compare with classification known in literature, e.g. in IIA (and IIB – Jerome’s story)
- Email DW and MP to ask who we should cite for (5.1)

1 Introduction

Consistent truncations are a powerful technique that simplifies the dynamics of 10-/11-dimensional supergravity by focusing on a restricted subset of fields. The key principle behind a consistent truncation is to truncate to a subsector of fields, such that all solutions of the truncated theory correspond to solutions of the original 10-/11-dimensional supergravity theory. By considering only a, typically finite, subset of fields, consistent truncations provide a powerful tool to find complicated new 10-/11-dimensional supergravity solutions and to study their deformations. Consistent truncations have proven particularly powerful in the AdS/CFT correspondence, since all well-understood AdS vacua of string theory do not admit scale-separation and, thus, cannot be studied using the usual tools of lower-dimensional effective theories.

Exceptional Field Theory (ExFT) is a reformulation of 10-/11-dimensional supergravity that unifies the metric and flux degrees of freedom and thereby makes manifest an exceptional symmetry group¹. Over the last decade, this has proven an extremely useful tool for constructing consistent truncations, leading to a number of new examples preserving various amounts of supersymmetry [2–18]. However, consistent truncations to three dimensions have, until recently [19], remained largely unexplored. One reason is that the $E_{8(8)}$ ExFT requires modifications for its local symmetry structure, i.e. the generalised Lie derivative, to close into an algebra [20, 21]. Another is that there are no maximally supersymmetric AdS_3 vacua, i.e. preserving 32 supercharges, of string theory, leaving no natural candidate for constructing a consistent truncation with vacua. Indeed, while the consistent truncations of 11-dimensional supergravity on S^7 and S^4 and of 10-dimensional supergravity on S^5 contain maximally supersymmetric AdS vacua and are captured by a universal Ansatz in ExFT [2, 3], the analogous S^8 truncation of 11-dimensional supergravity does not exist, while there are two S^7 truncations of 10-dimensional supergravity but neither contains a maximally symmetric vacuum [19, 22].

Nonetheless, there are half-maximal, i.e. $\mathcal{N} = (4, 4)$, supersymmetric AdS_3 vacua of string theory that are extremely intriguing. These are the $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ and $\text{AdS}_3 \times S^3 \times T^4$ of IIB string theory (realising the “large” and “small” $\mathcal{N} = (4, 4)$ superconformal symmetries), and, unlike in higher dimensions, can be supported by pure NS-NS flux. Not only does this mean that

¹For the purposes of this paper, we do not draw a distinction between ExFT and Exceptional Generalised Geometry, since these agree when the “section condition” is solved, which we will always assume here.

there are also heterotic versions of these vacua (preserving $\mathcal{N} = (4, 0)$ supersymmetry), but also that they can be readily studied via the string worldsheet CFT [23–26]. Moreover, the string sigma model in these vacua is also integrable [1]. Finally, the $\text{AdS}_3/\text{CFT}_2$ correspondence has seen much progress recently, with many new holographic duals being established [2].

Here we will focus on the $\mathcal{N} = (4, 4)$ $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ vacuum of IIB string theory and show that it admits a consistent truncation to a maximal gauged supergravity, which was first studied in [27]. All vacua of this theory break at least half the supersymmetries, reflecting the absence of a maximally supersymmetric AdS_3 vacuum in string theory. Key to constructing this consistent truncation is to add new fluxes to the S^1 reduction of the consistent truncation of IIB supergravity on $S^3 \times S^3$ [11].

Using our consistent truncation, we will show that the $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ has a rich moduli space of symmetry- and supersymmetry-breaking deformations, including some that are analogous to the “flat deformations” studied for $\text{AdS}_4 \times S^5 \times S^1$ [28–31]. While some of these deformations preserve some amounts of supersymmetry, others break all supersymmetries, yet, surprisingly, at least a subset of vacua continue to be perturbatively stable within IIB supergravity, similar to [31]. Our work also provides an uplift of 6-dimensional supergravity on S^3 studied in [1] to IIB string theory. As we will show, some deformations that appear non-compact in 6-dimensional supergravity [1] are compactified within the full 10-dimensional supergravity theory.

Another technical result of our work is the derivation of the potential of 3-dimensional gauged supergravity [32, 33] in terms of the embedding tensor, which was previously only expressed in terms of fermion shift matrices. As a by-product, we find that a constraint that must be obeyed by the embedding tensor of any 3-dimensional gauged supergravity that can be uplifted to 10-/11-dimensional supergravity. As we show, this allows us to prove a higher-dimensional origin of some 3-dimensional gaugings.

The outline of our paper is as follows. We begin with a review of $E_{8(8)}$ ExFT in section 2 and how to derive consistent truncations in this formalism in 3. In section 4 we derive the potential of 3-dimensional gauged supergravity in terms of the embedding tensor and prove the lack of higher-dimensional origin of some 3-dimensional theories. Then, we show how to add fluxes to consistent truncations in 5, allowing us to construct the consistent truncation of IIB supergravity on $S^3 \times S^3 \times S^1$. Using this truncation, we study the moduli space of the AdS_3 vacua in section 6, before concluding in section 7.

2 Review of $E_{8(8)}$ exceptional field theory

The $E_{8(8)}$ exceptional field theory, first constructed in ref. [20], is an $E_{8(8)}$ duality-covariant formulation of type II and 11d supergravities. It is defined on a set of $3 + 248$ coordinates made of three-dimensional external coordinates $\{x^\mu\}$ and internal coordinates $\{Y^M\}$ in the 248-dimensional adjoint representation of $E_{8(8)}$. The dependence of the fields on these coordinates is constrained by the “section constraints”²

$$\begin{cases} \eta^{MN} \partial_M \otimes \partial_N = 0, \\ f^{MN}{}_P \partial_M \otimes \partial_N = 0, \\ (\mathbb{P}_{3875})_{MN}{}^{KL} \partial_K \otimes \partial_L = 0, \end{cases} \quad (2.1)$$

²We use the notation \otimes to indicate that both derivatives may act on different functions.

where $f_{MN}{}^P$ are the structure constants of $E_{8(8)}$, $\eta_{MN} = \frac{1}{60} f_{MK}{}^L f_{NL}{}^K$ its Cartan-Killing metric and $(\mathbb{P}_{3875})_{MN}{}^{KL}$ is the projector on the representation **3875**:

$$(\mathbb{P}_{3875})_{MN}{}^{KL} = \frac{1}{7} \delta_{(M}{}^K \delta_{N)}{}^L - \frac{1}{56} \eta_{MN} \eta^{KL} - \frac{1}{14} f^P{}_{M(K} f_{PN}{}^{L)}. \quad (2.2)$$

Here and in the following, $E_{8(8)}$ indices are raised and lowered by the Cartan-Killing metric η_{MN} . The section constraints (2.1) ensure that the fields depends only on the 7- or 8-dimensional physical internal coordinates embeded in Y^M .

The theory describes the dynamics of the following bosonic fields:

$$\{g_{\mu\nu}, \mathcal{M}_{MN}, A_\mu{}^M, B_{\mu M}\}, \quad (2.3)$$

with $g_{\mu\nu}$ the 3-diminsinal external metric, \mathcal{M}_{MN} the generalized metric parametrizing the coset space $E_{8(8)}/\text{SO}(16)$ and the gauge fields $A_\mu{}^M$ and $B_{\mu M}$. It is a gauge theory, invariant under the generalized Lie derivative of parameters $\Upsilon = (\Lambda^M, \Sigma_M)$, whose action on a vector V^M of weight λ is given by

$$\mathcal{L}_\Upsilon V^M = \Lambda^N \partial_N V^M - 60 (\mathbb{P}_{248})^M{}_{NKL} V^N \partial_K \Lambda^L + \lambda V^M \partial_N \Lambda^N + f^{MN}{}_K \Sigma_N V^K, \quad (2.4)$$

with $(\mathbb{P}_{248})^M{}_{NKL} = (1/60) f^M{}_{NP} f^{PK}{}_L$ the projector on the adjoint representation. These transformations are well-defined (in particular, they close into an algebra) only if the parameters Σ_M and the fields $B_{\mu M}$ are covariantly constrained: they have to satisfy algebraic constraints similar to eq. (2.1) and be compatible with the partial derivatives. We require that

$$\begin{cases} \eta^{MN} C_M \otimes C'_N = 0, \\ f^{MN}{}_P C_M \otimes C'_N = 0, \\ (\mathbb{P}_{3875})_{MN}{}^{KL} C_K \otimes C'_L = 0, \end{cases} \quad \forall C_M, C'_M \in \{\partial_M, \Sigma_M, B_{\mu M}\}. \quad (2.5)$$

The bosonic action of $E_{8(8)}$ exceptional field theory is invariant under the transformations (2.4) and has the expression

$$S_{\text{ExFT}} = \int d^3x d^{248}Y \sqrt{|g|} \left(\hat{R} + \frac{1}{240} D_\mu \mathcal{M}_{MN} D^\mu \mathcal{M}^{MN} + \mathcal{L}_{\text{int}} + \frac{1}{\sqrt{|g|}} \mathcal{L}_{\text{CS}} \right). \quad (2.6)$$

\hat{R} is the $E_{8(8)}$ -covariantised Ricci scalar and the covariant derivative is defined as

$$D_\mu = \partial_\mu - \mathcal{L}_{(A_\mu, B_\mu)}. \quad (2.7)$$

\mathcal{L}_{int} is a potential term depending only on internal derivatives, explicitly

$$\begin{aligned} \mathcal{L}_{\text{int}} = & \frac{1}{240} \mathcal{M}^{MN} \partial_M \mathcal{M}^{KL} \partial_N \mathcal{M}_{KL} - \frac{1}{2} \mathcal{M}^{MN} \partial_M \mathcal{M}^{KL} \partial_L \mathcal{M}_{NK} \\ & - \frac{1}{7200} f^{NQ}{}_P f^{MS}{}_R \mathcal{M}^{PK} \partial_M \mathcal{M}_{QK} \mathcal{M}^{RL} \partial_N \mathcal{M}_{SL} \\ & + \frac{1}{2} g^{-1} \partial_M g \partial_N \mathcal{M}^{MN} + \frac{1}{4} \mathcal{M}^{MN} g^{-2} \partial_M g \partial_N g + \frac{1}{4} \mathcal{M}^{MN} \partial_M g_{\mu\nu} \partial_N g^{\mu\nu}. \end{aligned} \quad (2.8)$$

Finally, \mathcal{L}_{CS} is a Chern-Simons term required to impose the on-shell duality between scalar and vector fields, given by

$$\begin{aligned} \mathcal{L}_{\text{CS}} = & \frac{1}{2} \varepsilon^{\mu\nu\rho} \left(F_{\mu\nu}{}^M B_{\rho M} - f_{KL}{}^N \partial_\mu A_\nu{}^K \partial_N A_\rho{}^L - \frac{2}{3} f^N{}_{KL} \partial_M \partial_N A_\mu{}^K A_\nu{}^M A_\rho{}^L \right. \\ & \left. - \frac{1}{3} f_{MKL} f^{KP}{}_Q f^{LR}{}_S A_\mu{}^M \partial_P A_\nu{}^Q \partial_R A_\rho{}^S \right), \end{aligned} \quad (2.9)$$

where $F_{\mu\nu}{}^M$ is the covariant field strength of $A_\mu{}^M$ (see eq. (2.26) of ref. [20]).

3 Consistent truncations to 3-dimensional $\mathcal{N} = 16$ gauged supergravity

The $E_{8(8)}$ exceptional field theory is well suited to the construction of consistent truncation of type II and 11d supergravities to three-dimensional maximal supergravity. These truncations arise as generalised Scherk-Schwarz reductions, described by an $E_{8(8)}$ -valued twist matrix $U_M{}^{\bar{M}}$ and a scale factor ρ . The truncation Ansätze are as follows [19]:

$$\begin{aligned} g_{\mu\nu}(x, Y) &= \rho(Y)^{-2} \dot{g}_{\mu\nu}(x), \\ \mathcal{M}_{MN}(x, Y) &= U_M{}^{\bar{M}}(Y) U_N{}^{\bar{N}}(Y) \mathcal{M}_{\bar{M}\bar{N}}(x), \\ A_\mu{}^M(x, Y) &= \rho(Y)^{-1} (U^{-1})_{\bar{M}}{}^M(Y) \mathcal{A}_\mu{}^{\bar{M}}(x), \\ B_{\mu M}(x, Y) &= \Sigma_{\bar{M}M}(Y) \mathcal{A}_\mu{}^{\bar{M}}(x), \end{aligned} \quad (3.1)$$

with

$$\Sigma_{\bar{M}M} = \frac{\rho^{-1}}{60} f_{\bar{M}}{}^{\bar{P}\bar{Q}} (U^{-1})_{\bar{P}P} \partial_M (U^{-1})_{\bar{Q}}{}^P. \quad (3.2)$$

The fields with flat indices \bar{M} (in the **248** representation of $E_{8(8)}$) belong to the three-dimensional theory; all the dependence on the internal manifold is factored out in $U_M{}^{\bar{M}}$ and ρ . Note that with the definition of $\Sigma_{\bar{M}M}$ the condition (2.5) is automatically satisfied.

The Ansätze (3.1) describe a *consistent* truncation if the following condition is satisfied:

$$\mathcal{L}_{\mathfrak{U}_{\bar{M}}} \mathcal{U}_{\bar{N}}{}^M = X_{\bar{M}\bar{N}}{}^{\bar{P}} \mathcal{U}_{\bar{P}}{}^M, \quad (3.3)$$

with $\mathcal{U}_{\bar{M}}{}^M = \rho^{-1} (U^{-1})_{\bar{M}}{}^M$, $\mathfrak{U}_{\bar{M}} = (\mathcal{U}_{\bar{M}}, \Sigma_{\bar{M}})$ and $X_{\bar{M}\bar{N}}{}^{\bar{P}}$ a constant tensor. This condition ensures that the factorised form of the Ansätze (3.1) is preserved by the generalized Lie derivative, *e.g.*

$$\begin{aligned} D_\mu \mathcal{M}(x, Y) &= U_M{}^{\bar{M}}(Y) U_N{}^{\bar{N}}(Y) \mathcal{D}_\mu \mathcal{M}_{\bar{M}\bar{N}}(x), \\ \mathcal{L}_{(A_\mu, B_\nu)} A_\mu(x, Y) &= \rho(Y)^{-1} (U^{-1})_{\bar{M}}{}^M(Y) \llbracket \mathcal{A}_\mu, \mathcal{A}_\nu \rrbracket^{\bar{M}}(x), \end{aligned} \quad (3.4)$$

where **Note corrected sign in first equation!** (Sure? I took it from eq. (4.9) of ref. [19].)

$$\mathcal{D}_\mu \mathcal{M}_{\bar{M}\bar{N}} = \partial_\mu \mathcal{M}_{\bar{M}\bar{N}} + 2 \mathcal{A}_\mu{}^{\bar{P}} X_{\bar{P}(\bar{M}}{}^{\bar{Q}} \mathcal{M}_{\bar{N})\bar{Q}}, \quad \text{and} \quad \llbracket \mathcal{A}_\mu, \mathcal{A}_\nu \rrbracket^{\bar{M}} = X_{\bar{P}\bar{Q}}{}^{\bar{M}} \mathcal{A}_\mu{}^{\bar{P}} \mathcal{A}_\nu{}^{\bar{Q}}. \quad (3.5)$$

Thus, for a given twist matrix $U_M{}^{\bar{M}}$ satisfying the consistency condition (3.3), the action (2.6) reduces to three-dimensional $\mathcal{N} = 16$ gauged supergravity [32, 33]. The constant tensor $X_{\bar{M}\bar{N}}{}^{\bar{P}}$

plays the role of the embedding tensor of the three-dimensional gauged supergravity, as is clear from its appearance in (3.5) in the gauge-covariant derivative and the gauge algebra of the vector fields. From eq. (3.3), it has the following expression in terms of the twist matrix and ρ

$$\begin{aligned} X_{\overline{MN}}^{\overline{P}} = & -\rho^{-1} \Gamma_{\overline{MN}}^{\overline{P}} + \rho^{-1} f^{\overline{P}}_{\overline{NQ}} f^{\overline{QK}}_{\overline{L}} \Gamma_{\overline{KM}}^{\overline{L}} - \frac{1}{60} \rho^{-1} f^{\overline{PK}}_{\overline{N}} f_{\overline{ML}}^{\overline{Q}} \Gamma_{\overline{KQ}}^{\overline{L}} \\ & - \frac{1}{2} \rho^{-1} f^{\overline{P}}_{\overline{NQ}} f^{\overline{QK}}_{\overline{M}} \Gamma_{\overline{RK}}^{\overline{R}} + \left(\delta_{\overline{M}}^{\overline{K}} \delta_{\overline{N}}^{\overline{P}} - \frac{1}{2} f_{\overline{M}}^{\overline{LK}} f_{\overline{NL}}^{\overline{P}} \right) \xi_{\overline{K}}. \end{aligned} \quad (3.6)$$

Here, we defined the $E_{8(8)}$ current $\Gamma_{\overline{MN}}^{\overline{P}} = (U^{-1})_{\overline{M}}^{\overline{K}} (U^{-1})_{\overline{N}}^{\overline{L}} \partial_{\overline{K}} U_{\overline{L}}^{\overline{P}}$ and the trombone gauging

$$\xi_{\overline{M}} = 2 U_{\overline{M}}^{\overline{N}} \partial_{\overline{N}} \rho^{-1} - \rho^{-1} \Gamma_{\overline{NM}}^{\overline{N}}. \quad (3.7)$$

The embedding tensor is most nicely expressed once projected on the adjoint representation

$$\begin{aligned} X_{\overline{MN}} = & -\frac{1}{60} X_{\overline{MP}}^{\overline{Q}} f_{\overline{NQ}}^{\overline{P}} = -2 \rho^{-1} \Gamma_{(\overline{MN})} - \rho^{-1} \Gamma_{\overline{P}(\overline{M}}^{\overline{Q}} f_{\overline{N})\overline{Q}}^{\overline{P}} + \frac{1}{2} f_{\overline{MN}}^{\overline{P}} \xi_{\overline{P}} \\ = & -14 \rho^{-1} (\mathbb{P}_{3875})_{\overline{MN}}^{\overline{PQ}} \Gamma_{\overline{PQ}} - \frac{1}{4} \rho^{-1} \eta_{\overline{MN}} \Gamma_{\overline{P}}^{\overline{P}} + \frac{1}{2} f_{\overline{MN}}^{\overline{P}} \xi_{\overline{P}}, \end{aligned} \quad (3.8)$$

with the projection of the current $\Gamma_{\overline{MN}} = -\frac{1}{60} \Gamma_{\overline{MP}}^{\overline{Q}} f_{\overline{NQ}}^{\overline{P}}$. In the following, we will focus on gaugings with $\xi_{\overline{M}} = 0$.

The three-dimensional action follows from inserting the Ansätze (3.1) in the ExFT action (2.6). The expressions of the kinetic and Chern-Simons terms result immediately from eq. (3.4)

$$\begin{aligned} \mathcal{L}_{\text{kin}} = & \frac{1}{240} D_{\mu} \mathcal{M}_{MN} D^{\mu} \mathcal{M}^{MN} \stackrel{=}{=} \frac{\rho^2}{240} \mathcal{D}_{\mu} \mathcal{M}_{\overline{MN}} \mathcal{D}^{\mu} \mathcal{M}^{\overline{MN}}, \\ \mathcal{L}_{\text{CS}} \stackrel{=}{=} & -\rho^{-1} \varepsilon^{\mu\nu\rho} X_{\overline{MN}}^{\overline{P}} \mathcal{A}_{\mu}^{\overline{M}} \left(\partial_{\nu} \mathcal{A}_{\rho}^{\overline{N}} - \frac{1}{3} \llbracket \mathcal{A}_{\nu}, \mathcal{A}_{\rho} \rrbracket^{\overline{N}} \right). \end{aligned} \quad (3.9)$$

However, the case of the potential term \mathcal{L}_{int} , leading to a potential V for the three-dimensional scalars $\mathcal{M}_{\overline{MN}}$, is more subtle. We expect V to be quadratic in the embedding tensor, which makes the identification of $X_{\overline{MN}}^{\overline{P}}$ more complicated. Moreover, the potential of three-dimensional $\mathcal{N} = 16$ gauged supergravity in terms of the embedding tensor is unknown, with currently the potential only expressed in terms of the fermion shift matrices of the gauged supergravity [32, 33]. Thus, in the following, we will follow the truncation procedure carefully and thereby construct the potential of the three-dimensional $\mathcal{N} = 16$ gauged supergravity in terms of the embedding tensor.

4 Deriving the potential of $\mathcal{N} = 16$ gauged supergravity

Here we will use the $E_{8(8)}$ ExFT Lagrangian and the consistent truncation Ansatz (3.1) to derive the potential of three-dimensional $\mathcal{N} = 16$ gauged supergravity [32, 33] in terms of the embedding tensor. Not only is this an interesting application of ExFT, relying on purely bosonic considerations and bypassing the usual construction of the gauged supergravity potential using supersymmetry³,

³Note that the same strategy was recently used to impressively derive the potential of maximal two-dimensional gauged supergravity [34], where fermions are extremely poorly understood. As a result, in that case, the bosonic $E_{9(9)}$ ExFT [35, 36] provides the only currently accessible route for computing the gauged supergravity potential.

but the potential is crucial for finding vacua of the three-dimensional theory and uplifting these in the later parts of this paper.

In order to derive the potential, we adopt the following strategy. We want \mathcal{L}_{int} (2.8) to reduce to the embedding tensor squared upon inserting the generalised Scherk-Schwarz Ansätze (3.1). However, as in higher dimensions, e.g. [37–39], this does not have to match identically, but only up to total derivative terms and terms which violate the section condition. The possible boundary terms are

$$\frac{1}{\sqrt{|g|}} \partial_M \left(\sqrt{|g|} \partial_N \mathcal{M}^{MN} \right) \quad \text{and} \quad \frac{1}{\sqrt{|g|}} \partial_M \left(\mathcal{M}^{MN} \partial_N \sqrt{|g|} \right). \quad (4.1)$$

However, since the $\mathcal{N} = 16$ supergravity potential will be given by terms quadratic in $X_{\overline{MN}}^{\overline{P}}$, we do not want the boundary terms to involve any double derivative terms. Therefore, the two boundary terms (4.1) can only appear via the combination

$$\frac{1}{\sqrt{|g|}} \partial_M \left(\sqrt{|g|} \partial_N \mathcal{M}^{MN} + \frac{4}{3} \mathcal{M}^{MN} \partial_N \sqrt{|g|} \right). \quad (4.2)$$

Now, we first insert the generalized Scherk-Schwarz Ansätze (3.1) into the expression of \mathcal{L}_{int} (2.8) with the additional total derivative (4.2)

$$\begin{aligned} \mathcal{L}_{\text{int}} + \frac{a}{\sqrt{|g|}} \partial_M \left(\sqrt{|g|} \partial_N \mathcal{M}^{MN} + \frac{4}{3} \mathcal{M}^{MN} \partial_N \sqrt{|g|} \right) \\ \stackrel{\text{gSS}}{=} -\frac{1}{2} \mathcal{M}^{\overline{MN}} \Gamma_{\overline{MK}} \Gamma_{\overline{N}}^{\overline{K}} - \mathcal{M}^{\overline{MN}} \Gamma_{\overline{MK}} \Gamma_{\overline{N}}^{\overline{K}} - \frac{1}{2} \Gamma_{\overline{MN}} \Gamma^{\overline{NM}} - \frac{1}{2} \mathcal{M}^{\overline{MN}} \mathcal{M}^{\overline{KL}} \Gamma_{\overline{MK}} \Gamma_{\overline{NL}} \\ - \frac{1}{2} \mathcal{M}^{\overline{MN}} \mathcal{M}^{\overline{KL}} \Gamma_{\overline{MK}} \Gamma_{\overline{LN}} + \frac{3}{2} (a-1) \mathcal{M}^{\overline{MN}} \Gamma_{\overline{MN}}^{\overline{K}} \Gamma_{\overline{LK}}^{\overline{L}} - \frac{a}{2} \mathcal{M}^{\overline{MN}} \Gamma_{\overline{KM}}^{\overline{K}} \Gamma_{\overline{LN}}^{\overline{L}} \\ + \left(\frac{1}{2} - a \right) \mathcal{M}^{\overline{MN}} \Gamma_{\overline{KM}}^{\overline{L}} \Gamma_{\overline{LN}}^{\overline{K}} + \mathcal{M}^{\overline{MN}} \Gamma_{\overline{MK}}^{\overline{L}} \Gamma_{\overline{LN}}^{\overline{K}} - \frac{1}{2} \mathcal{M}^{\overline{MN}} \mathcal{M}^{\overline{KL}} f_{\overline{QL}}^{\overline{P}} \Gamma_{\overline{PN}}^{\overline{Q}} \Gamma_{\overline{MK}}. \end{aligned} \quad (4.3)$$

Secondly, knowing that the $\mathcal{N} = 16$ potential must be quadratic in $X_{\overline{MN}}$, we consider the most general quadratic function in $X_{\overline{MN}}$ and develop it in $E_{8(8)}$ currents using eq. (3.8). Note that in higher dimensions, there exist only two quadratic combinations of the embedding tensor

$$X_{\overline{MN}}^{\overline{P}} X_{\overline{QP}}^{\overline{N}} \mathcal{M}^{\overline{MQ}} \quad \text{and} \quad X_{\overline{MN}}^{\overline{P}} X_{\overline{QR}}^{\overline{S}} \mathcal{M}^{\overline{MQ}} \mathcal{M}^{\overline{NR}} \mathcal{M}_{\overline{PS}}. \quad (4.4)$$

However, in three dimensions, because the vector fields transform in the adjoint representation of $E_{8(8)}$, we can write two additional terms

$$X_{\overline{MN}} X_{\overline{PQ}} \eta^{\overline{MP}} \eta^{\overline{NQ}} \quad \text{and} \quad X_{\overline{MN}} X_{\overline{PQ}} \eta^{\overline{MN}} \eta^{\overline{PQ}}, \quad (4.5)$$

where the second term corresponds to the square of the singlet part of the embedding tensor.

It turns out that for gaugings that arise from consistent truncations, these two terms (4.5) can be related to the square of the trombone tensor. We can square eq. (3.8) to derive an expression for $X_{\overline{MN}} X^{\overline{MN}}$ in terms of the current Γ . We find **check coefficient of last term**

$$X_{\overline{MN}} X^{\overline{MN}} = 21 \rho^{-2} \Gamma_{\overline{MN}} \Gamma^{\overline{NM}} + 19 \rho^{-2} \left(\Gamma_{\overline{M}}^{\overline{M}} \right)^2 - 15 \xi_{\overline{M}} \xi^{\overline{M}}. \quad (4.6)$$

Similarly, squaring the trombone tensor ξ gives

$$\xi_{\bar{M}} \xi^{\bar{M}} = \rho^{-2} \left(\Gamma_{\bar{M}}^{\bar{M}} \Gamma_{\bar{N}}^{\bar{N}} + \Gamma_{\bar{M}\bar{N}} \Gamma^{\bar{N}\bar{M}} \right). \quad (4.7)$$

We can now deduce

$$X_{\bar{M}\bar{N}} X^{\bar{M}\bar{N}} = 6 \xi_{\bar{M}} \xi^{\bar{M}} - 2 \rho^{-2} \left(\Gamma_{\bar{M}}^{\bar{M}} \right)^2, \quad (4.8)$$

which can be expressed entirely in terms of $X_{\bar{M}\bar{N}}$ after tracing eq. (3.8)

$$X_{\bar{M}\bar{N}} X^{\bar{M}\bar{N}} = 6 \xi_{\bar{M}} \xi^{\bar{M}} - \frac{1}{1922} \left(X_{\bar{M}}^{\bar{M}} \right)^2. \quad (4.9)$$

Thus, (4.9) is a constraint on the embedding tensor of three-dimensional supergravity which must be satisfied in order for it to have an uplift to 10-/11-dimensional supergravity. Note that this constraint is not implied by the quadratic constraint. To see this, simply observe that the theory with $E_{8(8)}$ gauging has $X_{\bar{M}\bar{N}} = \eta_{\bar{M}\bar{N}}$ and $\xi_{\bar{M}} = 0$ [32, 33], violating (4.9).

Since we are focusing on gaugings with vanishing trombone $\xi_{\bar{M}} = 0$, (4.9) implies that the two terms in (4.5) are proportional and we only need to consider one of these terms in the gauged supergravity potential. Thus, the most general Ansatz for the supergravity potential, for gaugings that have an uplift to 10-/11-dimensional supergravity⁴, is

$$\begin{aligned} -V &= X_{\bar{M}\bar{N}} X_{\bar{P}\bar{Q}} \left(\alpha \mathcal{M}^{\bar{M}\bar{P}} \mathcal{M}^{\bar{N}\bar{Q}} + \beta \mathcal{M}^{\bar{M}\bar{P}} \eta^{\bar{N}\bar{Q}} + \delta \eta^{\bar{M}\bar{P}} \eta^{\bar{N}\bar{Q}} \right) \\ &\stackrel{\text{gSS}}{=} \rho^{-2} \left(14 \alpha \Gamma_{\bar{M}\bar{N}} \Gamma_{\bar{P}\bar{Q}} \mathcal{M}^{\bar{M}\bar{P}} \mathcal{M}^{\bar{N}\bar{Q}} + 14 \alpha \Gamma_{\bar{M}\bar{N}} \Gamma_{\bar{P}\bar{Q}} \mathcal{M}^{\bar{M}\bar{Q}} \mathcal{M}^{\bar{N}\bar{P}} \right. \\ &\quad + 2 \beta \mathcal{M}^{\bar{M}\bar{N}} \Gamma_{\bar{M}\bar{K}} \Gamma_{\bar{N}}^{\bar{K}} + \beta \mathcal{M}^{\bar{M}\bar{N}} \Gamma_{\bar{M}\bar{K}} \Gamma_{\bar{N}}^{\bar{K}} + (-12 \alpha + 2 \delta) \Gamma_{\bar{M}\bar{N}} \Gamma^{\bar{M}\bar{N}} \\ &\quad - 2 \beta \mathcal{M}^{\bar{M}\bar{N}} \Gamma_{\bar{M}\bar{K}}^{\bar{L}} \Gamma_{\bar{L}\bar{N}}^{\bar{K}} + \beta \mathcal{M}^{\bar{M}\bar{N}} \Gamma_{\bar{K}\bar{M}}^{\bar{L}} \Gamma_{\bar{L}\bar{N}}^{\bar{K}} + \left(7 \alpha + \frac{\beta}{2} \right) \mathcal{M}^{\bar{M}\bar{N}} \Gamma_{\bar{K}\bar{M}}^{\bar{K}} \Gamma_{\bar{L}\bar{N}}^{\bar{L}} \\ &\quad \left. + 14 \alpha \mathcal{M}^{\bar{M}\bar{N}} \mathcal{M}^{\bar{K}\bar{L}} f_{\bar{Q}\bar{L}}^{\bar{P}} \Gamma_{\bar{P}\bar{N}}^{\bar{Q}} \Gamma_{\bar{M}\bar{K}} \right). \end{aligned} \quad (4.10)$$

Finally, by requiring

$$\sqrt{|g|} \mathcal{L}_{\text{int}} \stackrel{\text{gSS}}{=} -\sqrt{|\dot{g}|} V, \quad (4.11)$$

the parameters in (4.3) and (4.10) are fixed to

$$\alpha = \frac{1}{28}, \quad \beta = \frac{1}{2}, \quad \delta = \frac{13}{28}, \quad a = 1. \quad (4.12)$$

With the generalized Scherk-Schwarz Ansätze (3.1), the action (2.6) becomes

$$\begin{aligned} S_{\text{ExFT}} \stackrel{\text{gSS}}{=} \int d^{248} Y \rho^{-1} \int d^3 x \left[\sqrt{|\dot{g}|} \left(\dot{R} + \frac{1}{240} \mathcal{D}_{\mu} \mathcal{M}_{\bar{M}\bar{N}} \mathcal{D}^{\mu} \mathcal{M}^{\bar{M}\bar{N}} - V \right) \right. \\ \left. - \varepsilon^{\mu\nu\rho} X_{\bar{M}\bar{N}} \mathcal{A}_{\mu}^{\bar{M}} \left(\partial_{\nu} \mathcal{A}_{\rho}^{\bar{N}} - \frac{1}{3} [\mathcal{A}_{\nu}, \mathcal{A}_{\rho}]^{\bar{N}} \right) \right], \end{aligned} \quad (4.13)$$

⁴If we do not require the three-dimensional gauged supergravity to arise from a consistent truncation, (4.9) may be violated and we need to include both terms of (4.5). However, here we are not interested in such gaugings.

with the potential

$$V = X_{\overline{MN}} X_{\overline{PQ}} \left(\frac{1}{28} \mathcal{M}^{\overline{MP}} \mathcal{M}^{\overline{NQ}} + \frac{1}{2} \mathcal{M}^{\overline{MP}} \eta^{\overline{NQ}} + \frac{13}{28} \eta^{\overline{MP}} \eta^{\overline{NQ}} \right). \quad (4.14)$$

As in higher dimensions, $X_{\overline{MN}}$ obtained from (3.3) automatically satisfies the linear constraint of the 3-dimensional maximal gauged supergravity [32, 33], and the section condition implies the quadratic constraints for $X_{\overline{MN}}$.

4.1 Ruling out gaugings

Let's check the $\text{GL}(6) \times \text{SO}(2, 2)$ gauging (half-maximal embedded in maximal)....

5 Adding fluxes to consistent truncations

In order to construct the consistent truncation around the $\mathcal{N} = (4, 4)$ $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ vacuum of IIB supergravity, a promising starting point is the dyonic $S^3 \times S^3$ truncation of IIB supergravity constructed in [11], and further reducing this on S^1 . However, this will not give the correct AdS_3 vacuum since the truncation is missing the required 7-form flux on $S^3 \times S^3 \times S^1$. Indeed, the 3-dimensional gauged supergravity that would be obtained this way does not have any AdS_3 vacua. We can remedy this situation by defining a new consistent truncation by adding a 7-form flux to the one obtained from the $S^3 \times S^3$ reduction constructed in $E_{7(7)}$ ExFT.

This motivates the following question: Given a consistent truncation, i.e. $\mathcal{U}_{\overline{M}}^M$, satisfying (3.3), when can we add a new flux component of string theory to the compactification to obtain a new consistent truncation? Adding a new flux component to the truncation is equivalent to twisting the generalised frame as follows [Email DW and MP to ask who we should cite for this?](#)

$$\mathcal{U}_{\overline{M}}^M \longrightarrow \mathcal{U}'_{\overline{M}}{}^M = \mathcal{U}_{\overline{M}}^N \exp(C)_N{}^M, \quad (5.1)$$

where C denotes the $E_{8(8)}$ generator corresponding to the potential we want to add to the compactification. The effect of the twist (5.1) is that the generalised Lie derivative of \mathcal{U}' satisfies

$$\mathcal{L}_{\mathcal{U}'_{\overline{M}}} \mathcal{U}'_{\overline{N}}{}^M = \left(\mathcal{L}_{\mathcal{U}_{\overline{M}}} \mathcal{U}_{\overline{N}}^N + F_{PQ}{}^N \mathcal{U}_{\overline{M}}^P \mathcal{U}_{\overline{N}}^Q \right) \exp(C)_N{}^M, \quad (5.2)$$

where $F_{MN}{}^P$ is a tensor in the $\mathbf{1} \oplus \mathbf{248} \oplus \mathbf{3875}$ of $E_{8(8)}$, i.e. the same representation as the embedding tensor, corresponding to the field strength of the potential C in (5.1). Using (3.3), we now have

$$\mathcal{L}_{\mathcal{U}'_{\overline{M}}} \mathcal{U}'_{\overline{N}}{}^M = \left(X_{\overline{MN}}{}^{\overline{P}} + \rho^{-1} F_{\overline{MN}}{}^{\overline{P}} \right) \mathcal{U}'_{\overline{P}}{}^M, \quad (5.3)$$

where we defined

$$F_{\overline{MN}}{}^{\overline{P}} \equiv (U^{-1})_{\overline{M}}{}^M (U^{-1})_{\overline{N}}{}^N U_P{}^{\overline{P}} F_{MN}{}^P, \quad (5.4)$$

and $X_{\overline{MN}}{}^{\overline{P}}$ is already constant. Therefore, we have a consistent truncation if and only if $\rho^{-1} F_{\overline{MN}}{}^{\overline{P}}$ is constant.

A particularly simple way of having constant $\rho^{-1} F_{\overline{MN}}{}^{\overline{P}}$ is to switch on fluxes which are stabilised by the twist matrix $U_M{}^{\overline{M}}$, which typically only lives in a subgroup $G \subset E_{8(8)}$. Therefore, in this case, we can simply tune the G -singlet components of the flux $F_{MN}{}^P$ to be proportional to ρ to obtain a new consistent truncation.

5.1 Adding fluxes to the S^3 truncation of 6-dimensional supergravity

As a warm-up for the $S^3 \times S^3 \times S^1$ truncation, let us demonstrate this methodology for the consistent truncation of $\mathcal{N} = (1, 1)$ 6-dimensional supergravity on S^3 , which was constructed in [1]. As discussed in [1], the consistent truncation of $\mathcal{N} = (1, 1)$ 6-dimensional supergravity to 3-dimensional half-maximal gauged supergravity can be described using the $\text{SO}(8, 4)$ ExFT [40, 41]. On the other hand, the consistent truncation on S^3 is conveniently described by twist matrices living in $\text{SL}(4) \simeq \text{SO}(3, 3)$ [2, 3, 6] I think the twist matrix of [2, 3, 6] coincide in the $\text{SL}(4)$ case, right? Camille, do you agree? (Yes.) with the embedding $\text{SO}(3, 3) \subset \text{SO}(4, 4) \subset \text{SO}(8, 4)$. However, the twist matrix in [1] differs from this $\text{SO}(3, 3)$ twist matrix [2, 3, 6] by an additional parameter, λ , which gives rise to an external 3-form flux or, equivalently, a new internal 3-form flux.

5.1.1 The S^3 twist matrix

Move to appendix?!?! (I think I would say yes)

Since we will use it throughout this paper, let us construct the $\text{SL}(4) \simeq \text{SO}(3, 3)$ twist matrix that describes the consistent truncation on S^3 . This can be viewed as a special case of a family of consistent truncations on S^n [2, 3, 6]. Here, we give a slightly different, but equivalent, form of this construction.

We use an $\text{SL}(n+1)$ ExFT, like in [3], which encodes an n -dimensional metric and volume-form flux, i.e. Freund-Rubin compactifications, and thus gives a natural description of S^n compactifications. The $\text{SL}(n+1)$ ExFT formally has coordinates in the antisymmetric representation of $\text{SL}(n+1)$, $y^{IJ} = -y^{JI}$, with $I = 1, \dots, n+1$, and similarly, generalised vector fields transform in the antisymmetric representation of $\text{SL}(n+1)$. The generalised Lie derivative on a generalised tensor in the fundamental V^I of weight λ is given by

$$\mathcal{L}_\Lambda V^I = \frac{1}{2} \Lambda^{JK} \partial_{JK} V^I - V^J \partial_{JK} \Lambda^{IK} + \left(\frac{\lambda}{2} + \frac{1}{n+1} \right) V^I \partial_{JK} \Lambda^{JK}, \quad (5.5)$$

and similarly for other generalised tensors. Closure of the generalised Lie derivative (5.5) requires the section condition

$$\partial_{[IJ} \otimes \partial_{KL]} = 0, \quad (5.6)$$

which restricts the dependence of all fields to a subset of physical coordinates.

We are interested in the maximal solutions of the section condition (5.6) which preserve a $\text{SL}(n) \subset \text{SL}(n+1)$ subgroup. Under this decomposition $\text{SL}(n+1) \rightarrow \text{SL}(n)$ with $V^I = (V^i, V^0)$, where $i = 1, \dots, n$, we solve the section condition by having physical coordinates y^{0i} on the n -dimensional manifold M , i.e. $\partial_{ij} = 0$ for all fields with only $\partial_{0i} = \partial_i \neq 0$. Correspondingly, the generalised tangent bundle, whose sections are in the antisymmetric representation of $\text{SL}(n+1)$ and carry weight $\frac{n-3}{n+1}$, decomposes as

$$E = TM \oplus \Lambda^{n-2} T^* M, \quad (5.7)$$

Note that for the case of interest to us, $n = 3$, (5.7) reduces to

$$E = TM \oplus T^* M, \quad (5.8)$$

and its fibres transform in the **6** of $\text{SL}(4) \simeq \text{SO}(3, 3)$. It is convenient to also introduce the generalised bundle with fibres in the anti-fundamental of $\text{SL}(n+1)$, which is, similarly, given by

$$N = T^* M \oplus \Lambda^n T^* M. \quad (5.9)$$

Sections of N are generalised tensors transforming in the anti-fundamental of $\text{SL}(n+1)$ and carrying weight $\frac{2}{n+1}$. For $V, V' \in \Gamma(E)$ and $W \in \Gamma(N)$, the generalised Lie derivative (5.5) reduces to

$$\begin{aligned}\mathcal{L}_V V' &= [v, v'] + L_v \omega'_{(n-2)} - \iota_{v'} d\omega_{(n-2)}, \\ \mathcal{L}_V W &= L_v \omega_{(1)} + L_v \omega_{(n)} + \omega_{(1)} \wedge d\omega_{(n-2)},\end{aligned}\tag{5.10}$$

where we write $V = v + \omega_{(n-2)}$, $V' = v' + \omega'_{(n-2)}$ and $W = \omega_{(1)} + \omega_{(n)}$ as a formal of vectors and p -forms, and L denotes the ordinary Lie derivative and $[v, v']$ the ordinary Lie bracket between vector fields v and v' .

We can now describe the S^n consistent truncation using the $\text{SL}(n+1)$ ExFT above. Let Y^I , $I = 1, \dots, n+1$ be the embedding coordinates of $S^n \subset \mathbb{R}^{n+1}$, such that $Y^I Y_I = 1$. On S^n , we can define a parallelisation of N using the generalised frame

$$\mathcal{U}^{\bar{I}} = dY^{\bar{I}} + Y^{\bar{I}} \text{vol}_{S^n} - dY^{\bar{I}} \wedge A,\tag{5.11}$$

where A is an $(n-1)$ -potential with field strength $dA = (n-1)\text{vol}_{S^n}$, and

$$\text{vol}_{S^n} = \frac{1}{n!} \epsilon_{I_1 I_2 \dots I_{n+1}} Y^{I_1} dY^{I_2} \wedge \dots \wedge dY^{I_{n+1}},\tag{5.12}$$

is the volume form on S^n . The frame in (5.11) has the important property that the $n+1$ generalised tensors are nowhere vanishing since $dY^I = 0$ only when $Y^I = 1$. Therefore, (5.11) provides a generalised parallelisation of S^n in the $\text{SL}(n+1)$ ExFT.

In a local basis, (5.11) gives us a $\text{GL}(n+1)$ matrix, $\mathcal{U}_I^{\bar{I}}$, whose determinant allows us to define the scalar density

$$\rho = \left(\det \mathcal{U}_I^{\bar{I}} \right)^{-1/2} = (\det \hat{g})^{-1/2},\tag{5.13}$$

with \hat{g} the round metric on S^n . Using (5.11) and (5.13) we can define the $\text{SL}(n)$ twist matrix

$$U_I^{\bar{I}} = \rho^{2/(n+1)} \mathcal{U}_I^{\bar{I}},\tag{5.14}$$

and hence a generalised frame for E or a generalised bundle in any other rep of $\text{SL}(n+1)$. For example, in the generalised tangent bundle (5.7), we have the generalised frame $\mathcal{U}_{IJ}^{IJ} = (\det \hat{g})^{1/2} \mathcal{U}_{\bar{I}}^I \mathcal{U}_{\bar{J}}^J$, with components given by

$$2\mathcal{U}_{IJ} = v_{IJ} + \star dY_{IJ} + \iota_{v_{IJ}} A,\tag{5.15}$$

where v_{IJ} are the $\text{SO}(n+1)$ Killing vectors of the round S^n , matching precisely the $\text{SL}(n+1)$ parallelisations constructed in [3].

Using (5.5), we can easily check that the generalised frame (5.15), (5.19) defines a generalised Leibniz parallelisation. In particular, using $dA = (n-1)\text{vol}_{S^n}$, we obtain the embedding tensor

$$\mathcal{L}_{\mathcal{U}_{IJ}} \mathcal{U}^{\bar{K}} = -X_{\bar{I}\bar{J}\bar{L}}^{\bar{K}} \mathcal{U}^{\bar{L}},\tag{5.16}$$

with

$$X_{\bar{I}\bar{J},\bar{L}}^{\bar{K}} = -\delta_{\bar{L}[\bar{I}} \delta_{\bar{J}]}^{\bar{K}}.\tag{5.17}$$

Finally, we can also connect the above with the $\text{SL}(n+1)$ Ansatz construction of [2]. We do this by evaluating (5.14) and (5.13) on the northern hemisphere $Y^I = (y^i, \sqrt{1-y^i y_i})$, $i = 1, \dots, n$ and using the gauge choice for the potential A

$$A_{i_1 \dots i_{n-1}} = \epsilon_{i_1 \dots i_{n-1} j} y^j (1 + K(v)), \quad (5.18)$$

with $v = y^i y_i$ and $\epsilon_{i_1 \dots i_{n-1} j}$ the volume form on S^n . With the above choices, we precisely recover the $\text{SL}(n+1)$ twist matrix of [2], i.e.

$$(U^{-1})_{\bar{I}}^I = \begin{pmatrix} (1-v)^{-1/(n+1)} (\delta_i^j + y_i y^j K(v)) & y_i (1-v)^{(n-1)/(2(n+1))} \\ y^j (1-v)^{(n-1)/(2(n+1))} K(v) & (1-v)^{\frac{n}{n+1}} \end{pmatrix}. \quad (5.19)$$

5.1.2 Adding flux

To demonstrate our methodology, we will now show how the parameter λ introduced in [1] can be obtained by the twisting procedure described above. Thus, we want to consider the $\text{SO}(3,3) \subset \text{SO}(4,4)$ twist matrix corresponding to S^3 (5.19) and add a 3-form flux via a $\text{SO}(4,4)$ twist. We begin by decomposing $\text{SO}(4,4) \rightarrow \text{SO}(3,3) \times \text{SO}(1,1)$, such that

$$\begin{aligned} \mathbf{8} &\rightarrow \mathbf{6_0} \oplus \mathbf{1_2} \oplus \mathbf{1_{-2}}, \\ \mathbf{28} &\rightarrow \mathbf{15_0} \oplus \mathbf{6_2} \oplus \mathbf{6_{-2}} \oplus \mathbf{1_0}. \end{aligned} \quad (5.20)$$

Correspondingly, we write a $\text{SO}(4,4)$ vector V^M as

$$V^M = (V^A, V^z, V^{\bar{z}}), \quad (5.21)$$

where $A = 1, \dots, 6$ labels the vector of $\text{SO}(3,3)$ and z, \bar{z} label the $\mathbf{1_2}$ and $\mathbf{1_{-2}}$, respectively.

We denote by $U_A^{\bar{A}}$ the $\text{SO}(3,3)$ twist matrix corresponding to the S^3 truncation constructed in [3, 6] and obtained from (5.14), (5.13), (5.19). Then, we can add a 3-form potential by twisting with the $\text{SO}(4,4)$ generator

$$(e^C)^{MN} = C_A (t^A)^{MN}, \quad (5.22)$$

with C_A an element of the $\mathbf{6_2}$, and $(t^A)^{MN}$ the generator corresponding to $\text{SO}(4,4)$ whose only non-zero component is

$$(t^A)^{Bz} = \eta^{AB}. \quad (5.23)$$

Here η_{MN} and η_{AB} are the $\text{SO}(4,4)$ and $\text{SO}(3,3)$ invariant metrics, respectively, and are used to raise/lower the corresponding vector indices. Decomposing with respect to the geometric $\text{SL}(3) \times \mathbb{R}^+$ of the S^3 , the coordinates on S^3 live in the $Y^A = (y^i, y_i)$, $i = 1, 2, 3$, while $C_A = (C_i, C^i)$ naturally contains a 2-form C^i . The field strength $H_3 = \partial^A C_A = \partial_i C^i$ is a singlet of $\text{SO}(3,3)$ and thus we see that $\rho^{-1} F_{\overline{MN}}^{\overline{P}}$ in (5.4) is constant and we obtain a consistent truncation with a new 3-form flux. Evaluating the twist matrix

$$U'_M{}^{\overline{M}} = (e^C)_M{}^N U_N{}^{\overline{M}}, \quad (5.24)$$

explicitly using (5.22) and setting $C^i = \lambda \xi^i$, (factor ρ^{-1} ?) with $\nabla_i \xi^i = 1$, in the notation of [1], we obtain precisely the twist matrix used in [1] for the consistent truncation of 6-dimensional $\mathcal{N} = (1,1)$ supergravity on S^3 with H_3 flux.

We will now follow this same procedure in the next section to obtain the consistent truncation of IIB supergravity on $S^3 \times S^3 \times S^1$ with 7-form flux.

5.2 Adding flux to the $S^3 \times S^3 \times S^1$ truncation of IIB supergravity

Our strategy for constructing the consistent truncation on $S^3 \times S^3 \times S^1$ with 7-form flux is to embed the $S^3 \times S^3$ truncation of $E_{7(7)}$ ExFT [11] into $E_{8(8)} \rightarrow E_{7(7)} \times \text{SL}(2)$, as described in [19]. This will give us the consistent truncation on $S^3 \times S^3 \times S^1$ without 7-form flux. Then we add a 7-form flux to this truncation as outlined above.

5.2.1 The $S^3 \times S^3$ truncation

Let us first review the dyonic $S^3 \times S^3$ truncation of IIB supergravity [11], which forms the starting point of our construction. The key step in the construction of [11] is that we can use the $\text{SL}(4)$ generalised frame (5.14) to form a generalised Leibniz parallelisation of $E_{7(7)}$ via the embedding

$$E_{7(7)} \rightarrow \text{SL}(8) \rightarrow \text{SL}(4)_1 \times \text{SL}(4)_2 \times \mathbb{R}^+. \quad (5.25)$$

The fundamental of $E_{7(7)}$ then decomposes as

$$\mathbf{56} \rightarrow \mathbf{28} \oplus \overline{\mathbf{28}} \rightarrow [(\mathbf{6}, \mathbf{1})_2 \oplus (\mathbf{1}, \mathbf{6})_{-2} \oplus (\mathbf{4}, \mathbf{4})_0] \oplus [(\mathbf{6}, \mathbf{1})_{-2} \oplus (\mathbf{1}, \mathbf{6})_2 \oplus (\overline{\mathbf{4}}, \overline{\mathbf{4}})_0], \quad (5.26)$$

where the first square brackets denote the branching of the $\mathbf{28}$ under $\text{SL}(4)_1 \times \text{SL}(4)_2 \times \mathbb{R}^+$ and the second square brackets that of the $\overline{\mathbf{28}}$. Crucially, for the generalised Leibniz condition (3.3) to hold, the coordinates and generalised vector fields corresponding to the $\text{SL}(4)_1$ and $\text{SL}(4)_2$ must be embedded within the $\mathbf{28}$ and $\overline{\mathbf{28}}$, respectively, which we will call “electric” and “magnetic” coordinates, following [11]. Thus, we write the 56 $E_{7(7)}$ coordinates as (*M already used for adjoint of $E_{8(8)}$*)

$$Y^M = (Y^{AB}, Y_{AB}), \quad (5.27)$$

with $A, B = 1, \dots, 8$ labelling the fundamental of $\text{SL}(8)$, corresponding to the decomposition $E_{7(7)} \rightarrow \text{SL}(8)$. Following the conventions of [11], the coordinates of the two $\text{SL}(4)$ ExFTs are embedded as $Y^{IJ} \subset Y^{AB}$, with $I, J = 1, 2, 3, 8$, and $Y_{AB} \subset Y_{AB}$, with $A, B = 4, 5, 6, 7$. Solving the $\text{SL}(4)$ ExFT section condition for Y^{IJ} and Y_{AB} guarantees a solution to the $E_{7(7)}$ ExFT, and we choose the solution where the six physical coordinates of IIB are

$$y^i = Y^{i8}, \quad \tilde{y}_a = Y_{a7}, \quad i = 1, 2, 3, \quad a = 4, 5, 6. \quad (5.28)$$

This solution of the section condition defines the “geometric” $\text{SL}(3)_1 \times \mathbb{R}_1^+ \subset \text{SL}(4)_1$ and $\text{SL}(3)_2 \times \mathbb{R}_2^+ \subset \text{SL}(4)_2$ subgroups, that will play an important role when adding flux in section 5.2.3.

We can now use one copy of the $\text{SL}(4)$ frame (5.14) for each $\text{SL}(4)$ subgroup of (5.25) to construct a generalised parallelisation for the full $E_{7(7)}$ ExFT, with an embedding tensor via (3.3) given as in (5.17) embedded in $E_{7(7)}$ as follows. Under $E_{7(7)} \rightarrow \text{SL}(8)$, the embedding tensor representation branches as

$$\mathbf{912} \rightarrow \mathbf{36} \oplus \overline{\mathbf{36}} \oplus \mathbf{420} \oplus \overline{\mathbf{420}}, \quad (5.29)$$

and we only generate the $\mathbf{36}$ and $\overline{\mathbf{36}}$ components, η_{AB}, η^{AB} . Thus, the $E_{7(7)}$ embedding tensor is given by

$$\begin{aligned} X_{AB,CD}{}^{\mathcal{EF}} &= \eta_{A[C} \delta_{D]B}^{\mathcal{EF}} - \eta_{B[C} \delta_{D]A}^{\mathcal{EF}}, \\ X^{AB}{}_{CD}{}^{\mathcal{EF}} &= -\eta^{A[\mathcal{E}} \delta_{CD}^{\mathcal{F}]B} + \eta^{B[\mathcal{E}} \delta_{CD}^{\mathcal{F}]A}. \end{aligned} \quad (5.30)$$

Because we are embedding the $\text{SL}(4)_1$ coordinates in the **28** but those of $\text{SL}(4)_2$ in the $\overline{\mathbf{28}}$, the twist matrix will generate an embedding tensor in the **36** for $\text{SL}(4)_1$ and $\overline{\mathbf{36}}$ for $\text{SL}(4)_2$, whose only non-vanishing components are

$$\eta_{IJ} = \delta_{IJ}, \quad \eta^{AB} = \delta^{AB}. \quad (5.31)$$

We can also rescale their S^3 's using their corresponding \mathbf{R}^+ generator. Without loss of generality, we can rescale S^3 to have radius α , using the \mathbf{R}_1^+ generator on the frame. The embedding tensor then becomes

$$\eta_{IJ} = \frac{1}{\alpha} \delta_{IJ}, \quad \eta^{AB} = \delta^{AB}. \quad (5.32)$$

5.2.2 The $S^3 \times S^3 \times S^1$ truncation

We now construct the consistent truncation on $S^3 \times S^3 \times S^1$ by embedding the $E_{7(7)}$ twist matrix corresponding to the $S^3 \times S^3$ truncation reviewed above in 5.2.1 in $E_{8(8)}$ via the branching $E_{8(8)} \rightarrow E_{7(7)} \times \text{SL}(2)$, such that $\mathbf{248} \rightarrow (\mathbf{133}, \mathbf{1}) \oplus (\mathbf{56}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3})$. As explained in [19], we can construct a consistent truncation on $M \times S^1$ to 3-dimensional gauged supergravity by embedding a consistent truncation on M to 4-dimensional supergravity, characterised by an $E_{7(7)}$ twist matrix and $E_{7(7)}$ scalar density σ , as well as the $\text{SL}(2)$ twist matrix given by

$$v_i^{\bar{i}} = \begin{pmatrix} \sigma & 0 \\ 0 & \sigma^{-1} \end{pmatrix}. \quad (5.33)$$

Finally, the $E_{8(8)}$ scalar density satisfies $\rho = \sigma^2$ to ensure a consistent truncation on $M \times S^1$.

Employing this procedure for $M = S^3 \times S^3$, gives us the consistent truncation on $S^3 \times S^3 \times S^1$. However, this does not have any AdS_3 vacua, since we still need to add the 7-form flux. To do this, let us first review the group theory of the $S^3 \times S^3$ truncation, since this will allow us to determine whether the 7-form flux is stabilised by the twist matrix.

5.2.3 Adding flux

To add a 7-form flux to the above truncation, we need to determine whether the 7-form flux is stabilised by the twist matrix on $S^3 \times S^3 \times S^1$ constructed in section 5.2.2. Thus, we need to understand whether IIB supergravity admits a 7-form field strength that is a singlet under the two $\text{SL}(4)$ groups in (5.25). To answer this question, we pick a gauge where the 6-form potential lives entirely in $S^3 \times S^3$ but depends on the S^1 coordinate, z . Therefore, in this gauge choice, the 6-form potential corresponds to an adjoint generator of $E_{7(7)}$, and since the S^1 coordinate is an $E_{7(7)}$ singlet, this adjoint generator must be a singlet under $\text{SL}(4)_1 \times \text{SL}(4)_2 \subset \text{SL}(8) \subset E_{7(7)}$ for us to have a consistent truncation.

IIB supergravity contains an S-duality doublet of 6-forms, which we can easily identify using the decomposition $E_{7(7)} \rightarrow \text{SL}(6) \times \text{SL}(2) \times \mathbb{R}_{\text{IB}}^+$, under which the $E_{7(7)}$ adjoint decomposes as

$$\begin{aligned} \mathbf{133} \rightarrow & (\mathbf{35}, \mathbf{1})_0 \oplus (\mathbf{1}, \mathbf{3})_0 \oplus (\mathbf{1}, \mathbf{1})_0 \oplus (\mathbf{1}, \mathbf{2})_{\pm 6} \\ & \oplus (\overline{\mathbf{15}}, \mathbf{2})_2 \oplus (\overline{\mathbf{15}}, \mathbf{1})_{-4} \oplus (\mathbf{15}, \mathbf{2})_{-2} \oplus (\mathbf{15}, \mathbf{1})_4. \end{aligned} \quad (5.34)$$

The 6-form doublet corresponds to the $(\mathbf{1}, \mathbf{2})_6$. To understand if one of the 6-forms in the doublet are singlets under $\text{SL}(4)_1 \times \text{SL}(4)_2 \subset E_{7(7)}$, we must decompose $E_{7(7)}$ with respect to the common subgroup of $\text{SL}(4)_1 \times \text{SL}(4)_2$ and $\text{SL}(6) \times \text{SL}(2)$, which is $\text{SL}(3)_1 \times \text{SL}(3)_2 \times \mathbb{R}_1^+ \times \mathbb{R}_2^+$, as defined by

the embedding of the physical coordinates in (5.28). Important for us is the correct identification of the \mathbb{R}^+ charges in both decompositions. We have, on the one hand,

$$\begin{aligned} E_{7(7)} &\rightarrow \mathrm{SL}(8) \rightarrow \mathrm{SL}(4) \times \mathrm{SL}(4) \times \mathbb{R}_{\mathrm{SL}(8)}^+ \\ &\rightarrow \mathrm{SL}(3)_1 \times \mathrm{SL}(3)_2 \times \mathbb{R}_1^+ \times \mathbb{R}_2^+ \times \mathbb{R}_{\mathrm{SL}(8)}^+, \end{aligned} \quad (5.35)$$

and, on the other,

$$\begin{aligned} E_{7(7)} &\rightarrow \mathrm{SL}(6) \times \mathrm{SL}(2) \times \mathbb{R}_{\mathrm{IIB}}^+ \\ &\rightarrow \mathrm{SL}(3)_1 \times \mathrm{SL}(3)_2 \times \mathbb{R}_{\mathrm{SL}(2)}^+ \times \mathbb{R}_{\mathrm{SL}(6)}^+ \times \mathbb{R}_{\mathrm{IIB}}^+. \end{aligned} \quad (5.36)$$

Here, we label the \mathbb{R}^+ 's with a subscript that refers to the groups they belong to. The \mathbb{R}^+ generators of the two decompositions are related as

$$\begin{aligned} \mathbb{R}_{\mathrm{SL}(6)}^+ &= -\frac{1}{2} \left(\mathbb{R}_{\mathrm{SL}(3)_1}^+ + \mathbb{R}_{\mathrm{SL}(3)_2}^+ \right), \\ \mathbb{R}_{\mathrm{SL}(2)}^+ &= \frac{1}{4} \left(\mathbb{R}_{\mathrm{SL}(3)_1}^+ - \mathbb{R}_{\mathrm{SL}(3)_2}^+ + \mathbb{R}_{\mathrm{SL}(8)}^+ \right), \\ \mathbb{R}_{\mathrm{IIB}}^+ &= \frac{1}{2} \left(\mathbb{R}_{\mathrm{SL}(3)_1}^+ - \mathbb{R}_{\mathrm{SL}(3)_2}^+ - 3 \mathbb{R}_{\mathrm{SL}(8)}^+ \right). \end{aligned} \quad (5.37)$$

The branching of the adjoint (5.34) under $\mathrm{SL}(6) \times \mathrm{SL}(2) \times \mathbb{R}_{\mathrm{IIB}}^+$ in (5.36) needs to now be compared with that via $\mathrm{SL}(4)_1 \times \mathrm{SL}(4)_2 \times \mathbb{R}_{\mathrm{SL}(8)}^+$, given by

$$\begin{aligned} \mathbf{133} &\rightarrow \mathbf{63} \oplus \mathbf{70} \\ &\rightarrow (\mathbf{15}, \mathbf{1})_0 \oplus (\mathbf{1}, \mathbf{15})_0 \oplus (\mathbf{1}, \mathbf{1})_0 \oplus (\mathbf{4}, \overline{\mathbf{4}})_2 \oplus (\overline{\mathbf{4}}, \mathbf{4})_{-2} \\ &\quad \oplus (\mathbf{1}, \mathbf{1})_{-4} \oplus (\mathbf{1}, \mathbf{1})_4 \oplus (\mathbf{4}, \overline{\mathbf{4}})_{-2} \oplus (\overline{\mathbf{4}}, \mathbf{4})_2 \oplus (\mathbf{6}, \mathbf{6})_0. \end{aligned} \quad (5.38)$$

Using (5.37), we can now identify each of the singlet generators $(1, 1)_{\pm 4}$ with one of the $\mathrm{SL}(2)$ doublet generators of charge ∓ 6 . Therefore, we find exactly one singlet $\mathrm{SL}(4)_1 \times \mathrm{SL}(4)_2$ generator, which we can write as t_{1238} , corresponding to a 6-form potential. The other $\mathrm{SL}(4)_1 \times \mathrm{SL}(4)_2$ singlet generator, t_{4567} , only differs by a compact generator and thus does not correspond to a different physical field. Finally, the other elements of the $\mathrm{SL}(2)$ doublet of 6-form potential can be mapped to the $(\mathbf{4}, \overline{\mathbf{4}})_2 \oplus (\overline{\mathbf{4}}, \mathbf{4})_{-2}$ generators, specifically t_7^8 and t_8^7 and are clearly not $\mathrm{SL}(4)$ singlets.

Therefore, we can add a 7-form flux to the twist matrix using

$$\mathcal{U}_{\overline{M}}^M \longrightarrow \mathcal{U}'_{\overline{M}}^M = \mathcal{U}_{\overline{M}}^N \exp(C)_N^M, \quad (5.39)$$

with $C_M^N = \lambda z \rho^{-1} t_{1238} M^N$ and λ a numerical parameter corresponding to the amount of 7-form flux. The resulting embedding tensor is most nicely written using the branching

$$\begin{aligned} E_{8(8)} &\longrightarrow E_{7(7)} \times \mathrm{SL}(2) \longrightarrow \mathrm{SL}(8) \times \mathrm{SL}(2) \\ \mathbf{248} &\longrightarrow (\mathbf{63}, \mathbf{1}) \oplus (\mathbf{70}, \mathbf{1}) \oplus (\mathbf{28}, \mathbf{2}) \oplus (\overline{\mathbf{28}}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{3}) \\ X^{\overline{M}} &\longrightarrow \left\{ X^{\overline{\mathcal{A}}}_{\overline{B}}, X^{\overline{\mathcal{A}BCD}}, X^{\overline{\mathcal{A}B}i}, X^{\overline{\mathcal{A}B}}_{\overline{i}}, X^{\overline{\alpha}} \right\}, \end{aligned} \quad (5.40)$$

and has the following non-vanishing components under the $\text{SL}(4)_1 \times \text{SL}(4)_2$ subgroups of $\text{SL}(8)$:

$$\begin{aligned}
X_{(\mathbf{63}, \mathbf{1}); (\mathbf{28}, \mathbf{2})} : & \begin{cases} X_{\bar{I}^{\bar{J}}; \bar{K}\bar{L}+} = -\frac{1}{\alpha} \delta_{\bar{I}[\bar{K}} \delta_{\bar{L}]}^{\bar{J}}, \\ X_{\bar{A}^{\bar{I}}; \bar{B}\bar{J}+} = -\frac{1}{2\alpha} \delta_{\bar{A}\bar{B}} \delta_{\bar{J}}^{\bar{I}}, \end{cases} \\
X_{(\mathbf{63}, \mathbf{1}); (\mathbf{28}, \mathbf{2})} : & \begin{cases} X_{\bar{A}^{\bar{B}}; \bar{C}\bar{D}+} = -\delta_{\bar{A}}^{[\bar{C}} \delta_{\bar{D}]}^{\bar{B}}, \\ X_{\bar{A}^{\bar{I}}; \bar{B}\bar{J}+} = -\frac{1}{2} \delta_{\bar{A}}^{\bar{B}} \delta_{\bar{J}}^{\bar{I}}, \end{cases} \\
X_{(\mathbf{70}, \mathbf{1}); (\mathbf{1}, \mathbf{3})} : & X_{\bar{I}\bar{J}\bar{K}\bar{L}; ++} = -\frac{\lambda}{12\sqrt{6}} \varepsilon_{\bar{I}\bar{J}\bar{K}\bar{L}}.
\end{aligned} \tag{5.41}$$

With this embedding tensor, the potential (4.14) has a vacuum at the scalar origin when $\lambda = 4\sqrt{3} \sqrt{1 + (1/\alpha^2)}$. We choose this value of λ from here onwards. **Presumably condition is on λ^2 and the sign of λ is fixed by SUSY (which we can ignore)?** (Exactly, the sign of λ exchanges \mathcal{N}_L and \mathcal{N}_R in $\mathcal{N} = (\mathcal{N}_L, \mathcal{N}_R)$)

6 Moduli of $\text{AdS}_3 \times S^3 \times S^3 \times S^1$

6.1 The undeformed $\mathcal{N} = (4, 4)$ $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ solution

At $\lambda = 4\sqrt{3} \sqrt{1 + (1/\alpha^2)}$, the three-dimensional solution described by the embedding tensor (5.41) corresponds to the consistent truncation of the pure NSNS ten-dimensional solution [42]

$$\begin{aligned}
ds^2 &= \ell_{\text{AdS}}^2 ds^2(\text{AdS}_3) + \alpha^2 ds^2(S^3) + ds^2(\tilde{S}^3) + dz^2, \\
H_{(3)} &= 2 \left(\ell_{\text{AdS}}^2 \text{Vol}(\text{AdS}_3) + \alpha^2 \text{Vol}(S^3) + \text{Vol}(\tilde{S}^3) \right),
\end{aligned} \tag{6.1}$$

with $H_{(3)} = dB_{(2)}$ and

$$\ell_{\text{AdS}}^2 = \frac{\alpha^2}{1 + \alpha^2}. \tag{6.2}$$

The S^1 coordinate z has periodicity $z \rightarrow z + 2\pi$.

The AdS_3 solution (6.1) preserves $\mathcal{N} = (4, 4)$ supersymmetry, with super-isometry group $\mathcal{G} = D^1(2, 1; \alpha)_L \times D^1(2, 1; \alpha)_R$, *i.e.* two copies of the large $\mathcal{N} = 4$ supergroup. The even part of $D^1(2, 1; \alpha)$ is isomorphic to $\text{SL}(2, \mathbb{R}) \times \text{SO}(3) \times \widetilde{\text{SO}(3)}$ and the bosonic isometries of the $\text{AdS}_3 \times S^3 \times \tilde{S}^3$ background are built from the even part of \mathcal{G} . The AdS_3 isometry group is $\text{SO}(2, 2) \simeq \text{SL}(2, \mathbb{R})_L \times \text{SL}(2, \mathbb{R})_R$, and the factors $\text{SO}(3)_L \times \text{SO}(3)_R \times \widetilde{\text{SO}(3)}_L \times \widetilde{\text{SO}(3)}_R \simeq \text{SO}(4) \times \text{SO}(4)$ combine into the isometry groups of the two spheres. The spectrum organizes into multiplets of \mathcal{G} , made out of products of long representations $[\ell, \tilde{\ell}]$ of $D^1(2, 1; \alpha)$ (see app. A for a review of the representations of $D^1(2, 1; \alpha)$). Explicitly, the spectrum is [43]

$$\bigoplus_{\ell, \tilde{\ell} \geq 0} [\ell, \tilde{\ell}] \otimes [\ell, \tilde{\ell}] \ominus [0, 0]_s \otimes [0, 0]_s. \tag{6.3}$$

In each factor the lowest conformal dimension is

$$h_L = h_R = -\frac{1}{2} + \frac{1}{2} \sqrt{1 + \frac{4\ell(\ell+1) + 4\alpha^2 \tilde{\ell}(\tilde{\ell}+1) + (2\pi n)^2}{1 + \alpha^2}}, \quad (6.4)$$

with n corresponding to the S^1 harmonics, and the lowest conformal dimension of $[\ell, \tilde{\ell}] \otimes [\ell, \tilde{\ell}]$ is then

$$\Delta_{\ell, \tilde{\ell}, n} = h_L + h_R = -1 + \sqrt{1 + \frac{4\ell(\ell+1) + 4\alpha^2 \tilde{\ell}(\tilde{\ell}+1) + (2\pi n)^2}{1 + \alpha^2}}. \quad (6.5)$$

For $n = 0$, the representations get shortened whenever $\ell = \tilde{\ell}$. The spectrum at the consistent truncation is then the one of ref. [27]. It corresponds to $\ell = \tilde{\ell} = n = 0$ in eq. (6.3):

$$[0, 0]_s \otimes [1/2, 1/2]_s \oplus [1/2, 1/2]_s \otimes [0, 0]_s \oplus [1/2, 1/2]_s \otimes [1/2, 1/2]_s. \quad (6.6)$$

The first two terms build the supergravity multiplet, each of them carrying one spin-2 field, four gravitini, seven vectors and four spin-1/2 fields, all massless. The only propagating degrees of freedom are in $[1/2, 1/2]_s \otimes [1/2, 1/2]_s$ (see the details in table 4 of [27]).

6.2 Moduli in gauged supergravity

Surprisingly, within the 3-dimensional gauged supergravity, the AdS_3 vacuum contains a large moduli space, which even has supersymmetric submanifolds. In particular, without any attempts at being exhaustive (**too strong? does this make us sound lazy?**) (**I think it's fine.**), we identified a continuous 11-parameter family of AdS_3 solutions, connected to the origin and given by the $E_{8(8)}$ coset representative (**$\overline{28} \leftrightarrow 28$ compared to the code, I think we made a mistake and confused down and up indices**)

$$\begin{aligned} \mathcal{V}_{\text{def}} = & \exp \left[\sqrt{2} \left(\chi_1(t_{(\mathbf{28}, \mathbf{2})})_{23, -} + \chi_2(t_{(\mathbf{28}, \mathbf{2})})_{81, -} + \tilde{\chi}_1(t_{(\overline{\mathbf{28}}, \mathbf{2})})_{56, -} + \tilde{\chi}_2(t_{(\overline{\mathbf{28}}, \mathbf{2})})_{74, -} \right) \right] \\ & \times \exp \left[\sqrt{2} \left(\zeta_1(t_{(\overline{\mathbf{28}}, \mathbf{2})})_{23, +} + \zeta_2(t_{(\overline{\mathbf{28}}, \mathbf{2})})_{81, +} + \tilde{\zeta}_1(t_{(\mathbf{28}, \mathbf{2})})_{56, +} + \tilde{\zeta}_2(t_{(\mathbf{28}, \mathbf{2})})_{74, +} \right) \right] \\ & \times \exp \left[\frac{\Omega}{120} \left((t_{(\mathbf{63}, \mathbf{1})})_1^1 - (t_{(\mathbf{63}, \mathbf{1})})_2^2 - (t_{(\mathbf{63}, \mathbf{1})})_3^3 + (t_{(\mathbf{63}, \mathbf{1})})_8^8 \right) \right. \\ & \quad \left. + \frac{\tilde{\Omega}}{120} \left((t_{(\mathbf{63}, \mathbf{1})})_4^4 - (t_{(\mathbf{63}, \mathbf{1})})_5^5 - (t_{(\mathbf{63}, \mathbf{1})})_6^6 + (t_{(\mathbf{63}, \mathbf{1})})_7^7 \right) \right] \\ & \times \exp \left[\psi \left(\frac{1}{120} (t_{(\mathbf{63}, \mathbf{1})})_I^I - \frac{1}{120} (t_{(\mathbf{63}, \mathbf{1})})_A^A - \frac{1}{2} (t_{(\mathbf{1}, \mathbf{3})})_1 \right) \right], \end{aligned} \quad (6.7)$$

where we denoted the generators according to the decomposition (5.40). The $(\chi_i, \tilde{\chi}_i, \zeta_i, \tilde{\zeta}_i)$ deformations excite the $(\mathbf{6}, \mathbf{1})_2$ and $(\mathbf{1}, \mathbf{6})_2$ generators in the branching (5.26). (**Describe what we do for $T = \exp \psi$? I think there is some freedom in the coeff there.**) (**Describe what we do for Ω ? Check coeff with code. Possible to change the signs to $(+, +, -, -)$, but very long to compute uplift.**)

Kinetic terms? Might be interesting for the S-fold deformations to see what kind of metric we have on the moduli space.

$$\begin{aligned}
ds_{\text{Zam.}}^2 = & -\frac{1}{2} \left[e^{2\psi} \left(e^{-\Omega} d\zeta_1^2 + e^{\Omega} d\zeta_2^2 + e^{\tilde{\Omega}} d\tilde{\zeta}_1^2 + e^{-\tilde{\Omega}} d\tilde{\zeta}_2^2 \right) \right. \\
& + \left(e^{-2\psi-\tilde{\Omega}} + \tilde{\zeta}_1^2 + e^{-\Omega-\tilde{\Omega}} \zeta_1^2 \right) \left(e^{\Omega+\tilde{\Omega}} + e^{2\psi+\Omega} \tilde{\zeta}_2^2 + e^{2\psi+\tilde{\Omega}} \zeta_2^2 \right) d\chi_1^2 \\
& + \left(e^{-2\psi-\Omega} + e^{-\Omega-\tilde{\Omega}} \tilde{\zeta}_2^2 + \zeta_2^2 \right) \left(1 + e^{2\psi+\tilde{\Omega}} \tilde{\zeta}_1^2 + e^{2\psi+\Omega} \zeta_1^2 \right) d\chi_2^2 \\
& + \left(e^{-2\psi-\tilde{\Omega}} + \tilde{\zeta}_1^2 + e^{-\Omega-\tilde{\Omega}} \zeta_1^2 \right) \left(1 + e^{2\psi+\tilde{\Omega}} \tilde{\zeta}_2^2 + e^{2\psi+\Omega} \zeta_2^2 \right) d\tilde{\chi}_1^2 \\
& \left. + \left(e^{-2\psi-\Omega} + e^{-\Omega-\tilde{\Omega}} \tilde{\zeta}_2^2 + \zeta_2^2 \right) \left(e^{\Omega+\tilde{\Omega}} + e^{2\psi+\Omega} \tilde{\zeta}_2^2 + e^{2\psi+\tilde{\Omega}} \zeta_1^2 \right) d\tilde{\chi}_2^2 + \dots \right]
\end{aligned} \tag{6.8}$$

6.3 Moduli in 10 dimensions

The consistent truncation Ansatz we constructed in this paper can be used to uplift the full 11-parameter family of AdS_3 vacua of the 3-dimensional gauged supergravity in (6.7) to IIB supergravity. However, this full 11-parameter solution can quickly become unwieldy. Therefore, here we will instead focus on a few interesting subsets of deformations.

$(\chi, \tilde{\chi}, \psi)$ Using Hopf coordinates for both S^3 's we can write the 5-parameter deformation involving $(\chi_i, \tilde{\chi}_i, \psi)$, $i = 1, 2$, as follows: **(Are those Hopf coordinates?)**

$$\begin{aligned}
ds^2 = & \ell_{\text{AdS}}^2 ds^2(\text{AdS}_3) + \alpha^2 (d\theta^2 + \cos^2(\theta) d\phi_1'^2 + \sin^2(\theta) d\phi_2'^2) \\
& + d\tilde{\theta}^2 + \cos^2(\tilde{\theta}) d\tilde{\phi}_1'^2 + \sin^2(\tilde{\theta}) d\tilde{\phi}_2'^2 + e^{2\psi} dz^2,
\end{aligned} \tag{6.9}$$

$$H_{(3)} = 2 \ell_{\text{AdS}}^2 \text{Vol}(\text{AdS}_3) + 2 \alpha^2 \cos(\theta) \sin(\theta) d\theta \wedge d\phi_1' \wedge d\phi_2' + 2 \cos(\tilde{\theta}) \sin(\tilde{\theta}) d\tilde{\theta} \wedge d\tilde{\phi}_1' \wedge d\tilde{\phi}_2',$$

where **Can we double check the α 's? I'm slightly confused (I think it's correct.)**

$$\begin{cases} d\phi_1' = d\phi_1 + \frac{1}{\alpha} \chi_1 dz, \\ d\phi_2' = d\phi_2 + \frac{1}{\alpha} \chi_2 dz, \end{cases} \quad \begin{cases} d\tilde{\phi}_1' = d\tilde{\phi}_1 + \tilde{\chi}_1 dz, \\ d\tilde{\phi}_2' = d\tilde{\phi}_2 + \tilde{\chi}_2 dz. \end{cases} \tag{6.10}$$

This 5-parameter deformation generically breaks $\text{SO}(3)_L \times \text{SO}(3)_R \times \widetilde{\text{SO}(3)}_L \times \widetilde{\text{SO}(3)}_R$ to $\text{U}(1)_L \times \text{U}(1)_R \times \widetilde{\text{U}(1)}_L \times \widetilde{\text{U}(1)}_R$ and all supersymmetries. The $\chi_i, \tilde{\chi}_i$, $i = 1, 2$ deformations are analogous to the two-parameter “flat deformations” [29] of $\text{AdS}_4 \times S^5 \times S^1$ [30, 31]. From the 4-dimensional perspective obtained after reducing on $S^3 \times S^3$, these deformations correspond to turning on Wilson lines for the $\text{SO}(3)_L \times \text{SO}(3)_R \times \widetilde{\text{SO}(3)}_L \times \widetilde{\text{SO}(3)}_R$ gauge fields along S^1 . The ψ deformation is even simpler: it just rescales the S^1 radius. These $\chi_i, \tilde{\chi}_i, \psi$ deformations are completely analogous to \mathbb{C} -structure deformations of T^2 , with the $\chi_i, \tilde{\chi}_i$ corresponding to the real part and the ψ deformation to the imaginary part of the \mathbb{C} -structure deformation.

Therefore, these deformations affect the Kaluza-Klein spectrum as follows: the conformal dimension of each physical field (*c.f.* eq. (6.5)) get shifted by replacing

$$2\pi n \longrightarrow e^{-2\psi} \left[2\pi n + \left((q_L + q_R) \chi_1 + (q_L - q_R) \chi_2 + \alpha (\tilde{q}_L + \tilde{q}_R) \tilde{\chi}_1 + \alpha (\tilde{q}_R - \tilde{q}_L) \tilde{\chi}_2 \right) \right], \tag{6.11}$$

where the q denote the charges under the different $\text{U}(1)$. **Somewhere here, we should give the character for all multiplets.**

Eq. (6.11) can be used to search for SUSY enhancement points within the 5-parameter landscape. At the undeformed origin, the massless gravitini sit in the two first multiplets in eq. (6.6) with $\text{SO}(3)_L \times \widetilde{\text{SO}}(3)_L \times \text{SO}(3)_R \times \widetilde{\text{SO}}(3)_R$ spins

$$(1/2, 1/2; 0, 0) \quad \text{and} \quad (0, 0; 1/2, 1/2). \quad (6.12)$$

and conformal dimension $\Delta = 3/2$. Turning on the deformation leads to a split into four groups of two modes, with the following $\text{U}(1)_L \times \widetilde{\text{U}}(1)_L \times \text{U}(1)_R \times \widetilde{\text{U}}(1)_R$ charges and conformal dimensions:

$$\begin{aligned} (\pm 1, \pm 1; 0, 0) : \quad \Delta &= \frac{1}{2} + \sqrt{1 + \frac{(\chi_1 + \chi_2 + \alpha(\tilde{\chi}_1 - \tilde{\chi}_2))^2}{4e^{2\psi}(1 + \alpha^2)}}, \\ (\pm 1, \mp 1; 0, 0) : \quad \Delta &= \frac{1}{2} + \sqrt{1 + \frac{(\chi_1 + \chi_2 - \alpha(\tilde{\chi}_1 - \tilde{\chi}_2))^2}{4e^{2\psi}(1 + \alpha^2)}}, \\ (0, 0; \pm 1, \pm 1) : \quad \Delta &= \frac{1}{2} + \sqrt{1 + \frac{(\chi_1 - \chi_2 + \alpha(\tilde{\chi}_1 + \tilde{\chi}_2))^2}{4e^{2\psi}(1 + \alpha^2)}}, \\ (0, 0; \pm 1, \mp 1) : \quad \Delta &= \frac{1}{2} + \sqrt{1 + \frac{(\chi_1 - \chi_2 - \alpha(\tilde{\chi}_1 + \tilde{\chi}_2))^2}{4e^{2\psi}(1 + \alpha^2)}}. \end{aligned} \quad (6.13)$$

SUSY enhancement points are then given by combinations of the parameters that yield $\Delta = 3/2$. This can either happen by leaving some modes with $\Delta = 3/2$ invariant, or where some other, originally massive gravitini outside the consistent truncation, i.e. with conformal dimension given by (6.5) with $n \neq 0$, obtain $\Delta = 3/2$. The different possibilities are listed below. (Shall we include the n ? Shall we give the multiplets? At least for the truncation? Not sure it's worth it.)

- $\mathcal{N} = 2$ The four-dimensional hypersurfaces

$$\chi_2 = \chi_1 \pm \alpha(\tilde{\chi}_1 + \tilde{\chi}_2) \quad \text{or} \quad \chi_2 = -\chi_1 \pm \alpha(\tilde{\chi}_1 - \tilde{\chi}_2), \quad (6.14)$$

give enhancements to $\mathcal{N} = (2, 0)$ and $\mathcal{N} = (0, 2)$, respectively. On this hypersurface (6.14), the spectrum then organizes into multiplets of $\text{SU}(1|1, 1)$.

- $\mathcal{N} = 4$ Choosing

$$\begin{cases} \chi_2 = \chi_1, \\ \tilde{\chi}_2 = -\tilde{\chi}_1, \end{cases} \quad \text{or} \quad \begin{cases} \chi_2 = -\chi_1, \\ \tilde{\chi}_2 = \tilde{\chi}_1, \end{cases} \quad (6.15)$$

leads to enhancements to $\mathcal{N} = (4, 0)$ and $\mathcal{N} = (0, 4)$, respectively. On these three-dimensional hyper-surfaces (6.15), the isometry group is $\text{SO}(4) \times \text{U}(1)^2$ and the spectrum reorganizes into multiplets of $D^1(2, 1; \alpha)$.

(Mention here the SUSY family of [1].)

Another possibility is to set

$$\begin{cases} \tilde{\chi}_1 = \pm \chi_1 / \alpha, \\ \tilde{\chi}_2 = \mp \chi_2 / \alpha, \end{cases} \quad \text{or} \quad \begin{cases} \tilde{\chi}_1 = \pm \chi_2 / \alpha, \\ \tilde{\chi}_2 = \mp \chi_1 / \alpha, \end{cases} \quad (6.16)$$

giving 3-parameter families of $\mathcal{N} = (2, 2)$ solutions, with multiplets of $\text{SU}(1|1, 1)_L \times \text{SU}(1|1, 1)_R$.

- $\mathcal{N} = 6$ Finally, for

$$\begin{cases} \chi_2 = \chi_1, \\ \tilde{\chi}_1 = \pm \chi_1/\alpha, \\ \tilde{\chi}_2 = \mp \chi_1/\alpha, \end{cases} \quad \text{or} \quad \begin{cases} \chi_2 = -\chi_1, \\ \tilde{\chi}_1 = \pm \chi_1/\alpha, \\ \tilde{\chi}_2 = \mp \chi_1/\alpha, \end{cases} \quad (6.17)$$

there are enhancements to $\mathcal{N} = (4, 2)$ and $\mathcal{N} = (2, 4)$, respectively. The isometry group is $\text{SO}(4) \times \text{U}(1)^2$ and the spectrum is then given by multiplets of $D^1(2, 1; \alpha)_{\text{L,R}} \times \text{SU}(1|1, 1)_{\text{R,L}}$.

In the above cases, it is easy to see from (6.11) that the spectrum is invariant under $\chi_i \rightarrow \chi_i + 2\pi k_i$, $\tilde{\chi}_i \rightarrow \tilde{\chi}_i + 2\pi \tilde{k}_i/\alpha$, for $k_i, \tilde{k}_i \in \mathbb{Z}$, with the SUSY enhancements also occurring if the relations (6.14), (6.15), (6.16) or (6.17) are satisfied up to integer shifts. We will return to this point below in section 6.4, when discussing compactness of the moduli space.

For generic values of $\chi_i, \tilde{\chi}_i$, the AdS_3 vacuum will be completely non-supersymmetric. However, it is easy to see from inspection of the spectrum (6.5) and (6.11) that the spectrum is bounded from below by the masses of the 3-dimensional gauged supergravity modes at the $\mathcal{N} = (4, 4)$ supersymmetric origin. Therefore, these non-supersymmetric AdS_3 vacua are still perturbatively stable within IIB supergravity and, similarly to [31] suggest that there may be a dual non-supersymmetric conformal manifold if there are no instabilities beyond supergravity, such as non-perturbative instabilities triggered by brane nucleation or non-vanishing β -functions of marginal multi-trace operators [add citations](#). (Mention the continuous limits to SUSY vacua?)

$(\zeta, \tilde{\zeta}, \Omega, \tilde{\Omega}, \psi)$ Again with spherical coordinates:

$$\begin{aligned} ds^2 = & \ell_{\text{AdS}}^2 ds^2(\text{AdS}_3) - \Delta \left(\cos^2(\theta) + e^\Omega \sin^2(\theta) \right) \left(e^{\tilde{\Omega}} \cos^2(\tilde{\theta}) + \sin^2(\tilde{\theta}) \right) dz^2 \\ & + \alpha^2 \left[d\theta^2 + \Delta \left(e^{\tilde{\Omega}} \cos^2(\tilde{\theta}) + \sin^2(\tilde{\theta}) \right) \left(e^\Omega \cos^2(\theta) d\phi_1^2 + \sin^2(\theta) d\phi_2^2 \right) \right] \\ & + d\tilde{\theta}^2 + \Delta \left(\cos^2(\theta) + e^\Omega \sin^2(\theta) \right) \left(\cos^2(\tilde{\theta}) d\tilde{\phi}_1^2 + e^{\tilde{\Omega}} \sin^2(\tilde{\theta}) d\tilde{\phi}_2^2 \right) \\ & + \Delta e^{2\psi} e^\Omega \cos^2(\theta) \cos^2(\tilde{\theta}) \left(\alpha \tilde{\zeta}_2 d\phi_1 - \zeta_2 d\tilde{\phi}_1 \right)^2 + \Delta e^{2\psi} e^{\tilde{\Omega}} \sin^2(\theta) \sin^2(\tilde{\theta}) \left(\alpha \tilde{\zeta}_1 d\phi_2 - \zeta_1 d\tilde{\phi}_2 \right)^2 \\ & + \Delta e^{2\psi} e^{\Omega+\tilde{\Omega}} \cos^2(\theta) \sin^2(\tilde{\theta}) \left(\alpha \tilde{\zeta}_1 d\phi_1 - \zeta_2 d\tilde{\phi}_2 \right)^2 + \Delta e^{2\psi} \sin^2(\theta) \cos^2(\tilde{\theta}) \left(\alpha \tilde{\zeta}_2 d\phi_2 - \zeta_1 d\tilde{\phi}_1 \right)^2 \\ & - 2\alpha \Delta e^{2\psi} \left(\zeta_1 \cos^2(\theta) d\phi_1 + e^\Omega \zeta_2 \sin^2(\theta) d\phi_2 \right) \left(e^{\tilde{\Omega}} \tilde{\zeta}_1 \cos^2(\tilde{\theta}) d\tilde{\phi}_1 + \tilde{\zeta}_2 \sin^2(\tilde{\theta}) d\tilde{\phi}_2 \right) \\ & + \Delta e^{2\psi} \left(e^{\tilde{\Omega}} \cos^2(\tilde{\theta}) + \sin^2(\tilde{\theta}) \right) \left[\cos^2(\theta) (\alpha \zeta_1 d\phi_1 + dz)^2 + e^\Omega \sin^2(\theta) (\alpha \zeta_2 d\phi_2 + dz)^2 \right] \\ & + \Delta e^{2\psi} \left(\cos^2(\theta) + e^\Omega \sin^2(\theta) \right) \left[e^{\tilde{\Omega}} \cos^2(\tilde{\theta}) \left(\tilde{\zeta}_1 d\tilde{\phi}_1 - dz \right)^2 + \sin^2(\tilde{\theta}) \left(\tilde{\zeta}_2 d\tilde{\phi}_2 - dz \right)^2 \right], \end{aligned} \quad (6.18)$$

with the warp factor

$$\begin{aligned} \Delta^{-1} = & \left(\cos^2(\theta) + e^\Omega \sin^2(\theta) \right) \left(e^{\tilde{\Omega}} \cos^2(\tilde{\theta}) + \sin^2(\tilde{\theta}) \right) \\ & + e^{2\psi} \left(\cos^2(\theta) + e^\Omega \sin^2(\theta) \right) \left(\tilde{\zeta}_2^2 \cos^2(\tilde{\theta}) + e^{\tilde{\Omega}} \tilde{\zeta}_1^2 \sin^2(\tilde{\theta}) \right) \\ & + e^{2\psi} \left(e^{\tilde{\Omega}} \cos^2(\tilde{\theta}) + \sin^2(\tilde{\theta}) \right) \left(e^\Omega \zeta_2^2 \cos^2(\theta) + \zeta_1^2 \sin^2(\theta) \right). \end{aligned} \quad (6.19)$$

(Expression of H ?)

SUSY deformation It is convenient to use the Hopf fibration to describe this, we denote by ϕ the coordinate on the S^1 fibre and by ξ the homogeneous coordinate on \mathbb{CP}^1 . The round metric on S^3 then reads

$$g = ds_{\mathbb{CP}^1}^2 + (\sigma + d\phi)^2 \quad (6.20)$$

where

$$\begin{aligned} ds_{\mathbb{CP}^1}^2 &= \frac{1}{(1 + |\xi|^2)^2} d\xi d\bar{\xi} \\ \sigma &= \frac{i}{2(1 + |\xi|^2)} (\xi d\bar{\xi} - \bar{\xi} d\xi) \\ vol_{S^3} &= vol_{\mathbb{CP}^1} \wedge d\psi. \end{aligned} \quad (6.21)$$

The deformation does not affect S_+^3 , but it mixes the S^1 factor in the internal space with the Hopf fibre of S_-^3 . It is parametrised by a real number Ω , such that the metric reads

$$\begin{aligned} ds^2 &= \alpha^2 (ds_{\mathbb{CP}^1}^2 + \kappa^2) + ds_{S_+^3}^2 + e^{-4\Omega} d\psi^2 \\ \kappa &= d\phi + \sigma - \frac{1}{\alpha} e^{-2\Omega} \sqrt{e^{2\Omega} - 1} d\psi. \end{aligned} \quad (6.22)$$

The three form flux gets changed according to the prescription $d\phi \rightarrow d\phi - \frac{1}{\alpha} e^{-2\Omega} \sqrt{e^{2\Omega} - 1} d\psi$ and we find:

$$H_{(3)} = \alpha^2 vol_{\mathbb{CP}^1} \wedge (d\phi - \frac{1}{\alpha} e^{-2\Omega} \sqrt{e^{2\Omega} - 1} d\psi) + vol_{S_+^3}. \quad (6.23)$$

Non-SUSY deformation Here we study a more general deformation depending on three parameters Ω , $\zeta_{1,8}$ and $\zeta_{2,3}$. It is convenient to write down the uplift in terms of spherical coordinates on S_-^3 , while again S_+^3 is not deformed.

For the metric we find

$$\begin{aligned} ds^2 &= \ell_{\text{AdS}}^2 ds^2(\text{AdS}_3) + ds^2(S_+^3) \\ &+ \alpha^2 (d\theta^2 + \Delta (2e^{2\Omega} + \zeta_{2,3}^2) \cos^2(\theta) d\varphi_1^2 + \Delta (2 + e^{2\Omega} \zeta_{1,8}^2) \sin^2(\theta) d\varphi_2^2) \\ &+ 2\sqrt{2} \alpha \Delta (-\zeta_{2,3} \cos^2(\theta) d\varphi_1 + \zeta_{1,8} \sin^2(\theta) d\varphi_2) d\psi \\ &+ 2 \Delta (\cos^2(\theta) + e^{2\Omega} \sin^2(\theta)) d\psi^2, \end{aligned} \quad (6.24)$$

where Δ is the warp factor

$$\Delta^{-1} = 2e^{2\Omega} + \zeta_{2,3}^2 + \cos^2(\theta) (2 - 2e^{2\Omega} - \zeta_{2,3}^2 + e^{2\Omega} \zeta_{1,8}^2). \quad (6.25)$$

The three-form flux is

$$H_{(3)} = vol_{S_-^3} + \alpha^2 \sin(\theta) \cos(\theta) (2 + e^{2\Omega} \zeta_{1,8}^2) (2e^{2\Omega} + \zeta_{2,3}^2) \Delta^2 d\theta \wedge \kappa_1 \wedge \kappa_2, \quad (6.26)$$

with the two one forms $\kappa_{1,2}$ defined by

$$\begin{aligned} \kappa_1 &= d\phi_1 - \frac{\zeta_{2,3}}{\alpha(2e^{2\Omega} + \zeta_{2,3}^2)} d\psi \\ \kappa_2 &= d\phi_2 + \frac{\zeta_{1,8}}{\alpha(2 + e^{2\Omega} \zeta_{1,8}^2)} d\psi. \end{aligned} \quad (6.27)$$

Note that setting ... recovers the supersymmetric deformation described earlier.

6.4 Compactification of the moduli space

We can use the Kaluza-Klein spectrum and the uplift of the 3-dimensional gauged supergravity models to IIB supergravity to understand properties of the dual CFTs that are obscured in the lower-dimensional supergravities. One such aspect is the compactness of the conformal manifold. Consider the deformations $\chi_i, \tilde{\chi}_i$, which appear non-compact in 3-dimensional $\mathcal{N} = 16$ gauged supergravity, with only the point $\chi_i = \tilde{\chi}_i = 0$ having $\mathcal{N} = (4, 4)$ supersymmetry. The subset of these deformations corresponding to the SUSY lines $\chi_1 = \chi_2 = \chi, \tilde{\chi}_i = 0$, *Is this right?* was analysed in the half-maximal theory in [1]. Again, within 3-dimensional $\mathcal{N} = 8$ supergravity, this deformation is non-compact, and remains non-compact even after uplifting to 6-dimensional supergravity on S^3 , with no periodicity in χ observed in the KK spectrum of the 6-dimensional solution [1].

However, from (6.5), (6.11), (??) we see that the full Kaluza-Klein spectrum is invariant under the shifts

$$\begin{aligned}\chi_i &\rightarrow \chi_i + 2\pi k_i, \\ \tilde{\chi}_i &\rightarrow \tilde{\chi}_i + 2\pi \tilde{k}_i / \alpha,\end{aligned}\tag{6.28}$$

for $k_i, \tilde{k}_i \in \mathbb{Z}$. In these cases, higher KK modes on the S^1 outside the $\mathcal{N} = 16$ truncation replace those within the consistent truncation. Therefore, even though neither the 3-dimensional gauged supergravity, nor 6-dimensional supergravity is invariant under these shifts, upon lifting to the full 10-dimensional IIB theory, the higher KK modes on S^1 restore the invariance under the shifts, just as in the analogous deformations studied in $\text{AdS}_4 \times S^5 \times S^1$ [30, 31]. This highlights the importance of the S^1 direction, which is often ignored in holographic applications *Maybe remove this last sentence?. (Maybe the last part, but I would keep the first part.)*

The compactification of the moduli space in 10 dimensions can also be understood geometrically. Each of $\chi_i, \tilde{\chi}_i$ corresponds to the real part of the complex structure modulus of a T^2 fibration with local coordinates $\phi_i, \tilde{\phi}_i$ together with z , respectively. Explicitly, the complex structure is given by *Not sure about this, see comment about α above*

$$\tau = i \frac{e^\psi}{\alpha} + \frac{1}{\alpha} \chi.\tag{6.29}$$

Under this identification, the shifts (6.28) correspond to Dehn twists, or T -transformations, of the torus, which belong to the modular group $\text{SL}(2, \mathbb{Z})$, just as for $\text{AdS}_4 \times S^5 \times S^1$ [30, 31].

Given the interpretation of the shifts (6.28) as T -transformations of a T^2 -fibration, it is natural to ask whether S -transformations of these T^2 would also leave the $\text{AdS}_3 \times S^3 \times S^3 \times S^1$ solution invariant. The S -transformation acts on the moduli as

$$\psi \rightarrow -\psi,\tag{6.30}$$

thus inverting the radius of the S^1 . The KK spectrum (6.5), (6.11) is clearly not invariant under this transformation. However, the S -transformation takes us out of the regime of validity of supergravity, with new string degrees of freedom potentially becoming light. Indeed, in this case, after the transformation (6.30), strings wound around S^1 will become light, and one would expect them to replace the KK modes that have become heavy, realising the T-duality on S^1 . Thus, in the full IIB string theory, the moduli space of each T^2 fibration with complex structure (6.29) should be identified under the Möbius group $\text{SL}(2, \mathbb{Z})$. *Can we check the T-dual explicitly? Maybe we need a different gauge choice for C_6 so that it does not depend on z . Can we describe the $T^4 \times S^1$ a bit better instead of as a collection of T^2 's?*

7 Conclusions

Acknowledgements

We are grateful to XXX for useful discussions and correspondence. CE is supported by the FWO-Vlaanderen through the project G006119N and by the Vrije Universiteit Brussel through the Strategic Research Program “High-Energy Physics”. MG and EM are supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) via the Emmy Noether program “Exploring the landscape of string theory flux vacua using exceptional field theory” (project number 426510644).

A Representations of $D^1(2, 1; \alpha)$

The supergroup $D^1(2, 1; \alpha)$ has

$$\mathrm{SL}(2, \mathbb{R}) \times \mathrm{SU}(2) \times \widetilde{\mathrm{SU}(2)} \quad (\mathrm{A.1})$$

as even part. Its multiplets are labelled by the spins ℓ and $\tilde{\ell}$ of each $\mathrm{SU}(2)$ factor and the eigenvalue h of the $\mathfrak{sl}(2, \mathbb{R})$ generator. Generic short multiplets $[\ell, \tilde{\ell}]_s$ are given by [42, 43]

$$\begin{array}{ll} h_0 & (\ell, \tilde{\ell}) \\ h_0 + 1/2 & (\ell + 1/2, \tilde{\ell} - 1/2) \oplus (\ell - 1/2, \tilde{\ell} - 1/2) \oplus (\ell - 1/2, \tilde{\ell} + 1/2) \\ h_0 + 1 & (\ell, \tilde{\ell} - 1) \oplus (\ell, \tilde{\ell}) \oplus (\ell - 1, \tilde{\ell}) \\ h_0 + 3/2 & (\ell - 1/2, \tilde{\ell} - 1/2), \end{array} \quad (\mathrm{A.2})$$

with $h_0 = (\alpha \ell + \tilde{\ell})/(1 + \alpha)$. Shortenings occur for $\ell < 1$ or $\tilde{\ell} < 1$ [42]. For instance:

$$\begin{array}{llll} & h_0 = 1/2 & (1/2, 1/2) \\ [0, 0]_s : & h_0 = 0 & (0, 0), & [1/2, 1/2]_s : & 1 & (1, 0) \oplus (0, 0) \oplus (0, 1) \\ & & & & 3/2 & (1/2, 1/2) \\ & & & & 2 & (0, 0). \end{array} \quad (\mathrm{A.3})$$

Two short multiplets can combine into a long multiplets $[\ell, \tilde{\ell}]$ as follows:

$$[\ell, \tilde{\ell}] = [\ell, \tilde{\ell}]_s \oplus [\ell + 1/2, \tilde{\ell} + 1/2]_s. \quad (\mathrm{A.4})$$

The value of h is then not constrained. The explicit content of the long representation $[\ell, \tilde{\ell}]$ is

$$\begin{array}{ll} h & (\ell, \tilde{\ell}) \\ h + 1/2 & (\ell + 1/2, \tilde{\ell} + 1/2) \oplus (\ell + 1/2, \tilde{\ell} - 1/2) \oplus (\ell - 1/2, \tilde{\ell} + 1/2) \oplus (\ell - 1/2, \tilde{\ell} - 1/2) \\ h + 1 & (\ell + 1, \tilde{\ell}) \oplus (\ell, \tilde{\ell} - 1) \oplus (\ell, \tilde{\ell}) \oplus (\ell, \tilde{\ell}) \oplus (\ell - 1, \tilde{\ell}) \oplus (\ell, \tilde{\ell} + 1) \\ h + 3/2 & (\ell + 1/2, \tilde{\ell} + 1/2) \oplus (\ell + 1/2, \tilde{\ell} - 1/2) \oplus (\ell - 1/2, \tilde{\ell} + 1/2) \oplus (\ell - 1/2, \tilde{\ell} - 1/2) \\ h + 2 & (\ell, \tilde{\ell}). \end{array} \quad (\mathrm{A.5})$$

References

- [1] C. Eloy, G. Larios, and H. Samtleben, *Triality and the consistent reductions on $AdS_3 \times S^3$* , *JHEP* **01** (2022) 055, [[arXiv:2111.01167](#)].
- [2] O. Hohm and H. Samtleben, *Consistent Kaluza-Klein Truncations via Exceptional Field Theory*, *JHEP* **01** (2015) 131, [[arXiv:1410.8145](#)].
- [3] K. Lee, C. Strickland-Constable, and D. Waldram, *Spheres, generalised parallelisability and consistent truncations*, *Fortsch. Phys.* **65** (2017), no. 10-11 1700048, [[arXiv:1401.3360](#)].
- [4] E. Malek and H. Samtleben, *Dualising consistent IIA/IIB truncations*, *JHEP* **12** (2015) 029, [[arXiv:1510.03433](#)].
- [5] A. Baguet, O. Hohm, and H. Samtleben, *Consistent Type IIB Reductions to Maximal 5D Supergravity*, *Phys. Rev.* **D92** (2015), no. 6 065004, [[arXiv:1506.01385](#)].
- [6] A. Baguet, C. N. Pope, and H. Samtleben, *Consistent Pauli reduction on group manifolds*, *Phys. Lett.* **B752** (2016) 278–284, [[arXiv:1510.08926](#)].
- [7] K. Lee, C. Strickland-Constable, and D. Waldram, *New gaugings and non-geometry*, [arXiv:1506.03457](#).
- [8] E. Malek, *7-dimensional $\mathcal{N} = 2$ Consistent Truncations using $SL(5)$ Exceptional Field Theory*, *JHEP* **06** (2017) 026, [[arXiv:1612.01692](#)].
- [9] F. Ciceri, A. Guarino, and G. Inverso, *The exceptional story of massive IIA supergravity*, *JHEP* **08** (2016) 154, [[arXiv:1604.08602](#)].
- [10] D. Cassani, O. de Felice, M. Petrini, C. Strickland-Constable, and D. Waldram, *Exceptional generalised geometry for massive IIA and consistent reductions*, *JHEP* **08** (2016) 074, [[arXiv:1605.00563](#)].
- [11] G. Inverso, H. Samtleben, and M. Trigiante, *Type II supergravity origin of dyonic gaugings*, *Phys. Rev.* **D95** (2017), no. 6 066020, [[arXiv:1612.05123](#)].
- [12] E. Malek and H. Samtleben, *Ten-dimensional origin of Minkowski vacua in $N=8$ supergravity*, *Phys. Lett.* **B776** (2018) 64–71, [[arXiv:1710.02163](#)].
- [13] E. Malek, *Half-Maximal Supersymmetry from Exceptional Field Theory*, *Fortsch. Phys.* **65** (2017), no. 10-11 1700061, [[arXiv:1707.00714](#)].
- [14] E. Malek, H. Samtleben, and V. Vall Camell, *Supersymmetric AdS_7 and AdS_6 vacua and their minimal consistent truncations from exceptional field theory*, *Phys. Lett.* **B786** (2018) 171–179, [[arXiv:1808.05597](#)].
- [15] E. Malek, H. Samtleben, and V. Vall Camell, *Supersymmetric AdS_7 and AdS_6 vacua and their consistent truncations with vector multiplets*, *JHEP* **04** (2019) 088, [[arXiv:1901.11039](#)].
- [16] D. Cassani, G. Josse, M. Petrini, and D. Waldram, *Systematics of consistent truncations from generalised geometry*, *JHEP* **11** (2019) 017, [[arXiv:1907.06730](#)].

- [17] E. Malek and V. Vall Camell, *Consistent truncations around half-maximal AdS_5 vacua of 11-dimensional supergravity*, *Class. Quant. Grav.* **39** (2022), no. 7 075026, [[arXiv:2012.15601](#)].
- [18] D. Cassani, G. Josse, M. Petrini, and D. Waldram, *$\mathcal{N} = 2$ consistent truncations from wrapped M5-branes*, *JHEP* **02** (2021) 232, [[arXiv:2011.04775](#)].
- [19] M. Galli and E. Malek, *Consistent truncations to 3-dimensional supergravity*, *JHEP* **09** (2022) 014, [[arXiv:2206.03507](#)].
- [20] O. Hohm and H. Samtleben, *Exceptional Field Theory III: $E_{8(8)}$* , *Phys.Rev.* **D90** (2014) 066002, [[arXiv:1406.3348](#)].
- [21] M. Cederwall and J. A. Rosabal, *E_8 geometry*, *JHEP* **07** (2015) 007, [[arXiv:1504.04843](#)].
- [22] T. Fischbacher, H. Nicolai, and H. Samtleben, *Nonsemisimple and complex gaugings of $N=16$ supergravity*, *Commun. Math. Phys.* **249** (2004) 475–496, [[hep-th/0306276](#)].
- [23] J. M. Maldacena and H. Ooguri, *Strings in $AdS(3)$ and $SL(2,R)$ WZW model 1.: The Spectrum*, *J. Math. Phys.* **42** (2001) 2929–2960, [[hep-th/0001053](#)].
- [24] L. Eberhardt, M. R. Gaberdiel, and R. Gopakumar, *The Worldsheet Dual of the Symmetric Product CFT*, *JHEP* **04** (2019) 103, [[arXiv:1812.01007](#)].
- [25] L. Eberhardt and M. R. Gaberdiel, *Strings on $AdS_3 \times S^3 \times S^3 \times S^1$* , *JHEP* **06** (2019) 035, [[arXiv:1904.01585](#)].
- [26] L. Eberhardt, M. R. Gaberdiel, and R. Gopakumar, *Deriving the AdS_3/CFT_2 correspondence*, *JHEP* **02** (2020) 136, [[arXiv:1911.00378](#)].
- [27] O. Hohm and H. Samtleben, *Effective actions for massive Kaluza-Klein states on $AdS(3) \times S^{**3} \times S^{**3}$* , *JHEP* **05** (2005) 027, [[hep-th/0503088](#)].
- [28] A. Guarino, C. Sterckx, and M. Trigiante, *$\mathcal{N} = 2$ supersymmetric S-folds*, *JHEP* **04** (2020) 050, [[arXiv:2002.03692](#)].
- [29] A. Guarino and C. Sterckx, *Flat deformations of type IIB S-folds*, *JHEP* **11** (2021) 171, [[arXiv:2109.06032](#)].
- [30] A. Giambrone, E. Malek, H. Samtleben, and M. Trigiante, *Global properties of the conformal manifold for S-fold backgrounds*, *JHEP* **06** (2021), no. 111 111, [[arXiv:2103.10797](#)].
- [31] A. Giambrone, A. Guarino, E. Malek, H. Samtleben, C. Sterckx, and M. Trigiante, *Holographic evidence for nonsupersymmetric conformal manifolds*, *Phys. Rev. D* **105** (2022), no. 6 066018, [[arXiv:2112.11966](#)].
- [32] H. Nicolai and H. Samtleben, *Maximal gauged supergravity in three-dimensions*, *Phys.Rev.Lett.* **86** (2001) 1686–1689, [[hep-th/0010076](#)].
- [33] H. Nicolai and H. Samtleben, *Compact and noncompact gauged maximal supergravities in three-dimensions*, *JHEP* **04** (2001) 022, [[hep-th/0103032](#)].

- [34] G. Bossard, F. Ciceri, G. Inverso, and A. Kleinschmidt, *Consistent Kaluza-Klein Truncations and Two-Dimensional Gauged Supergravity*, *Phys. Rev. Lett.* **129** (2022), no. 20 201602, [[arXiv:2209.02729](#)].
- [35] G. Bossard, F. Ciceri, G. Inverso, A. Kleinschmidt, and H. Samtleben, *E_9 exceptional field theory. Part I. The potential*, *JHEP* **03** (2019) 089, [[arXiv:1811.04088](#)].
- [36] G. Bossard, F. Ciceri, G. Inverso, A. Kleinschmidt, and H. Samtleben, *E_9 exceptional field theory. Part II. The complete dynamics*, *JHEP* **05** (2021) 107, [[arXiv:2103.12118](#)].
- [37] D. S. Berman, E. T. Musaev, and D. C. Thompson, *Duality Invariant M-theory: Gauged supergravities and Scherk-Schwarz reductions*, *JHEP* **1210** (2012) 174, [[arXiv:1208.0020](#)].
- [38] E. T. Musaev, *Gauged supergravities in 5 and 6 dimensions from generalised Scherk-Schwarz reductions*, *JHEP* **1305** (2013) 161, [[arXiv:1301.0467](#)].
- [39] C. D. A. Blair and E. Malek, *Geometry and fluxes of $SL(5)$ exceptional field theory*, *JHEP* **1503** (2015) 144, [[arXiv:1412.0635](#)].
- [40] O. Hohm, E. T. Musaev, and H. Samtleben, *$O(d+1, d+1)$ enhanced double field theory*, *JHEP* **10** (2017) 086, [[arXiv:1707.06693](#)].
- [41] H. Samtleben and O. Sarioğlu, *Consistent S^3 reductions of six-dimensional supergravity*, *Phys. Rev. D* **100** (2019), no. 8 086002, [[arXiv:1907.08413](#)].
- [42] J. de Boer, A. Pasquinucci, and K. Skenderis, *AdS / CFT dualities involving large 2-D $N=4$ superconformal symmetry*, *Adv. Theor. Math. Phys.* **3** (1999) 577–614, [[hep-th/9904073](#)].
- [43] L. Eberhardt, M. R. Gaberdiel, R. Gopakumar, and W. Li, *BPS spectrum on $AdS_3 \times S^3 \times S^3 \times S^1$* , *JHEP* **03** (2017) 124, [[arXiv:1701.03552](#)].