

Homework 4

Caroline Hutchings

09/14/2026

Conceptual Questions

```
library(tidyverse) arrange(filter(select(as_tibble(iris), starts_with("Petal"), Species),  
Petal.Length < 1.55), Species)
```

1. Rewrite using baseR chaining operator

```
library(tidyverse)
```

```
-- Attaching core tidyverse packages ----- tidyverse 2.0.0 --  
v dplyr      1.2.0      v readr      2.1.6  
v forcats    1.0.1      v stringr    1.6.0  
v ggplot2    4.0.2      v tibble     3.3.1  
v lubridate  1.9.5      v tidyr      1.3.2  
v purrr      1.2.1
```

```
-- Conflicts ----- tidyverse_conflicts() --
```

```
x dplyr::filter() masks stats::filter()
```

```
x dplyr::lag()     masks stats::lag()
```

```
i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become
```

```
iris |>  
  as_tibble() |>  
  select(starts_with("Petal"), Species) |>  
  filter(Petal.Length<1.55) |>  
  arrange(Species)
```

```
# A tibble: 37 x 3
```

```
  Petal.Length Petal.Width Species  
      <dbl>         <dbl> <fct>
```

```

1          1.4          0.2 setosa
2          1.4          0.2 setosa
3          1.3          0.2 setosa
4          1.5          0.2 setosa
5          1.4          0.2 setosa
6          1.4          0.3 setosa
7          1.5          0.2 setosa
8          1.4          0.2 setosa
9          1.5          0.1 setosa
10         1.5          0.2 setosa
# i 27 more rows

```

2. What is meant by long format data and wide format data?

1. Long format is a data table organization where each row is one observation and each column is a variable. Repeated measures are stored in additional rows. This is the ideal format when working with structured data for statistical analysis.
2. Wide format is where there can be multiple observations per row. Repeated measures are stored in additional columns instead of additional rows. Sometimes this makes the data easier to read in a table format.
3. Which do we generally prefer for statistical analysis? We generally prefer the long format to do analyses on. This enables us to do analysis on the variables and include each observation separately, not necessarily each individual though as some rows may be repeated measures on the same individual.

Task 2

Glass Data

Read in the glass data to a tibble. Change the variable Type to be a factor variable with character names and filter the rows to include only the tableware and headlamps type and Fe less than 0.2.

```

#Read in the glass data to a tibble
glass_data <- read_csv("https://www4.stat.ncsu.edu/online/datasets/glass.data", col_names=c(

```

```

Rows: 214 Columns: 11

```

```

-- Column specification -----

```

```

Delimiter: ","

```

```

dbl (11): Id, RI, Na, Mg, Al, Si, K, Ca, Ba, Fe, Type

```

i Use ``spec()`` to retrieve the full column specification for this data.
i Specify the column types or set ``show_col_types = FALSE`` to quiet this message.

```
glass_data
```

```
# A tibble: 214 x 11
```

	Id	RI	Na	Mg	Al	Si	K	Ca	Ba	Fe	Type
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>
1	1	1.52	13.6	4.49	1.1	71.8	0.06	8.75	0	0	1
2	2	1.52	13.9	3.6	1.36	72.7	0.48	7.83	0	0	1
3	3	1.52	13.5	3.55	1.54	73.0	0.39	7.78	0	0	1
4	4	1.52	13.2	3.69	1.29	72.6	0.57	8.22	0	0	1
5	5	1.52	13.3	3.62	1.24	73.1	0.55	8.07	0	0	1
6	6	1.52	12.8	3.61	1.62	73.0	0.64	8.07	0	0.26	1
7	7	1.52	13.3	3.6	1.14	73.1	0.58	8.17	0	0	1
8	8	1.52	13.2	3.61	1.05	73.2	0.57	8.24	0	0	1
9	9	1.52	14.0	3.58	1.37	72.1	0.56	8.3	0	0	1
10	10	1.52	13	3.6	1.36	73.0	0.57	8.4	0	0.11	1

```
# i 204 more rows
```

```
#Change the variable Type to be a factor and filter the rows
```

```
glass_data <- glass_data |>
  mutate(Type = factor(Type, levels=c(1,2,3,4,5,6,7), labels= c('building_windows_float_proce
  filter(Fe<0.2, Type %in% c('tableware' , 'headlamps'))
glass_data
```

```
# A tibble: 38 x 11
```

	Id	RI	Na	Mg	Al	Si	K	Ca	Ba	Fe	Type
	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<dbl>	<fct>
1	177	1.52	14	2.39	1.56	72.4	0	9.57	0	0	tableware
2	178	1.52	13.8	2.41	1.19	72.8	0	9.77	0	0	tableware
3	179	1.52	14.5	2.24	1.62	72.4	0	9.26	0	0	tableware
4	180	1.52	14.1	2.19	1.66	72.7	0	9.32	0	0	tableware
5	181	1.51	14.4	1.74	1.54	74.6	0	7.59	0	0	tableware
6	182	1.52	15.0	0.78	1.74	72.5	0	9.95	0	0	tableware
7	183	1.52	14.2	0	2.09	72.7	0	10.9	0	0	tableware
8	184	1.52	14.6	0	0.56	73.5	0	11.2	0	0	tableware
9	185	1.51	17.4	0	0.34	75.4	0	6.65	0	0	tableware
10	186	1.51	13.7	3.2	1.81	72.8	1.76	5.43	1.19	0	headlamps

```
# i 28 more rows
```

Yeast Data